On April 2nd she had stopped taking the glands, and the urine was increasing in quantity.

I have had great difficulty in inducing the patient to continue the treatment with regularity, and to give me opportunity for accurate observations. I feel that this case has been very insufficiently watched and reported, but I have done the best I could under the circumstances.

I was led to try the effect of the adrenals by remembering the large nerve supply accompanying their blood vessels, which might or might not influence the nerves governing the blood supply of the kidneys, both being derived from the semilunar ganglia. Whether the effect is thus produced, or whether it is the result of the tonic effect alleged to be induced by the ingestion of the adrenal substance on the blood vessels of the system generally, I cannot say, but in the case before me I find that:

1. The urine has been reduced in amount from 4 gallons to 3 pints in the twenty-four hours.

2. The general health and strength have been much improved. The patient, who before could hardly move about the house, can now walk several miles, can do all the work of the house, and attend to her children.

3. So long as she takes the glands the urine is moderate in quantity, but when she ceases taking them the urine increases in amount, and great thirst and weakness result.

What also strikes me is the fact that though the amount of urine passed has so much diminished, the quantity of urea excreted should remain so small. Though so much stronger her weight has not increased. I may also mention that the glands from frozen mutton appeared to have much less effect than those from mutton recently killed.

A CASE OF DISSEMINATED SARCOMA, WITH HÆMORRHAGES INTO THE SKIN AND ELSEWHERE.¹

By T. C. RAILTON, M.D.LOND., M.R.C.P.,
Physician to the Manchester Clinical Hospital.

HISTORY.

J. W., a boy 12 months old, was brought to the Manchester Clinical Hospital in March, 1894, on account of certain soft projections which had appeared upon his head in the course of the previous two months, accompanied by increasing weakness and wasting. He only lived ten days after admission.

He was born naturally and of healthy parents. He was the second child, the other being alive and well. It was not thought that he had received any injury. The projections on the forehead were the first symptom of anything being wrong, and they had made their appearance gradually and painlessly, one or two first showing themselves and continuing to grow steadily, while others followed in the neighbourhood. One of these, prominently situated upon the left side of the forehead, had attained the size of a hen's egg, being surrounded by smaller ones. Another had appeared upon the right side a little more than a fortnight before he was brought to the hospital and this ultimately became the largest. While these swellings were appearing the child deteriorated perceptibly; he became pale, lost flesh and at the time of his admission he weighed 22½ lbs., having weighed 23 lbs. when five months old.

When he was admitted, he presented an appearance of the most profound anæmia, his complexion was waxen, his lips and conjunctivæ bloodless. There were traces of hæmorrhage in the skin of the forehead in various stages of absorption, a bluish ecchymosis in the right upper eyelid towards the inner canthus, and two smaller ones upon the inner and outer sides of the left upper lid. He did not appear to be in pain, but lay quietly in bed with his eyes apathetically looking into vacancy. The numerous bosses upon the anterior part of the head and forehead were soft and fluctuating, the skin over and around them was tinged with blue from blood extravasation. Over the right frontal eminence there was a rounded fluctuating projection nearly 2 inches in diameter, which yielded chocolate-coloured semi-fluid blood when explored with a

needle. Three smaller swellings of similar character were placed behind it. On the left side there were several swellings of moderate size over the frontal and parietal bones. None of the projections were tender to pressure. Blue veins stood out in the middle and at the sides of the forehead. The patient had beads on his ribs and the radii slightly enlarged at the wrists. He had six teeth. The gums were pale but otherwise normal. The bridge of the nose was somewhat depressed. The liver appeared considerably enlarged, and felt decidedly harder than usual. The spleen could be felt a little below the costal border.

There was no evidence of effusion of blood about the limbs nor of deep-seated swelling around the shafts of any of the long bones. He moved freely and showed no tenderness. The urine contained no blood during the time he was under observation. There were no purpuric spots on either body or limbs, nor was there hæmorrhage from any mucous membrane.

Three days after admission a fresh effusion of blood was observed upon the right temple. On the fifth day proptosis of both eyes occurred, the right eye being the more prominent. On the seventh day a fresh swelling made its appearance on the left side of the frontal bone near the suture. The following day the temperature, which had previously been hovering between 100° and 101°, suddenly rose to 104.2°, but without any fresh physical signs. The chest showed no evidence of disease; there was no alteration of the pulse-respiration ratio, and no diarrhoea. On the ninth day the right eye appeared considerably more prominent, and the ecchymosis in the upper lid much deeper in colour. The patient moved both hands, occasionally bringing the right one up to his ear and then putting it down by his side again. The temperature rose to 105.2° F. There was a slight trace of albumen in the urine. The following day he died, the temperature before death being 105.8° F. He had no sickness; the pupils were equal and not unduly dilated. No symptoms of paralysis were observed.

At the *post-mortem* examination made the following day the scalp was found to strip off readily from the pericranium. The tumours beneath the latter consisted of a soft, excessively vascular growth, and were seen under the microscope to be composed of small round cells, evidently of a sarcomatous nature. Between the dura mater and the bone in the anterior part of the skull there was a large, diffuse, lobulated swelling. The bone was roughened both externally and internally at the seat of the growths. The chief mass of the internal growth was at the left side of the frontal region, but

¹ Read before the Pathological Society of Manchester.

it extended also to the right side, and also backwards as far as the occipital region. The brain was compressed by the growth, but otherwise normal.

The liver was not found to be very much enlarged, but there was a large, rounded, encapsulated tumour intimately adherent to the under surface of the right lobe, with its outer surface continuous with that of the liver. This growth measured 4 inches from before backwards, and $3\frac{1}{2}$ inches from side to side at its broadest part. Upon the upper surface of the liver there were three nodules showing a white, cheesy-looking substance when cut into.

The lungs were congested, but there was no consolidation nor evidence of any new growth. There were no hæmorrhages in the pleura. The heart, kidneys, and spleen were apparently healthy. There were some enlarged glands at the roots of the lungs which when cut open showed a soft pigmented substance, and the retroperitoneal glands at the root of the mesentery formed a large lobulated mass presenting the same appearance on section.

The stomach and intestines appeared healthy and neither growths nor hæmorrhages were to be found in them. One long bone—the right femur—was removed for examination. This was selected at hazard, as there was neither swelling nor bruise about any of the limbs. On cutting, however, through the periosteum of this bone, blood extravasation was found beneath in two places, around the upper end of the shaft below the trochanters, and around the lower end above the condyles. The periosteum was not thickened.

Microscopic examination showed the sarcomatous growth within the skull to be infiltrating the dura mater, though not extending to the inner surface. Extensive hæmorrhages had taken place between the cells in both internal and external tumours. Sections of the nodules from the upper surface of the liver showed a similar small-celled growth, the liver substance being shrunken and attenuated in their midst into thin bands and certain parts existed in which no trace of secreting cells was to be seen. Sections from the retroperitoneal glands and bronchial glands showed similar collections of round cells interspersed with the remains of hæmorrhages. The kidneys and lungs showed no abnormality.

REMARKS.

The subject of disseminated sarcoma associated with cutaneous and other hæmorrhages was dealt with by the late Dr. Hilton Fagge in *Guy's Hospital Reports* for 1881, his paper containing an account of six cases. Further cases have since been reported by Drs. Kingsford, Harris, Anderson, and others. It is the relation between the growths and the hæmorrhages which renders the matter peculiarly interesting. Dr. Fagge pointed out the fact that a large proportion of the fatal cases of purpura hæmorrhagica occurring at Guy's during a period of twenty years were of this nature. In some of his cases, notably his first case, some of the purpuric spots were raised, forming indurated nodules, which, when microscopically examined, were found to consist of new growths, probably originating as he suggested, in minute emboli from a primary tumour. In a most interesting communication published in the *BRITISH MEDICAL JOURNAL* by Mr. Jonathan Hutchinson in 1888, a picture of the leg of a gentleman is shown with an eruption upon it of more than a hundred sarcomatous nodules, possibly due to embolism from a growth in connection with the lumbar vertebrae.

This view, however, as Dr. Fagge admitted, does not exhaust the subject. The profound alteration in the composition of the blood which obtains in disseminated sarcoma may have a considerable part to play in the production of the hæmorrhages in mucous or cutaneous surfaces. The ecchymoses in the eyelids and about the forehead of the patient I have described were not raised, and forcibly reminded me of the cutaneous hæmorrhages one sees in scurvy, congenital occlusion of the bile duct, and other diseases of a like character.

The subperiosteal hæmorrhage around the shaft of the femur is to be noted in this connection, bringing as it does the disease, with its intense anæmia and hæmorrhages into the skin, into close relation with infantile scurvy in its clinical aspect.

OBSERVATIONS ON THE TREATMENT OF EPILEPSY.¹

By T. SYDNEY SHORT, M.D.,

Assistant-Physician to the General Hospital, and Visiting Physician to the City Infirmary, Birmingham.

FIFTY cases were selected from the female wards of the Epileptic Department of the Birmingham City Infirmary. Of the male cases under my care, so many work in the grounds that I thought perhaps the exact number of fits that occurred amongst them might not be so accurately recorded. All the women work indoors, either in the wards or in the laundry, and are therefore more closely under observation. These 50 cases were chosen haphazard, in order to avoid the fallacy arising from special selection. Their ages varied from 16 to 70 years. They were all subject to true epileptic convulsions with loss of consciousness. Many of them also, as one would expect, suffered from the minor attacks which occur in epilepsy. The fits recorded in this paper refer to attacks in which the loss of consciousness and struggling were sufficient to attract the attention of a second person, so that they could be entered in the book as a fit. Of the fifty patients originally chosen, a few passed out of observation by reason of removal from the infirmary, or on account of some other unforeseen cause; but I was able to keep a full and complete record of 43, and these form the basis of my observations.

For about twelve months—the actual period being 348 days—the ordinary routine treatment of the cases was continued. The diets varied; most of the patients were taking 3 oz. of cooked meat a day, and a few took as much as 5 oz. Medicinally a mixture containing $7\frac{1}{2}$ gr. each of bromide of potassium and bromide of ammonium was given according to the case and the severity of the attacks; and in the event of numerous fits occurring within a short period of time other drugs were given, such as those contained in the following mixture: 10 gr. of bromide of potassium, 15 gr. of chloral hydrate, in 1 oz. of chloroform water. This period then formed a fair example of the life of these people, such as many of them had been living for years.

During the 348 days the total number of fits recorded for the 43 cases was 5,001, or, in other words, 14.37 fits occurred on an average amongst the whole number for each day. This is not the average number of each individual case, but the average number amongst all the cases together. I have put the average in this way because if the number of fits experienced by each person were taken as an average, the number which I must presently compare would be expressed as fractions of unity instead of whole numbers, and numbers below unity are more difficult to compare rapidly. Books were kept in the rooms or wards in which the women worked, and daily the number of fits were recorded by the nurse on duty. This number, then, 14.37, may be taken as the average number of fits occurring daily amongst the 43 cases during a period of twelve months.

The first change made was this: For eleven weeks—the actual time being seventy-nine days—all the cases were put upon a uniform diet, which I call "A" diet, containing: Meat, 3 oz.; bread, 18 oz.; potatoes, 12 oz.; butter, 1 oz.; tea, 2 pints; gruel, 1 pint; and for medicine they were ordered a mixture of soda and gentian. It happened occasionally that during the occurrence of a succession of severe fits a mixture containing bromide of potassium was given for a short period, usually from one to three days. The main difference, then, between this period and the last was that the diet was made uniform, and the bromide was only given occasionally, so occasionally indeed that only a few had it at all. It is interesting to note that during these seventy-nine days the whole number of fits was 1,119, or 14.16 a day; a record which is practically the same as the average taken as a standard.

The amount of meat in the diet was now decreased during a period of six weeks, the actual number of days being forty-three; all the patients being again on a uniform diet. This diet I call "B" diet: Meat, 2 oz.; bread, 14 oz.; potatoes,

¹ Abstract of a paper read before the Birmingham and Midland Counties Branch of the British Medical Association.