A castration wound was also tested when the drainage tube was taken out forty-eight hours after the operation. Although the result was septic, the wound healed by first intention.

A fishing-gut suture was taken on the seventh day from a wound on the back of the hand, after the removal of a ganglion. The broth in which it had been placed was found by subcultures to contain staphylococcus epidermidis albus. The wound had healed by first intention. At the operation the patient's skin and the ligature were aseptic, but the skin of the house-surgeon, sister, and nurse were all septic.

On another occasion a limb wound was septic. A nævus was removed from the inner side of the knee of a child aged 8 years. When dressed on the seventh day it infected an agar-agar tube. The wound was soundly healed, but the skin around was raised in vesicles containing clear serous fluid.

Lastly, a wound of the neck, made for the removal of cancerous glands, was also septic. At the operation the patient's skin was septic, but that of the surgeon, together with fishing-gut ligature, after having been used, and wool sponge were all aseptic.

A wound for the removal of a cancerous breast and axillary lymphatic glands was dressed at the end of forty-eight hours for the removal of a drainage tube inserted to let out blood, which continued to ooze. The clot in the drainage tube and serum, which ran from the aperture were both septic. The depths of this wound healed perfectly, but owing to the tension needed to bring the enormous wound together the skin sloughed a very little in places, and a minute quantity of pus formed about the sloughs.

These are all the wounds tested with culture media. They were taken as they came without selection. They comprise 22 operation wounds, of which 10 were aseptic and 12 were septic. This is, I believe, a higher proportion of a sepsis than has yet been recorded. Of the septic cases 1 had slight suppuration. This proportion of 1 in 22 marks about the proportion of suppuration which I have of late had in hospital practice in cases in which there was no pre-existing septic focus.

As the house-surgeons are often changing, and likewise the nurses who prepare the sponges, bowls, and towels, this proportion cannot be easily diminished. With such a small proportion of suppuration it is particularly galling to have to report that one occurred after the wiring of a fractured patella. The operation was done because it was clear that the prepatellar aponeurosis had got betwixt the fragments. The patient, too, was anxious to be operated on, although the risk was clearly explained. The suppuration was very subacute, as is usually the case when asepsis is so nearly attained. The patient was never ill or in much pain, and she promises to have a movable joint. The patella itself is united and movable from side to side. At the operation the skin of the surgeon, of the house-surgeon, and of the patient were aseptic; the skin of the nurse was septic. During the operation the 1 in 4,000 biniodide of mercury lotion was aseptic after an hour and a half's exposure to air, and also the same lotion mixed with blood clot. An inexperienced sister introduced an undisinfected bowl into the field of operation before we were aware of the error. No iodoform was dusted on the skin. Also it may seem wise after the event, to remark that before the operation the patient's temperature was raised and rather irregular, suggesting auto-inoculation. But my own belief is that some bleeding took place into the joint and wound, and thus the clot became infected with bacteria of inconsiderable pathogenic properties, but enough, however, to disintegrate the clot. On another occasion I shall drain and use iodoform.

AN OUTBREAK OF OPHTHALMIA IN A POOR-LAW SCHOOL.

By SYDNEY STEPHENSON, M.B., F.R.C.S. EDIN.,

Ophthalmic Surgeon to the North-Eastern Hospital for Children; Surgeon to the Ophthalmic School, Hanwell, W.

THE following account of an outbreak of ophthalmia in the Central London District School is published for two reasons: first, because it emphasises the value of prompt isolation; secondly, because it deals with an epidemic that was practically under my care from the outset.

The essential facts are these. At about midday on Sunday, April 12th, 1896, an infant boy, named W. N., was found by a nurse to have some discharge in the corner of his eyes. Two hours later another child (R. H.) showed some redness of the conjunctiva. Both children belonged to the same ward (D6). They were kept apart pending the arrival of my colleague, Dr. S. G. Litteljohn, who saw them shortly afterwards. Under his direction they were at once taken to the

school infirmary, where they were put to bed and carefully watched.

Both patients were seen by me on the following morning (April 13th). They then appeared to be suffering from ordinary phlyctenular conjunctivitis, but, in accordance with my routine practice, cover-glass preparations were made from the eye discharges. The specimens obtained from R. H. were found to contain a few of the short bacilli described by Koch, Weeks, Kartulis, Morax, and myself, as characteristic of acute muco-purulent or catarrhal ophthalmia ("blight," "cold in the eyes," or "pink eyes," to use the vulgar phraseology). This patient was therefore sent off forthwith to the Ophthalmic School, which is close to the Central London District School. No bacilli having been discovered in the specimens taken from W. N., that lad was kept in the infirmary until the precise nature of his ailment declared itself.

On the following day (April 14th) W. N.'s condition showed no improvement. His eyes were bloodshot; phlyctenulae were present around each cornea; and there was a notable amount of yellowish-white discharge. Bacteriological examination of the latter, moreover, disclosed the existence of numerous clumps of short bacilli, which took ordinary basic aniline colours, but did not retain the stain by Gram's method. A diagnosis of acute muco-purulent ophthalmia was accordingly made, and the lad sent to the Ophthalmic School.

Upon inspecting the other children belonging to the dormitory from which the foregoing patients had come, I found three lads (D. S., C. C., and S. N.) with more or less conjunctival discharge; these were at once isolated. A fourth boy (E. M.) from another ward, who was also found to have muco-purulent ophthalmia, was sent without loss of time to the Ophthalmic School. It was ascertained upon inquiry that this patient (E. M.) had two days previously gone with a message to the infected ward (D6). He had, furthermore, access to a playing yard in common with the inmates of that ward, so that two distinct exposures of E. M. to infection were at once revealed.

Now the dormitory (D6) whence all but one of the original patients had come, had been remodelled so recently as 1890, when the Hanwell School, formerly consisting of an unbroken three-storeyed frontage of 600 feet, was broken up into six blocks by means of air spaces, each 16 feet wide, driven through the building. Its length was 54.75 feet, its breadth 18 feet, and its height 12.25 feet; its floor space consequently amounted to 985.5 square feet. On a basis of 36 feet super, it was capable of accommodating 27 inmates, but as it contained at the time of the outbreak 24 children only, overcrowding could in no sense be said to exist. The ward was ventilated on the south and east sides by six large external windows, and upon the north side by three others. Its windows were never wholly closed, either by day or by night; those upon its north side opened into a covered corridor, which had a length of 77.25 feet, a width of 10 feet, and a height of 12.25 feet, with open ends. The dormitory was further provided with eleven large Tobin's tubes, some of which came direct from the open air, while others ventilated into the corridor, and in all cases they were so arranged that it was well nigh impossible to block them. The room also contained three inlets for warmed air, together with a similar number of extracting flues in connection with three of Teale's open fire grates. Its floor was stained and polished. There was, in short, no reason for thinking that the ward was in anything but a satisfactory sanitary condition; there was certainly nothing in its state to account for the generation (if I may employ so unscientific a word) of muco-purulent ophthalmia.

It should be explained that the children inhabiting D6 were classed as "baby boys"—that is to say, male infants whose ages ranged from 3 to 5 years. In common with the so-called "baby girls"—who occupied a ward upon the same floor as D6—they used a dining and playroom apart from the elder children. In wet weather only, when both "baby boys" and "baby girls" went to a large dayroom for purposes of recreation, did they mix to any extent with others. They had a separate school. A lavatory, arranged upon what is known as the "jet system," was set aside for them, and it goes almost without saying that each child was provided with a towel of his own. It will thus be seen that the "babies" of D6 and D5 formed a small community living apart from the older

infants and girls who were lodged upon the same side of the school premises.

The disease having declared itself so definitely among a particular group of inmates, it was deemed advisable to quarantine those not yet affected. Hence, school attendance was done away with for the time being, and the children sent into the fields as much as possible; a lavatory was appropriated to their exclusive use, and towels were changed upon each occasion of washing. In a word, the inmates of D6 were not suffered to come into any sort of contact with the rest of the school.

The attendants were warned as to the nature of the disease, and told to notify me immediately the least sign of discharge or redness was noticed in any eye. A special observation ward was got ready in the Ophthalmic School, to which any suspicious case might be sent without an instant's delay. Directions were given that the bedding, towels, etc., of any infected child should go to the steam disinfecter, and also that his bedstead should be carbolised. Lastly, the "babies" were examined by me twice a day.

It is most instructive to trace the further course of the malady. On April 15th 3 and on April 16th 6 more cases were detected among the children of D6. On the latter day another lad (J. R.) was taken from the infirmary, to which he had been sent twenty-four hours previously for some general ailment. On April 17th 3 and on April 18th 2 other children were taken from D6. On April 19th the 2 remaining infants were found to have contracted the disorder, and were forthwith separated. In fact ophthalmia had made a clean sweep of the infants of the ward, for the only inmates who remained unaffected were two schoolgirls who helped in the dormitory work. Presumably they escaped owing to the fact that at their age they had naturally less tendency to contract disease than the infants, who from their habits are particularly prone to spread infection from one to the other. Furthermore, one older lad (E. M.) had been separated. The total thus far affected was therefore 23.

On April 19th a suspicious case (E. W.) was found among the "baby girls," who, as already stated, occupied a ward on the same floor as D6. The child was sent to the Ophthalmic School. On the following day another "baby girl" (J. G.) developed the disease, and was isolated without loss of time.

The ward occupied by these "baby girls" was known as D5. It was, save in size, similar in all essential points to D6, and hence need not be described in detail, further than to say that its length was 67.25 feet, its width 18 feet, and its height 12.25 feet. At the time of these occurrences it contained 30 children, of whom 28 were infants and 2 ward girls. Upon the 36 feet super basis, then, there was no overcrowding.

As I feared that the disease might spread among the children of this dormitory, as it had among those of D6, the whole of the "baby girls," with their beds and bedding, were sent to a detached building that lay upwards of 100 yards from the main school, where they were strictly quarantined. The two ward girls went with them, as did also the two who had remained unaffected in D6.

The wards thus vacated by the removal of children were fumigated with sulphur and then thrown freely open to the air. Their walls and ceilings were washed and re-distempered. The floors were scrubbed, and afterwards polished with beeswax and turpentine. An opportunity was also taken of disinfecting, cleansing, and redecorating the attached lavatory.

Judged in the light of after-events, the entire separation of those belonging to D5 had a happy result. Thus, they were kept apart from April 20th to April 30th, yet during those ten days no fresh case was found among them.

We may sum up the facts as follows: Between April 12th and April 25th, 25 children became affected with acute mucopurulent ophthalmia. Of that number, 22 came directly or indirectly from a single ward, and the other 3 from dormitories more or less closely connected with it.

In conclusion, I think it important to state clearly that this is the only outbreak of acute ophthalmia that has taken place in the Central London District School during the last six years. That the recent cases were not of a serious nature will be evident from the facts (a) that those affected are now

convalescent: (b) that in no single instance was the cornea involved; and (c) that no sequelæ have developed.

[Besides the foregoing, 10 girls and 2 boys were sent from the school to the observation ward between April 18th and April 24th, 1896. These children were removed simply as a measure of precaution. Most proved to be instances of phlyctenular conjunctivitis or of episcleritis, but a few were of anomalous nature. Not a single one suffered from mucopurulent ophthalmia.]

NOTES ON A CASE OF ACROMEGALY.

BY

R. ROXBURGH, M.B., and A. J. COLLIS, M.B.,
F.R.C.S. Edin., M.A. Cantab.,
Weston-super-Mare.

THE aggregation of clinical phenomena, which received from Marie the designation of "acromegaly," though constituting a well-recognised pathological entity, still belongs to the rarities of medical observation, and as *post-mortem* examinations of cases have been rarer still, the following example may be of interest.

Mrs. M., aged 35, lived in Scotland till just before her marriage at the age of 19, and from then till the time of her death at Weston-super-Mare. There is no history of a similar complaint in the family, and the family history is good. The mother is living and healthy, aged 60; the father died in India (cause unknown). The patient had had no previous illness of any importance except variola at the age of 13. She had six children in nine years, all of whom are alive and healthy. The last was born seven years ago, and from that time (when aged 28) menstruation ceased entirely.

Six or seven years ago she observed that her hands and feet were getting larger, and five times over she required to have her rings filed off her fingers and made of an increased size. She had to have boots specially made to order during the last three years. Her friends noticed a great alteration in her appearance, and say that her nose grew bigger. No photograph of her as a younger woman is obtainable. She seems to have first come under medical observation in May, 1893, when she complained of failing sight. Double optic neuritis was then found, but no cause was assigned. The size of the hands and feet does not seem to have been such as to attract attention at that time.

The case first came to our notice in December, 1894, when the appearances were those generally associated with this malady—large and ponderous hands and feet, heavy and thickened features, the nose especially so, prominent frontal eminences, the skin of the face and neck coarse and thick. Her weight was 15 stone, though she was not corpulent; two years previously she weighed a little over 13 stone. Her general appearance was anæmic. The enlargement was clearly due to bony growth and not to any subcutaneous deposit. The lower jaw was noticeably thickened, giving the face the ovoid form with the broad diameter downwards which is characteristic. The heart was normal, and the thyroid and spleen presented no increase of size. Constipation was troublesome, and she complained of indigestion and frequently of great thirst; double optic atrophy was present. Owing to various circumstances little was seen of the case till July, 1895, when it was found that the amaurosis had advanced to a condition of total blindness in the left eye and well-marked hemianopia in the right, the nasal half of the field of vision being defective. She complained at this time of occasional headaches and pains in the lower limbs. Towards the end of July she fell into a very drowsy condition lasting about a fortnight, so that she was unable to do her work. One day she was found asleep on the stairs, and when aroused said she had gone to sleep unintentionally whilst going up. She frequently went to sleep whilst being spoken to. This period of drowsiness was followed by a persistent headache, which remained till her death on September 21st. When this headache first came on, it was very severe and prevented her sleeping; she took to her bed, and lay groaning day and night unless relieved by morphine injections. Sulphonal, phenacetin, potassium iodide, arsenic, and cannabis indica gave no relief. During the last week in August