experimentation. When a number of facts seem to indicate that two or more phenomena are correlated and due to the action of a certain cause, we feel generally, when dealing with biological problems, that we may have overlooked many factors, and therefore we have to test our views by experimentation. Experimentation is not, however, always guided by direct observation, for it often happens that the causation of certain lesions is inferred from what we know of the causation of other more or less analogous lesions.

The nature, the duration, and the difficulty of many of the experiments which have been made for the purpose of testing theories touching the causation of disease and the production of lesions make it difficult to teach the whole of this part of pathology by means of demonstrations and practical classes, and it is in this case that systematic lectures or textbooks are the most justifiable and, in fact, necessary.

Etiology and pathogenesis cannot be put aside, for much of the advances which have been made in the art of prognosis, in the treatment, and in the prevention of disease are based on our improved knowledge of these subjects.

Bacteriology, however, lends itself specially well to practical work, and I believe it would be beneficial to most medical men to have a more practical knowledge of this subject than they are generally made to acquire in the course of their studies.

The conclusion which I draw from all these considerations is that students would obtain a more useful knowledge of medicine if they had fewer lectures and more practical courses. They should be made to attend thorough practical courses on (1) pathological anatomy, histology, and chemistry; (2) bacteriology as applied to the study of infectious diseases; (3) general clinical medicine and surgery, with special courses in special branches of clinical work such as diseases of women and obstetrics, diseases of children, infectious diseases, mental diseases, diseases of the eye, of the ear, of the throat, etc.

Systematic lectures should be confined to courses on (1) etiology of disease and pathogenesis; (2) general considerations regarding the practice of medicine and of surgery. These courses should be as short as is compatible with a clear exposition of those principles which but very few students would be able to formulate for themselves from the study of facts.

Finally, I wish to say that if, in expressing my views, I have wounded the feelings of any who have opinions differing from mine I will ask him to apply to me the words of Harvey: "I would not charge with wilful falsehood anyone who was sincerely anxious for truth, nor lay to anyone's door as a crime that he had fallen into error."¹²

AN ADDRESS

DELIVERED AT THE OPENING OF

THE SECTION OF OPHTHALMOLOGY.

By DAVID LITTLE, M.D., F.R.C.S.E.,

Lecturer on Ophthalmology, Owens College; Senior Surgeon, Royal Eye Hospital, Manchester.

At the Annual Meeting of the British Medical Association at Carlisle, July, 1896.

OBSERVATIONS ON THE TREATMENT OF CERTAIN FORMS OF CATARACT.

MY first duty, gentlemen, on taking this chair, is to tender my thanks to the Council of the Association for the great honour that has been conferred upon me by asking me to preside over this Section. However much I may feel the compliment that has been paid me, I am nevertheless very sensible of the great responsibility attached to it. To preside over a meeting of my colleagues is no small undertaking; but I confidently rely on your kind indulgence and help to make the work of our Section a success in spite of all my deficiencies.

The Council of this Association did a wise thing in establishing, about fifteen years ago, a Section of Oph-

thalmology as part of its annual programme. The success attending these meetings has fully justified its action, for no one can doubt the large attendances at most of them, and the interest and value of the papers and discussions on ophthalmology. And how could it be otherwise? The science of ophthalmology has advanced to such an extent during the last twenty-five or thirty years that its work is admitted to be of the highest value, not only for its own sake, but for the beneficial influence which its aims and methods exercise over the whole field of medicine and surgery.

One cannot, therefore, wonder that within the last dozen years or so a very large addition has been made to workers in this special branch of practice; eye hospitals or ophthalmic departments in connection with general hospitals have become established all over the country; in fact, most of the populous towns and districts are well provided with special men, very capable of dealing with all eye disease, and so the public, and particularly the poor, are spared the trouble and expense of going long distances to the great centres for advice, as was the case in former years.

I am an advocate for well-equipped ophthalmic departments in general hospitals, if for no other reason than for the convenience of students. It is hardly possible to get these young men to attend an eye hospital, no matter how great the inducements there offered for instruction, because it is usually placed at some inconvenient distance from their medical school or teaching hospital, and they have neither the time nor the inclination to go out of their way; at least I have found this the case in Manchester. To be able to get all his clinical instruction under one roof is of the greatest advantage. Besides, in addition to the student's point of view, you are well aware that many eye affections are closely associated with some nervous disease, and it often happens that the ophthalmic surgeon desires the opinion of a physician, and a physician that of the ophthalmic surgeon, and these aids are not at hand except in the general hospital.

Speaking of the education of the student, it is very properly becoming more and more the practice of teachers of practical subjects at medical schools to impart instruction by clinical demonstration and less by systematic lectures. Of all the branches of our profession, none demands this line of treatment more than our own. It is hopeless to give a student a notion of eye diseases by lectures alone, and so I have always been in the habit of devoting a part of my lecture hour at Owens College to demonstration on living subjects, as illustrating the main points of the lecture.

In view of the important papers before us, it is not my intention to occupy your time by a lengthened address, but I should like, with your permission, to refer very briefly to some personal experience in one or two practical subjects not touched upon in our programme, and, first, as regards the treatment of immature senile cataract. For some important information on this subject I would refer you to Mr. Anderson Critchett's Presidential Address at Leeds in 1889, to a discussion at the Ophthalmological Society in its *Transactions* for 1890, and to Mr. McHardy's contribution in *King's College Hospital Reports*, vol. i, 1893-94.

When incipient cataract has advanced to a stage when vision is so impaired as to render the patient useless for any occupation, it is clearly our duty to relieve him if we have reasonable means of doing so. To be kept waiting in a semi-blind condition for long and indefinite periods—perhaps years—is a great hardship to those people in many ways and is very depressing and often injurious to health. There is abundant evidence that such cases can be safely dealt with by operation. The question appears to be what is the best operation to perform, whether to extract the immature cataract at once or mature it before extraction by iridectomy and trituration of the lens, as recommended by Forster. The objection, I think, to the former method is the liability to retention of cortex and its consequences. In old people cortex left in the pupil is slow of absorption, and during that tedious process, iritis is apt to occur, sometimes very severely.

With a view to avoid irritation from retained cortex, Mr. Tweedy ruptures the capsule at the periphery after making the iridectomy, so that if cortex should be left behind, it is retained within the capsule and is likely to be harmless. The main objection some surgeons have to trituration of the

¹² Harvey, *An Anatomical Dissertation upon the Movement of the Heart and Blood in Animals*. Dedication 1628. Rendered into English for G. Moreton. Canterbury. 1894.

lens is the fear the operation may damage the eye by undue irritation and iritis and synechia, a condition of things which might militate against the success of the after-extraction.

I have had a fairly good experience in both operations, sufficient, I think, to enable me to form an impartial judgment on their respective merits. Up to five years ago I was in the habit of extracting unripe cataracts in necessitous cases every now and again by the combined method in the ordinary way, with this exception—to avoid as much as possible leaving cortex in the pupil, I made the section unusually long so as to admit an easy escape of the lens with the least possible amount of pressure. In some cases the result was all that could be desired, in others again the cortex found afterwards in the pupil often gave trouble and anxiety by iritis. Without troubling you with any statistics I may just say that the results on the whole were not equal to those of extraction of a naturally mature lens.

For the last five years I have adopted Forster's method in about 40 cases in hospital and private practice. To give it a fair trial I made a selection of what I thought were really suitable cases, following Forster's recommendation as closely as possible. They were cases of uncomplicated senile cataract with soft cortex, in which cortical opacity more or less existed.

Forster considers it essential that cortical opacity must have begun. The vision of these eyes varied from 10 to 20 Jaeger, and the progress of the opacity was known to have been exceedingly slow, extending in some cases over some years. After making a small iridectomy well back in the sclera, I massaged or rubbed the centre of the cornea with a vulcanite or tortoise-shell spoon for about two minutes, exercising care and judgment in the amount of pressure. Beyond a little hæmorrhage occasionally in the anterior chamber, nothing unusual happened in the operation. In no case did undue irritation follow; nothing approaching iritis—the opacity was complete in most cases within a fortnight, in a few it was nearly a month; no atropine was used before operation, and seldom afterwards; the after-extractions were all normal, no escape of vitreous, the healing process went on favourably, there was no loss of the eye, and the ultimate results were on a par with extractions of a naturally mature lens. I have been much impressed with the safety and efficiency of the operation. If it be done with reasonable care and judgment, I feel sure it is as safe as any operation can be, and I shall continue to practise it and recommend it with perfect confidence. There are cases, however, in which I doubt if it would answer. Cataracts of deep amber colour, uniformly hard throughout and no soft cortex, with a good deal of transparency, sufficient sometimes to enable one to see the details of the fundus—these cases, in my experience, can be safely extracted at once, and with excellent results, and I have therefore never tried Forster's method in that class of case. I think trituration might possibly cause undue irritation, or, what is more likely, it might fail to complete the opacity.

In zonular cataract, and in some cases of traumatic or soft cataract in young subjects, it is a favourite practice of mine and has been so for many years, to extract the lens with a Teale's suction curette, in preference to linear extraction. I can rely upon it to clear a pupil more effectually than by linear extraction, and I have often been surprised it is not more generally used. It is said there is a danger of introducing septic matter through the tube, and of sucking iris and vitreous, but if the curette is made clean and aseptic, and kept within the area of the dilated pupil and not passed too deeply, accidents of that kind need not be feared. If I have reason to suspect a rupture of the hyaloid, I do not use it. In zonular cataract I have rarely found excision of a small portion of the pupillary margin of the iris improve vision, at least to any appreciable extent, and for many years I have given it up. I either leave the case alone, or resort to discission and suction. In cases of tough capsule obstructing vision after cataract operations, if the needle or double needle operation fails to cut it or displace it permanently, I employ a Wecker's scissors, and have found it answer admirably, better than a cataract knife, or cutting needle, with the additional advantage that there will be no dragging of the iris when adhesions are present.

Wecker's scissors is also best adapted in making a pupil when it is partially closed and drawn up from iritis after extraction.

We sometimes see an eye with a white irregular mass of shrivelled up calcareous-looking capsule and lens, filling up the whole pupil, and presenting rather an unsightly appearance, usually the result of some old injury. These people consult one for cosmetic reasons, and ask if it cannot be removed. I have operated upon several such cases in my time, with good cosmetic effect. I make the ordinary cataract incision with iridectomy above; if there are iritic adhesions I separate them; very often there are none. I then introduce a stout pair of capsule forceps (specially made for the purpose, ribbed on the inner side of each blade), and by gentle traction draw the white mass out; sometimes I have had to pass one blade behind and the other in front to get a firm hold. Sometimes vitreous escapes, sometimes not. It may not be bad practice to leave such cases alone, in case of inflammation and greater disfigurement. In the few cases I have done nothing of the kind has happened. If the iris is healthy looking and the tension normal I have no hesitation in recommending operation.

In traumatic prolapse of the iris the best practice I have found is to slit up the sac throughout the whole length of the wound with a Graefe's knife, under antiseptics, and if iris remains in the lips of the wound to excise it. Rapid healing takes place, and the result is a firm and flat cicatrix, with little or no synechia, and the eye is left pretty safe from future infection. Such good results are not by any means always obtained by the other recognised methods, as excision, bandage, etc. I have never found eserine or atropine reduce a prolapse, even when it has been very small and recent.

Errors of refraction form a large proportion of ophthalmic work nowadays. We have various ways of estimating refraction errors. In my own practice I rely on retinoscopy and testing the vision with Snellen's test types. My object in referring to this subject is to say that I paralyse the accommodation with atropine in a large proportion of young subjects in the first examination. I do not mean in every case, but in a large number of them, and it is astonishing how great the spasm that is often found to exist.

By paralysing the ciliary muscle we ascertain the full amount of error, and are able to correct the defect most accurately. These young patients, on their first visit, are often nervous and unreliable, and atropine gives one independence; besides, the examination can be made more easily and rapidly, and time is saved.

The Camberwell Guardians have, on the motion of Dr. Partridge, appointed Miss Mary Helen Cruikshank, L.R.C.S. and P. Edin., and L.F.P.S. Glasg., to the post of third assistant medical officer at Constance Road Workhouse.

A discussion was recently initiated at the London School Board by Mrs. Holman, who proposed that a committee should report on the advisability of engaging a fully-qualified female practitioner as assistant to Dr. Smith, the Board's medical officer. The proposal was rejected by a substantial majority, some members arguing that an assistant was not required and others holding that a lady doctor was out of place.

MEDICAL MAGISTRATE.—Surgeon-Lieutenant-Colonel Robert Anderson (retired) has been named a Justice of the Peace for the County of Suffolk. The new magistrate was gazetted from Netley to the 3rd King's Own Hussars, and served with that regiment through the whole of its Indian tour. He was on the special escort of the Prince of Wales when His Royal Highness was in the Bombay Presidency and he was also on the staff of His Excellency the Commander-in-Chief of the Bombay Army. Surgeon-Lieutenant-Colonel Anderson served in the Egyptian campaign, and has the decoration of the Khedive star and medal.

HOSPITAL SUNDAY FUND.—A meeting of the Council of the Hospital Sunday Fund was held at the Mansion House on July 28th. The report, which was adopted, showed that the total amount available for distribution was £44,029, of which £41,829 was recommended for distribution among 125 hospitals and 55 dispensaries. £2,200 was set apart for the purchase of surgical appliances during the ensuing year.