with its central putrid patch, unless further absorption were at once stopped.

I commenced the operation with the intention of merely resecting a rib and establishing a free drainage of the pleura, and I am bound to admit that there seemed little hope of a successful issue to the case.

It was not till I had seen and felt the condition of the lung that it occurred to me that the chances of recovery would be greatly augmented by further surgical interference. I was led to drain the lung by a consideration of the extent to which Nature might be aided by artificial means; in short, I treated the infiltrated septic lung as I would have treated septic infiltration of the cellular tissue over the buttock—by free drainage.

It is possible that some will consider the patient's recovery due rather to good luck than to good surgery. I am aware that my treatment of the lung was attended with great risk. Infection of the pleura had, however, already occurred, and had there been hæmorrhage I have no doubt that it could have been arrested. Hæmorrhage from the lung has several times been successfully treated by plugging, and in the centre of an area of solid exudation it could not surely have been very difficult.

Since this case came under my care Mr. Paget² has put before us the records of many cases of operations on the lung, and a careful consideration of all the cases recorded convinces me that the only right treatment of this case was that which I adopted.

ON THE SUBSEQUENT HISTORY OF CHILDREN WHO HAVE SUFFERED FROM TYPHLOITIS.

By R. W. MURRAY, F.R.C.S.,
Surgeon to the Liverpool Infirmary for Children.

THE cases I am about to deal with have all been treated at the Children's Infirmary during the last nine years. Some were under the care of the late Mr. Rhinall¹ Pughe, most of them under the care of my colleagues (Drs. Peter Davidson and Marsh), and others under my own care; and I take this opportunity of thanking Dr. Davidson and Dr. Marsh for allowing me to make use of their cases, for I have thus been enabled to collect a series of 23 cases of an affection not very uncommon in children, but an affection which I take it is of interest both from a medical and from a surgical point of view. These 23 patients by no means represent all the children suffering from typhloitis treated at the infirmary since 1886, but they are all the cases of which I have notes, and whose subsequent history I have been able to trace.

With one exception I have purposely refrained from including the more recent cases, that is to say, those under treatment during the last two years, so that one may the more fairly deal with the question of prognosis. In order to satisfy myself as to the after-history of the patients, I have recently called and seen them at their own homes. In the majority of instances I saw the children themselves, and when not enabled to do so either from their being at work or at school, I learnt from the mother the child's present state of health and progress since discharged from hospital.

The cases for the most part naturally arrange themselves into the three following classes:

CLASS I.—Inflammation of the vermiform appendix, most probable associated with a localised peritonitis.

CLASS II.—A localised peritoneal abscess associated with an inflamed appendix.

CLASS III.—The appendix rapidly become gangrenous or perforated, resulting in general suppurative peritonitis.

CLASS I.—INFLAMED VERMIFORM APPENDIX.

The ages of these children varied from five years to twelve years, the average age being nine years. The history was usually the following: For a week or ten days before admission the child complained of pain in the abdomen, and from time to time vomited. The bowels, as a rule, were constipated, though in some instances the motions had been loose and offensive. The vomiting continued off and on, and there was loss of appetite and increase of abdominal pain. The

legs, especially the right, were drawn up, and any attempt to straighten it caused pain. The tongue was furred, and the temperature ranged from 99° to 102°. There was no very marked abdominal distension, but acute localised tenderness in the right iliac region. The treatment almost invariably adopted was, rest in bed, milk diet, fomentations to the right iliac fossa, and an enema or dose of castor oil. The symptoms usually subsided rapidly, and at the end of a few weeks all tenderness had entirely disappeared and the child was again in its usual health. In 6 of these 9 cases there was no previous history of any similar trouble, but 3 had suffered from similar attacks, and in fact might be called "relapsing cases." A boy of 11 years—the relapses occurred in the older children—was in hospital three times for the same complaint, the last attack being far the most severe. A girl of 12 years was in hospital twice for the same trouble; and in the case of a boy of 10 years there was a history of several milder but similar attacks previous to the one for which he was admitted. In every one of these 9 cases there was an accompanying gastro-intestinal disturbance, and the local trouble in the right iliac fossa was, I take it, almost certainly due in the first instance to a catarrhal condition of the vermiform appendix, the swelling of its mucous membrane causing some retention of the morbid secretion, which induced inflammation of the appendix as a whole, together with, most probably, a slight and strictly localised peritonitis. All these 9 children have remained in good health since their discharge from hospital, and, as this includes a period ranging from three to seven years, the results must be extremely gratifying to the physicians under whose care they were.

The following case, recently under the care of Dr. Davidson, illustrates the occasional difficulty in diagnosis.

Edward P., aged 6 years, was admitted into the infirmary on October 5th, 1895. His only previous illness was measles, two years ago. For a fortnight before admission he complained of pain in the abdomen, and lost his appetite. His bowels were regular, and he had not vomited. The abdominal pain had occasionally been so severe as to "double him up." When I saw the boy, with Dr. Davidson, three days after his admission, during which time he had been kept in bed, he appeared to be in good health, and had had no return of the abdominal pain. His temperature, which on admission was 100°, had fallen to and remained normal. His bowels were constipated, but his tongue was clean. There was no abdominal distension or tenderness, but in the right iliac fossa a firm irregular-shaped mass was noticed by Dr. Davidson the first time he saw the child. In order to make a more thorough examination chloroform was administered. The swelling could then be readily felt. It was about the size of a hen's egg, irregular in outline, remarkably hard, and apparently fixed to the bone. What appeared to be a thickened cæcum and ascending colon could be detected attached to the upper limit of the tumour. A rectal examination confirmed the hardness and fixity of the mass. It was difficult to say what the nature of the tumour was, and in the absence of a previous history of trouble in this region, and on account of its hardness and the absence of tenderness, it was thought unlikely to be of an inflammatory origin. I inclined to the view that we were dealing with a new growth. However, the diagnosis being uncertain, an exploratory operation was performed on October 16th, and I found not a new growth but the vermiform appendix much thickened and enlarged, bound down to the iliac fossa by a great mass of firm fibrous tissue. I liberated the appendix, and removed it by performing practically a circular amputation near its base. The boy subsequently made an uninterrupted recovery.

CLASS II.—LOCALISED PERITONEAL ABSCESS.

The clinical histories of the 10 patients in this class were much the same as in the 9 non-suppurative cases, with this difference, either the inflammatory trouble was from the first more acute or the children were not brought for treatment sufficiently early. There was, as in the simpler cases, some preceding and accompanying gastro-intestinal disturbance. The children, as a rule, were decidedly ill, the temperature raised and the tongue furred. The right thigh was acutely flexed upon the abdomen, there was swelling, tenderness, and marked induration in the right iliac fossa extending sometimes from Poupart's ligament to the umbilicus, with, in the more advanced cases, some local oedema of the skin.

It is rare to find actual fluctuation, and certainly most undesirable to attempt to do so, for fear of rupturing the abscess and so causing general suppurative peritonitis. This localised suppuration is, of course, merely an aggravation of the more simple inflammatory trouble, and I do not draw any very hard and fast lines between the two conditions. The indications upon which I chiefly rely as evidence of suppuration are, the general condition of the patient and the local condition in the right iliac fossa. If there is suppuration the temperature as a rule is raised, but by no means necessarily so, for in several

² *Surgery of the Chest.* By Stephen Paget, F.R.C.S. Bristol: J. Wright and Co.

instances in which I have evacuated one or two ounces of pus, the temperature has been very little above normal.

Much more usually if there is suppuration, the general condition of the patient either remains stationary or deteriorates. The local trouble becomes aggravated, and the swelling and induration as distinct from mere tenderness more pronounced. However, it is by the combination of symptoms rather than upon any individual symptom that one must be guided; always remembering that we may be dealing with a localised collection of pus in the immediate neighbourhood of the general peritoneal cavity, which, if not evacuated, may at any moment open into that cavity, and cause probably a fatal issue.

At the same time it is well not to be in too great a hurry to open these abscesses. As Mr. Treves has pointed out, they have fortunately a tendency to travel towards the surface, and if an attempt be made to open them before the induration is pronounced the pus, if present, can only be reached by opening the general peritoneal cavity, and must subsequently be drained through that cavity. I have never made an exploratory puncture with a needle in order to ascertain the presence of pus, for I consider the practice a most dangerous one. Important structures may thus easily be wounded, and unless the needle be a large one the pus is usually so thick that the result of the exploration, so far as the presence or absence of pus is concerned, would probably be negative. One of the chief determining factors of suppuration in this region is probably the condition of the vermiform appendix. A healthy gut will not allow the passage of micro-organisms through its walls, but with an inflamed gut it may be otherwise. And the facility with which micro-organisms may and do pass through the walls of the vermiform appendix appears to bear some relation to the intensity of the inflammation of the appendix. In this respect I much regret that no bacteriological examination of the pus was made in any of my cases, but Mr. Treves, to whom we owe so much in this and other branches of surgery, in relating before the Harveian Society a series of cases of relapsing typhilitis, said:

It has been now apparently demonstrated that that form of localised peritonitis.....is due, in the great majority of instances, to the bacterium coli commune. Thus Tavel and Lanz examined twenty-three cases of trouble in the appendix. The examination was made either of the pus within or without the appendix or of the exudation into the peritoneal cavity immediately around the diseased process. The material for this bacteriological investigation was obtained during the patient's life. In three no micro-organisms were found. In nineteen examples the colon bacillus was found either alone or in association with other pathogenic germs. The streptococcus was found in several cases; the pneumococcus was met with in two examples, but not in pure culture. A staphylococcus was in one instance the only micro-organism discovered.

In the 10 suppurative cases an incision was made over the centre of the area of induration, usually about an inch above and parallel with the outer one-third of Poupart's ligament, simply with a view of opening the abscess. If the pus was sweet a drainage tube was inserted without irrigating; if, as more frequently happened, the pus was offensive, gentle irrigation was practised. In none of these 10 cases was any attempt made, either at the time of operation or subsequently, to search for the vermiform appendix. The abscess was simply opened and drained. The general treatment adopted was the same as in the simple cases.

In 9 out of the 10 cases the children rapidly improved. The drainage tube was usually discarded during the second week, and the wound then soon healed. The patients were sent home wearing a Thomas's hip splint on the affected side in order to keep the parts at rest. These 9 children were not only discharged as cured, but have remained well ever since. The time that has elapsed since the operation varying from three to six years, a result as gratifying to me as it was astonishing. Their ages at the time of operation were from 3 to 12 years, the average age being 8 years. One of these 10 cases died, and it teaches an obvious moral.

M. M., 4½ years of age, was taken into the infirmary on July 6th, 1892. Four days before admission she had complained of pain in the abdomen and limped. On admission she was a well-nourished child, but walked with a considerable limp and great pain. The right thigh was flexed, almost to a right angle, and any attempt to straighten it increased the pain. The hip-joint was healthy, but there was acute tenderness, with some swelling and induration in the right iliac fossa. The tongue was furred and the temperature 99.2°. In spite of rest in bed and general treatment the swelling and induration increased, and on July 17th (eleven days after her admission) an incision was made over the centre of the area of induration, and 4 ounces of thick offensive pus evacuated. The abscess cavity was gently irrigated and drained.

She was doing well when on the morning of August 4th (eighteen days after the operation) during the syringing of the wound (it would appear

that the syringing was somewhat forcible), the child suddenly complained of pain in the abdomen, and became collapsed. She frequently vomited, and died twenty-eight hours afterwards with a greatly distended abdomen. I did not see her after the accident until a few hours before death, when it was too late to do anything. At the examination after death we found both cæcum and appendix bound down by adhesions. A peritoneal abscess localised to this region had been ruptured opening into the peritoneal cavity, causing general suppurative peritonitis. The mucous surfaces of the lower end of the small intestine, the cæcum and appendix, presented an unhealthy catarrhal condition, but the appendix, which was evidently the centre of the mischief, contained no foreign body, nor was there any perforation.

CLASS III.—GANGRENE AND PERFORATION.

In 3 cases the vermiform appendix rapidly became gangrenous or perforated. These three children died.

CASE A.—A girl, aged 8 years, of whom I regret to say I have a very scanty record, was admitted on January 15th, 1886, and died on February 11th, 1886. She had hip disease, a tumour of the cerebellum, and pneumonia, but the immediate cause of death was some acute abdominal trouble; after death a pin was found to have perforated the vermiform appendix.

CASE B.—A boy, aged 5 years; is the only case in the whole series in which the onset of the trouble was apparently due to an injury. He was admitted on the evening of August 2nd, 1889, and died the following day. He had not been well since he had measles six months before admission; at the same time he had no very definite symptoms. Five days before his admission he fell while running, and afterwards complained of pain in the abdomen. When admitted he did not appear to be very ill, but there was great pain and tenderness in the right iliac region, and he kept his right thigh drawn up. The temperature was 99°. During the night he screamed with pain after passing water, but a sound did not detect a stone. The following morning the child was very much worse. There was distinct swelling in the right iliac fossa, and the tenderness had also increased. The acute pain after micturition persisted. During the evening he became suddenly collapsed, the abdomen distended and tympanitic, and he died in a few hours. At the *post-mortem* examination we found general peritonitis. The vermiform appendix, which was adherent to the back of the bladder, was gangrenous, perforated at its extremity, and lying just outside this perforation was a concretion the size of a pea. Whenever the bladder was emptied it pulled upon the inflamed appendix, thus accounting for the acute pain after micturition. I would draw attention to the fact that this child died in a few hours after the general peritoneal cavity became affected, suggesting death from some virulent poison rather than inflammatory trouble alone.

CASE C.—A girl, aged 5 years, was admitted under Dr. Marsh on the evening of October 7th, 1893. She was quite well on October 4th, but on October 5th—two days before admission—she suddenly complained of pain in the abdomen and vomited. The vomiting continuing and the abdomen becoming more swollen and acutely tender, the child was brought to the infirmary. There was no history of an injury. On admission, she had a pained and anxious expression as she lay on her back with her legs drawn up, and complained of pain in her abdomen, which was swollen and very tender. The respirations were hurried and entirely thoracic; the temperature was 100° F. Next morning she was much worse, and had vomited everything she had taken since admission. As the child was evidently suffering from acute and suppurative peritonitis, the cause being doubtful, it was decided to explore the abdomen. This was done at mid-day. General suppurative peritonitis was found, the coils of small intestine being acutely inflamed and adherent one to the other by layers of lymph. Several adherent coils were separated, thus liberating some clear serous fluid. It was not considered advisable to make any prolonged search for the seat of origin of the peritonitis. The abdomen was irrigated with warm water, a drainage tube inserted, and the wound closed. Death took place two hours and a-half later. At the *post-mortem* examination no perforation of any portion of the gut was discovered, but the vermiform appendix was coiled up upon itself, acutely inflamed, and fixed by adhesions of recent lymph; its extremity was of a dark colour and almost gangrenous. A most careful examination was made for a perforation of the diseased process, but none was discovered.

THE HEALTH OF THE SPANISH TROOPS IN CUBA.—The *Times* Special Correspondent, writing from Havana says: "Sickness amongst the troops, and especially yellow fever, has increased notably of late. In the Military Hospital (Havana) there are this week 2,500 patients, none of these being convalescents. On reaching the convalescent stage the men are immediately sent to La Cabana, there to recover sufficiently before being returned as fit for duty. From all sections of the island the health reports are equally discouraging. The brother of General Ahumada has died of yellow fever in Havana, and the deaths of many other officers from this cause are reported from different places. The worst of it is that for two months to come the sickly season will continue."

TYPHOID FEVER AT BARKING.—An outbreak of typhoid fever has occurred at Barking. Dr. C. F. Fenton informs us that from July 13th to August 31st 23 cases were notified; 13 of these cases were removed to the Upney Isolation Hospital, and among these 4 deaths have taken place. The statement which has appeared in some of the newspapers to the effect that the Upney Isolation Hospital is full is, we learn, incorrect; the hospital is not and has not at any time been full.