prevent any dragging upon the skin at these points. As the main portion of the threads lie over and at right angles to the line of union, they may be utilised for keeping in position the deeper dressing, which may be passed beneath them, but this is not an essential feature of the method.

A CASE OF HODGKIN'S DISEASE.

By WILLIAM E. L. ALLEN, M.B., C.M.GLAS.,
Hawkshead, near Ambleside.

A. E., aged three years and a-half, previously healthy, with the exception of a mild attack of scarlet fever followed by measles in October, 1895, was first seen on March 7th, 1896. She was suffering from a scratch by a cat on the ring finger of the right hand, which was very much inflamed and suppurating. About a week later an abscess formed in the right axilla, which when opened discharged a dark brown offensive fluid. This healed under ordinary treatment, but was accompanied by some sloughing. About this time some glands below the left ear became slightly enlarged, but the swelling disappeared on painting with weak iodine liniment.

About the middle of April the cervical and submaxillary glands, axillary and inguinal glands were found to be swollen on both sides. Under iodide of potash internally and iodine externally there was considerable diminution in the size of the cervical glands on the right side, and subsequently slight improvement in those of the left side. None of the glands exceeded the size of an almond, many not being larger than a pea. It was impossible to count them, so many being involved, but all were freely movable—distinct and free from pain. The complexion became very waxy and there was marked leukæmia, the spleen was easily felt about an inch below the ribs, about the size of an orange. There was also considerable enlargement of the liver. The face and body were very much swollen, but there was no pitting on pressure and the swelling was more a puffiness than actual œdema. The bowels and kidneys acted freely. The temperature was slightly above normal, but never more than 102° F., shortly before death. At the beginning of May there was epistaxis, but not severe, though it continued more or less the whole of one day. There were very marked purpuric spots on the arms, legs and lower part of the body, the largest being about the size of a threepenny piece. On May 6th pneumonia set in, which proved fatal on May 10th. Four days before death there was suppuration from the right ear.

This case is interesting owing to the early age of the patient and to its rapid termination, as there was only a period of nine weeks from the first distinct appearance of the complaint—dating back to the first enlargement of the cervical glands—before death took place. It is also interesting on account of the suppuration from the ear, which in this case was a sequel of the complaint and not a cause of it as suggested by Trousseau. Though there is a history of a poisoned (?) wound, it can scarcely be looked upon as a case of septicæmia, but rather as the cause of the scratch taking on an unhealthy action, as the symptoms point conclusively in favour of Hodgkin's disease.

As to treatment, arsenic and phosphorus have been used, but without any marked benefit except in Billroth's case, where arsenic was used. In this case the glands diminished under pot. iodid., but that has been observed under other forms of treatment, shortly before death and apparently independent of treatment. In a future case I should be inclined to give the gland treatment a trial.

A CASE OF HEAT STROKE: HYPERPYREXIA: RECOVERY.

By GERALD SICKEL, SURGEON, R.N.

[Communicated by the DIRECTOR GENERAL, Medical Department, R.N.]

THE following interesting case occurred on board H.M.S. *Gibraltar* after she had been in the Mediterranean just over a month:

On July 13th, when at Palma, in Majorca, at 6 P.M., J. W., an engine-room artificer, aged 27, came down to the sick-bay,

complaining of headache and pains in the legs. His temperature was 100.4°, his tongue slightly furred, he looked pale and ill and was put in a cot in the sick-bay.

He lay there sleeping until 9 P.M. when the sick-bay steward thought he seemed slightly delirious and took his temperature which was 105.2°, shortly afterwards his breathing became stertorous, and he lay comatose with a temperature 109.2°, full rapid pulse, pale face, closed eyes, pupils equal and fixed about half-way between dilatation and contraction: his venous system was engorged, his body dry and felt burning hot: at times he groaned loudly and struggled to get out of bed.

He was bled to 6 oz., but it apparently did him no good, and his breathing becoming rather embarrassed he was given a hypodermic injection of 5 minims of liquor strychninæ; meanwhile a bath was being prepared, into which he was put, the temperature of the water being brought by ice from tepid to about 65° in half an hour. This brought his temperature down to 101.2°, otherwise his condition remained unaltered. At 11.30 P.M., he was given a large soap enema, which emptied the lower bowel of a great deal of feces, in part scybalous. Later his temperature rose to 103.4°, at 12.30, 104.6° and he began to have epileptiform fits, in one of which he bit his tongue: later he had fits every 10 minutes.

At 2 A.M. on July 14th the temperature was 107.8°. He was put in a cold pack and his body rubbed with ice, the temperature came down, till at 5 A.M. it was normal, with pulse 104, respirations 22, he began to grow a little conscious and recognised being called by name. He was sick several times during the night. Later he became fairly conscious and his temperature did not go above 99°.

On July 15th the temperature kept down, but he was sick after everything he ate; he had a very restless night; morphine gr. $\frac{1}{4}$, with atropine gr. $\frac{1}{15}$ did no good. Urine: acid, specific gravity 1024, no albumen.

July 16th, the temperature was 99° and the pulse 76 in the morning. Vomiting had ceased, but he was still very restless; he was not fully conscious but complained of diplopia and difficulty of hearing. Towards evening he became more and more restless and in the end he struggled so violently that he had to be tied down in bed. Morphine, gr. $\frac{1}{2}$, and bromide potassium 3ss did no good. During the ensuing night the temperature went up 102.2°: and he was put in a wet pack.

On July 17th, at 1.35 A.M. he was given 30 grs. of chloral hydrate, after which he had a sound sleep. After he woke his mind was wandering, and he passed urine under him twice, the tongue very dry and furred: with sordes in teeth and lips. He was given another dose (20 grains) of chloral, after which he slept all day, having to be waked to take his food (beef tea and eggs beaten up) which he kept down well. At 9 P.M. the temperature 99.2° the pulse 68 and the respirations 20.

On July 18th, he had had a very good night, but required another 20 grains of chloral in the evening.

From this date he steadily improved until on the 22nd he was allowed to sit up for two hours in the evening.

Until July 19th, he had a decided impediment in his speech, but this then almost entirely disappeared—he had some slight peculiarity of speech before the stroke—and this of course remained. From this date onwards he made an uninterrupted recovery.

An ophthalmoscopic examination was made on July 23rd, there was no optic neuritis, or atrophy, or retinitis.

COMMENTS.—There was nothing very peculiar about the atmospheric conditions on the day of the stroke. The barometer at 4 P.M. was 30.12 in., thermometer 87° in the shade and relative humidity (100-saturation) 52: but we had just come from Malta, where everyone appeared to get "run down." The thermometer night or day hardly ever fell below 80°. From 7 A.M. until 3 P.M. he had been working in the engine-room where the temperature was 100°. About the same time that this man was ill, we had several minor cases of heat stroke, the majority of them among those who worked in the engineers' department.

It is interesting to notice that, as is so often the case, the stroke came on in the comparatively cool evening.

Sir Joseph Fayrer objects to bleeding in these cases as a rule, but in this man the nervous system was so obviously engorged,

that it appeared advisable to relieve it by moderate bleeding and quite possibly although only 6 oz. were abstracted, and although there was no immediate good result this step was important towards his recovery, for it immediately relieved the pressure on the nervous centres, if only to a very small extent, while it was not enough to retard his convalescence by unduly weakening him. Temperatures over 108° are usually fatal, and the present case is one more in evidence of the efficacy of the direct application of cold to the body in cases of hyperpyrexia. He was given the bath as soon as it could be got ready.

For the restlessness and delirium, both morphine and bromide failed, but chloral always had the desired effect: in my experience a somewhat surprising fact after the two former drugs have failed.

The relapse on July 16th was by no means unexpected, and here again cold in the shape of a wet pack acted well.

The impairment of speech, sight, and hearing after the first attack are all evidence of the great strain the entire central nervous system must have sustained.

In testing his urine, which was done on the earliest occasion possible, it is to be regretted that it was not possible to test for sugar, as the reagents had been destroyed by the warm weather: it would have been interesting to observe if temporary glycosuria, due to intense central nervous disturbance, were present.

A SUCCESSFUL CASE OF DRAINING THE LUNG.

By HAROLD B. MEAKIN, M.D.LOND.

THE following case¹ came under my care in the London County Asylum at Banstead:

E. C., aged 31, a very stout, well-fed woman, working in the laundry of the asylum, developed an attack of acute pneumonia at the base of the right lung on May 20th, 1896. The patient had been in the asylum for eleven years, and was subject to frequent and severe epileptic fits. With the exception of an attack of influenza in 1892 she had been in good health since her admission.

On May 22nd the patch of consolidation showed no increase. It was about 5 inches in diameter. There was a resonant interval between the edge of the dull area and the spine, and there were no signs of pleural effusion. Her evening temperature was 103.5° .

On the evening of May 23rd the patient appeared to be more distressed, and her breath was noticed to have a pungently foetid odour. An examination showed that the area of dulness had extended in every direction, and there was no resonant interval between it and the spine. The temperature was 104° . Later in the night the patient's condition was distinctly worse, and her breathing was short, laboured, and very rapid. The dulness had reached the clavicle, and though harsh breath sounds were heard everywhere, the character of the percussion note was that met with when an effusion is present. The dulness extended one inch to the left of the sternum and the heart's apex beat was outside the nipple line. The temperature was 104.5° . The patient was sweating freely, and was obviously very ill. She had some cough, but could not be persuaded to expectorate the sputum.

The chest was punctured with an exploring syringe in the sixth space in the posterior axillary line. The syringe filled easily with an opalescent fluid, with the peculiar odour which is associated with gangrene. The needle of an aspirator was then introduced, and about 3jss of thick and deeply blood stained fluid were withdrawn, which was even more stinking than the fluid removed by the exploring syringe. As the cannula was slowly withdrawn from the chest there was a sudden rush of fluid, not blood stained, and similar in appearance to that withdrawn by the exploring syringe. Altogether 3viij of fluid were withdrawn and the stream then ceased spontaneously.

The removal of the fluid made no difference to the dulness and seemed to cause but temporary improvement. Four

hours later (6 A.M., May 24th) the patient's condition became very grave. Her chest was again explored, and finally about 3vj of opalescent stinking fluid, not blood stained, was obtained through a cannula introduced high up in the axilla. It seemed certain that she would die unless a free drainage of the pleura were established.

The patient was taken to the theatre, and since her condition did not justify the employment of a general anaesthetic, local anaesthesia was induced over the intended seat of operation by Schleich's method of infiltration. The sixth and seventh ribs were resected in the anterior axillary line, about $1\frac{1}{2}$ inch being removed from each, and the pleural cavity was opened. A spot so far anterior was chosen because the patient's breathing was so laboured that it did not seem safe to turn her further over to the sound side of her chest. This would have been necessary had a spot further back been chosen.

On the pleura being opened, a quantity of greyish-black, semipurulent fluid, with an abominably foetid smell, gushed out. The lung could be clearly seen and felt. It did not collapse, and was never more than $\frac{3}{4}$ inch away from the opening in the pleura. It was bluish-black in colour. To the touch it was generally hard, though immediately opposite the opening was a spot where it felt doughy and inelastic; over this spot was a small punctured wound in the visceral pleura, which, it seemed likely, was due to one of the previous exploratory punctures. Doubtless it was from a puncture of this sort that the thick blood-stained fluid of the first tapping was obtained. The point of a pair of closed pressure forceps was introduced through this, and readily found its way to a depth of about 2 inches. The forceps were withdrawn unopened, and were then pushed in again, having first gripped the end of a piece of india-rubber drainage tube. The drainage tube was left in, reaching into the lung tissue a depth of about 2 inches. A second drainage tube was put into the pleural cavity, and the ends of both tubes were sewn to a transverse piece of tubing to keep them in position. The wound was well dusted with iodoform powder and dressed. The patient was then put back to bed.

After the operation the patient's distress was considerably diminished, and her condition improved materially. About five hours after the operation she developed a status epilepticus. Her temperature was 105.5° , her pulse reached 160, and then became so running that it could not be counted. She had frequent opisthotonos, and was acutely cyanosed. Rectal injections of chloral and bromide of potassium had no apparent effect on the fits. In order to reduce her temperature she was sponged first with tepid water, then with iced water, and finally rubbed with ice. The temperature fell to 104.8° , the fits ceased, and the patient dropped into a heavy sleep.

From this point onward she made a steady recovery. The discharge from the tube in the lung was, for the first 24 hours, dark red, almost black, and of a tarry consistency. It then gradually took on the characteristics of ordinary pus. The tube in the pleura at first drained a fluid similar to that which escaped at the operation when the pleura was first opened. The stench from the wound was foetid for the first four days, after which it disappeared. The tube in the lung was shortened at intervals, and finally removed on June 3rd. The tube in the pleura was finally removed on July 4th. On July 11th the wound was quite healed. The breath-sounds were normal over the whole chest, and the patient, who had been putting on flesh rapidly for a fortnight or so, returned to her work in the laundry.

This case seems to me to be of interest in respect of (first) the diagnosis and (second) the treatment.

The clinical history of the case is best explained by regarding the rapidly-occurring foetid effusion as due to infection of the pleura from a patch of pulmonary gangrene. Gangrene of the lung following pneumonia is comparatively common among the insane. It is unfortunate that the patient swallowed her sputum. An examination of it might have assisted the diagnosis materially.

In regard to the treatment adopted I cannot but think that the patient's recovery depended greatly upon the draining of the lung in addition to the free opening of the pleura. At the time of operation she appeared to be thoroughly poisoned by septic absorption, and it seemed unlikely that she would live long enough for Nature to deal with the extensive pneumonia

¹ I am indebted to Dr. Clay Shaw for permission to publish this case, and for advice and support in the treatment I adopted.