CLINICAL LECTURE ON ANEURYSM OF THE POPLITEAL ARTERY.

Delivered at the Middlesex Hospital

By ANDREW CLARK, F.R.C.S.,
Surgeon to the Hospital.

GENTLEMEN,—The number of cases of aneurysm admitting of surgical treatment that present themselves at the hospital nowadays are few; I therefore think that it is desirable to devote this clinical lecture to the case of A. S., a young man, aged 26, who has recently been discharged convalescent after cure of a popliteal aneurysm by ligature of the femoral artery in Hunter's canal.

It is not common to meet with aneurysm in so young a subject, but in this case the origin seems to have been a strain, since, as we shall see presently, there is a distinct history of his having hurt himself when lifting a side of beef (he is a butcher), and two days after he felt a lump behind his knee. He comes of a healthy family, and he has healthy children. There is no history of syphilis, gout, or any other dyscrasia which might have led to inflammation of arteries so as weaken them at any spot; neither is he old enough for any of those degenerative changes to have taken place which come with age. Ever since he can remember he has always enjoyed good health with the exception of an attack of pleurisy seven years ago and, very recently—about a fortnight ago—a slight attack of inflammation of the bowels, due, probably, to the presence of a large tapeworm, which he got rid of about that time.

The history that he gives is as follows:—About Christmas, 1894, he felt something give way behind his left knee when lifting a heavy side of beef. The pain he experienced did not last long, and he continued his work and forgot all about it; but two days after, feeling a decided pain, he put his hand behind his knee, and distinctly felt there a lump about the size of a marble. He did not attach any importance to it, and said he thought it would go away as it came, but he admitted that it gradually, though decidedly, increased in size. He worked and lived as usual until November 19th last, when he was confined to bed for a few days by an attack of enteritis. He had, however, got over this and was at work again, and went to bed in his usual health on November 25th. He was awake the next morning by an acute pain behind his knee, and on putting his hand there felt that the lump had more than doubled its size, and then thought it wise to call the attention of his medical man (Dr. W. G. Murray, of Clacton-on-Sea) to it, who, recognising the condition, arranged for his speedy removal here for treatment.

On admission we found a man apparently in perfect health in every way with the exception of a pulsating expansile tumour, about the size of a large duck's egg, occupying his left popliteal space, pressure on the femoral artery in Scarpa's triangle completely arrested the pulsation, and it was obvious without further examination that it was an aneurysm of the popliteal artery from which the patient was suffering. There are some cases of aneurysm the diagnosis of which is attended with very great difficulty, cases, for instance, in which there has been inflammation in the tissues round the sac, or where the aneurysm has been partly consolidated, or where it has become diffused; and there are others, again, like the one we are discussing, in which no one, I think, who knew what an aneurysm was could make a mistake. Nevertheless, before deciding on the method of treatment to adopt, it was necessary to go more fully into the case, and I will ask you to notice the following further particulars about it: The knee is habitually slightly bent, although it can be straightened without giving pain; the tumour is soft and elastic, and can be reduced very much in size by manipulation, and if pressure be kept up on the femoral artery for about a minute, it can be made to entirely disappear, reappearing the moment the pressure is removed. On placing a stethoscope over the tumour the purring sound characteristic of aneurysm can be distinctly heard, and an examination of the arteries lower down in the leg, the anterior and posterior tibial on both sides, does not show any difference in the force of the circulation.

In addition, too, it was observed that the presence of the tumour had had no ill effect on other structures from pressure; there was no œdema from pressure on the vein, no pain from pressure on a nerve, and no interference with the cutaneous circulation. With the above symptoms we came to the conclusion that we had an aneurysm in which there had been no attempt at spontaneous cure, and which had not in any way interfered with the circulation of the blood through the limb. The next point in the case to which I will direct your attention is the sudden increase of size after it had existed nearly a year; when first felt, you remember, it was the size of a marble, slowly increased until November 26th, and then suddenly doubled in size. In all probability the aneurysm was of the sacculated variety, which is by far the most common in the limbs and, indeed, everywhere, and it projected from the course of the artery in the way that sacculated aneurysms do; supposing the aneurysm projects from the front of the vessel the dense bone against which it presses—you remember that the artery is close to the femur and tibia—prevents its attaining any great size; if it projects from the side or back, there is nothing to prevent its enlargement, and the pressure from within, having practically no counterpressure from without, soon causes a large tumour; in this case it probably sprang from the front, but near one side, and as soon as it escaped from the pressure of the bone, the much more rapid enlargement ensued. I do not suppose that the tumour really doubled its size in a few hours, as the patient thought; we must always take patients' statements for what they are worth; for, with the best intentions of giving an accurate history, their powers of observation are not sufficiently cultivated to enable them to do so, and I think it exceedingly probable that only when he felt pain he put his hand down. Had the aneurysm ruptured, it would very likely have suddenly increased, but then the effusion of blood would have continued, and, instead of the circumscribed tumour, we should have had a diffuse swelling in all probability presenting far greater difficulties for diagnosis. The only things that a circumscribed aneurysm can be mistaken for are bursæ, circumscribed abscesses, and solid tumours, such as enlarged glands; but, when it has become diffused, it may be mistaken for acute suppuration or a soft malignant growth, the difficulty being greatly enhanced by the absence of pulsation, and there are several cases recorded in which experienced surgeons have made serious mistakes.

This was obviously a case in which surgical treatment was necessary and in which the prognosis was very good, and we had no hesitation in assuring the patient that, if he would help us, he would in all probability get perfectly cured. There were several methods open to us for treatment, any of which were applicable in this particular case. I mean manipulation, flexion, compression—either digital or instrumental—and ligature of the artery above the tumour, either the popliteal or the femoral in Hunter's canal or in Scarpa's triangle; but before resorting to either of these modes of cure I thought it prudent to keep the patient in bed and employ medical treatment, partly with a view to accustoming a man who has been leading an active life to lie in bed, and partly in the hope of assisting the consolidation of the tumour. Accordingly he was put on a mixture containing gr. v pot. iod. three times a day and a low diet—treatment calculated to lessen the blood pressure and diminish the quantity of blood in the body.

I just now mentioned four amongst the various surgical procedures in vogue for the treatment of aneurysm, either of which seemed applicable in this case. The first of these was manipulation, a method introduced about forty years ago by the late Sir W. Fergusson, and practised successfully in cases that were, from their situation, incapable of being treated by more certain means, and has been employed as an adjunct to other methods, even in popliteal aneurysm, and the result has, I believe, proved satisfactory. The idea is that a portion of clot is detached from the consolidating tumour, and, escaping into the blood current at the distal end, blocks up the artery, and thus acts much in the same way as the distal ligature. It must be admitted, however, that this is not devoid of danger, as the clot may be sufficiently small to pass through the larger arteries, and eventually get stopped in one of the cerebral arteries and lead to untoward cerebral

symptoms and even hemiplegia; and, again, the arterial coat being weakened, rupture of the sac is not an impossible accident.

I did not intend seriously to try this method, but it occurred to me that during the ten days that elapsed between the patient's admission and any other surgical treatment, it was possible that consolidation might have taken place from the manipulation to which the tumour was subjected, not only by myself, but by a good many of the students for diagnostic purposes. I was not able, however, to detect any difference in it when on December 10th I decided to commence compression. This sometimes can be effected by the employment of the method known as flexion, introduced into this country by Mr. Ernest Hart about the same time as manipulation, but I believe practised on the Continent some years before. It is a mode of treatment said to be specially applicable to aneurysm in the popliteal space, and can only be used at such places as the ham or the elbow, where one part of a limb can be forcibly bent on the other, and its effect is, not only to obstruct the circulation through the main artery, either completely or nearly, but also to exercise compression on the tumour, and when this method is successful, it cures by so obstructing the circulation that layers of clot form on the inside of the sac until it is quite blocked up. It is very easy of application: the leg is firmly and evenly bandaged up to the knee, and then the heel brought up to the buttock and tied there, but it is very uncomfortable, and after a time the forced unusual position becomes painful, and the pain is frequently so great that the patient cannot endure it long enough for it to be effectual. This was one reason why I did not have recourse to this method of treatment; and the other was because I could not, on bending the limb, sufficiently obstruct the circulation, very likely because the aneurysm was so large, and it is generally said that it is in cases in which the tumour is small that flexion is successful. Digital compression, introduced about 50 years ago, might have been employed; in this case it would have been the femoral artery, either at the base or apex of Scarpa's triangle that would have had to be compressed, and I suppose that is about as easy an artery to compress as any in the body, provided the tips of the fingers press in the right direction, that is, in such a way that the artery may be between them and the bone. Students often forget this in compressing an artery, and dip their fingers deeply into the soft tissues, carrying the artery before them; this certainly controls the blood flow to a certain extent, but is attended with considerable pain to the patient and unnecessary fatigue to the operator. Popliteal aneurysms have been cured by digital compression, but I did not employ it here partly on account of the difficulty of getting relays of fingers sufficiently accustomed to that kind of work to lead one to think that the result would have been satisfactory, and also because of the facility with which instrumental compression can be effected. Compression with tourniquets then was the treatment resolved on, and before speaking of the details of its application, let me tell you why I selected it in place of ligaturing the artery, which in the case of popliteal aneurysm, is recommended by some surgeons as the quicker, more certain, and less painful mode of procedure, and the risk of accidents in ligaturing the femoral artery in such a case is said to be *nil*. My chief reasons were: first, that if the compression were eminently successful, it was possible that consolidation might take place in a few hours, and the average time is less than twenty-four; if it were not successful, as it was not, the patient would not be in any worse position for the ligature to be subsequently applied, as even if there had been some adhesions between artery and vein or artery and its sheath, set up by the compression (and this is one of the reasons suggested against compression), it would not have interfered with the operation, as I did not propose to tie the artery at the same place that the pressure was applied. Rather than worse off I think he would be better off, because the fact of the circulation through the main vessels having been for some time cut off, enlargement of other vessels would have begun, and a collateral circulation would be more readily and more rapidly established. As to compression being more painful, I do not think that is a reason against it, as the pain may be combated by opiates, or an anæsthetic may be employed during the whole time the

pressure is kept up, but the pain is often not excessive, and pressure can be borne by many patients for a considerable time; in this case it was over three hours before the patient complained at all, and through the whole time that the compression was kept up the pain was combated by opiates.

I hardly like to say that the risks attending ligature are *nil*, and I think the risk of gangrene of the limb, which is perhaps the most likely accident and the most serious, is lessened if pressure has been employed. Statistics show that the results of ligature are much better than they used to be, and there seems to be a gradual progressive improvement which is probably due to the improved methods of operating; still, for all that, I think that compression is the safer.

On December 11th, at 9.45 A.M., the patient having been prepared and the leg well powdered, a flannel bandage was applied from the foot to the knee, and Skey's tourniquet arranged on the thigh as high up as possible, Signoroni's horseshoe tourniquet being placed about 8 inches lower down. The upper one was immediately tightened, and without any great pressure, and without giving any pain, completely controlled the circulation. At 10 o'clock the lower one was tightened and the upper one relaxed; this did not control the blood flow until great pressure was exercised, which was attended with a good deal of pain, so the instrument had to be a little altered in position, after which for about a couple of hours the circulation was completely controlled with the two alternately.

Beyond the superficial veins being a good deal distended, the notes tell us that everything went on perfectly satisfactorily up to 12.45, when the pressure exercised by the lower tourniquet produced a good deal of pain, so a mixture containing 15 minims of liq. opii sedat. was prescribed, with orders that it should be repeated at intervals of not less than two hours if required, and a Carte's tourniquet substituted for Signoroni's, with a view to making the pressure a little more elastic. It had the desired effect, and the patient bore it well until half-past 7, when it got shifted, and did not control the blood-flow satisfactorily. At this hour it is noted that the thigh had become oedematous and the surface covered with small petechiæ. Patient said that he had a good deal of pain all the way down the leg, but not more than he can bear.

The lower instrument was rearranged, and pressure continued, the two instruments being tightened and loosened about every fifteen minutes until 10 o'clock, when I saw him; and finding that the oedema had much increased, as well as the petechiæ, and that the whole limb was quite numb, I thought it expedient to remove the pressure altogether; accordingly the tourniquets were removed, and the leg enveloped in cotton wool and bandaged from the foot to the groin, another dose—making the fourth—of the opiate mixture given, and the patient left to go to sleep. There was a good deal of oedematous swelling over the aneurysm, which appeared to have consolidated, and no pulsation could be detected in it. The compression had been practically continuous for a little over twelve hours. About an hour after this it is noted that the patient had not slept owing to pain along the line of the artery, and that there was some pulsation in the tumour.

The next morning (December 12th) it is noted that the patient had a restless night, and his temperature had risen to 99.8° F. The numbness had been gradually passing off, and there was a severe aching pain through the whole of the limb. On examination there was considerable tenderness where the pressure had been applied, and still a good deal of oedema, but it was evidently subsiding; there was obvious pulsation in the aneurysm, but it was harder and apparently partly consolidated. I did not, however, think it advisable to return to the pressure, chiefly on account of the extent of the oedema that had been produced, and decided as soon as the limb had regained its normal condition to ligature the artery leading to the aneurysm.

The question then arose: Where was it to be tied? There was the choice of the common femoral, the superficial in Scarpa's triangle or Hunter's canal, the popliteal from the inner side of the thigh, or in the popliteal space. The last-named is practically the old operation, which is supposed to be superseded by the Hunterian method, chiefly on the ground that the artery is more liable to be diseased so near

the aneurysm. It has, however, been revived within the last dozen years, and practised in cases in which the size of the aneurysm was producing injurious pressure effects, as it enabled the surgeon to remove the tumour at the time of the operation, instead of leaving it to disappear gradually.

Then, again, it has been shown that in many cases where the artery has been tied at a distance from the aneurysm it has remained pervious, and in some, though a temporary cure has been effected, pulsation has returned in the aneurysm, and further treatment has had to be resorted to. This did not seem to me to be a case in which there was any need to get rid of the swelling rapidly, and as laying open the sac would have certainly been a more serious operation than either of the others, I decided against that. As regards ligaturing the common femoral, I at once put that out of the question, as, though it is easily got at, and is a mode of procedure recommended by some experienced surgeons, it does not meet with general sanction, partly, no doubt, because the results have not been shown to be generally satisfactory, and partly because, being a short vessel and having a good many (though small) branches, it is very likely that a firm clot will not form. The choice, then, lay between the superficial femoral (in Scarpa's triangle or in Hunter's canal, and the popliteal on the inner side of the thigh. I chose the second, as being easier to reach than the popliteal, and nearer to the aneurysm than the vessel in Scarpa's triangle. This was the original operation performed by John Hunter, but for a good many years it was abandoned in favour of the operation in Scarpa's triangle, where the artery is more superficial, and said to be more easily got at.

On December 18th, just a week after the compression, the œdema had disappeared, and the general condition of the patient being quite satisfactory, the rise of temperature noted the morning of the 12th not having recurred, I proceeded to the operation of ligature of the femoral artery in Hunter's canal. I need not go into all the details and steps of the operation. The usual antiseptic precautions were adopted, the usual incision made, and the artery exposed without any difficulty; the sheath was carefully opened, and a ligature of Chinese twist used to encircle the vessel. The moment it was tied pulsation quite stopped in the aneurysm, the wound was closed with silk sutures, and a small drainage tube inserted at the lower angle, and iodoform and dry dressing applied. The limb was enveloped in cotton wool, carefully bandaged with a flannel roller, and the patient returned to bed.

The notes the next day say the patient kept awake a good deal during the night by pain along the line of the incision; no pulsation in the tumour; no œdema of the leg, which is quite warm and comfortable. Temperature at 10 A.M. 100°; the drainage tube withdrawn. The same evening the temperature was 100.4°, after which there was a gradual fall till it became normal four days after the operation, where it remained. On the 22nd the wound was looked at and found to have healed by first intention; the stitches were taken out. No pulsation in the tumour and no untoward symptom whatever. On January 1st the patient was allowed to be up a few hours, and a note says "slight pulsation in tumour," so he was kept in bed for four days longer, after which he was up a little every day, and on examining him on January 10th I made a note, "the tumour is firmly consolidated, and now about the size of a Tangerine orange; there is no pulsation in it, neither can pulsation be felt in the tibials below the tumour; the leg is quite warm and apparently in its normal condition, and the patient's general health has not apparently suffered from his stay in the hospital." Discharged January 11th cured; whether the cure will be permanent I cannot say, but I see no reason why it should not be.

RUSSIAN CONGRESS OF SYPHILIS.—A Congress of Russian Syphilographers will be held in St. Petersburg on December 18th. The object of the Congress is to discuss the means of checking the spread of syphilis in Russia. The Medical Department of St. Petersburg has in the meantime appointed a special Committee to examine the official reports sent to the Government from different parts of the Russian Empire, dealing with the subject.

AN ADDRESS

ON

SOME SURPRISES AND MISTAKES.

Delivered at the Surgical Section of the Royal Academy of Medicine in Ireland, on Friday, November 15th, 1896.

By WILLIAM THOMSON,

President of the Royal College of Surgeons in Ireland; Direct Representative for Ireland on the General Medical Council; Surgeon to the Richmond Hospital; etc.

[AFTER some preliminary observations, in the course of which Mr. Thomson expressed his thanks for the honour done him by his election to the Presidency, he continued as follows:—]

I think, as a rule, that most students have in their early days a good deal of trust in the sequence of events. They are enamoured by the newness of all around them in the hospitals; they have plenty of work to do, and those who do it have not time to be sceptical. They take what the books tell them so definitely, as though under no circumstances is it possible that there should be any deviation from the course of things as there described. Later on they have their faith disturbed as experience grows; but yet to the oldest of us there still remains a belief that under certain conditions certain things must happen. There is deep down in our minds the conviction "that when the brains were out a man must die, and there an end," but how that is upset by our experience! There, for example, is the famous case of the man through whose frontal lobes a heavy iron jumper was driven by a blast, and who recovered. I have seen a man who had lost a portion of his occipital bone, as large as the palm of my hand, and a cupful of brain substance, and yet years afterwards he did not seem to be anything the worse. It is true that these, what I may call "accidental experiments," have helped us to be less fearful of interference with the brain, but always with the exactitude which has been helped by deliberate research. Still, for the greater part, these rough tearings and bruising of brain tissue are fatal; and even when with our later and better knowledge they are recovered from, they are "surprises." As our experience grows we encounter them. We—and I include physicians as well as surgeons—see cases which according to all rule ought not to live. The prognosis is a fatal result; and yet they do well, and disappoint our certainties by their recovery. And then there are those cases which die, and we do not really know why. A man gets a scratch which ought not to hurt a child, and he dies none the less surely of shock than if he had suffered some terrible crushing injury of internal organs. What are the curious conditions in the two classes of cases—the powerful vital resistance, as we call it, in the one, which is effective against forces which we regard as unconquerable, and that other class in which vitality is so low and the nervous system so sensitive that it yields at once to what is almost a pretence of injury?

Just let me leave these suggestions for inquiry or speculation, and give you one or two examples of many which have come under my own notice. An old man was sent to me as suffering from appendicitis. He had had an attack undoubtedly, but it had cleared up. The only signs remaining were a slightly raised temperature and a thin strip of dulness extending upwards to the iliac crest. I made an opening, and found a little pus and an extensive granulation-lined cavity. I scraped it out and drained in the usual way. All went well; but in about a week I was startled to hear that there had been a noisy discharge of flatus from the wound and a sign of feces on the dressings. When I saw him there was no escape of gas, but the next day I found on the dressings a faecal concretion quite like a large nutmeg in shape and colour. Of course it had come from the intestine into the abscess cavity, and had perforated the gut wall. But how had this great mass passed through without leaving a large fistulous track, with faecal extravasation and perhaps worse? There was never again any sign of escape of bowel contents, either solid or gaseous, and the patient, having been delivered in this way of his concretion, got quite well without further trouble. It is, I think, plain that, although the final act of extrusion took place when the flatus escaped