unsuitable for a condition like an extrauterine gestation—for such a case, for instance, as the one I have related to you in some detail.

I have not mentioned attempts to kill the foetus by puncture of its sac or by means of electricity, because I consider such methods far more dangerous than an exploratory operation.

A CASE OF SUDDEN DEATH DUE TO EMBOLISM OF ONE OF THE CORONARY ARTERIES OF THE HEART.

By H. D. ROLLESTON, M.A., M.D., F.R.C.P.,

Assistant Physician and Lecturer on Pathology, St. George's Hospital; Assistant Physician to the Victoria Hospital for Children, Chelsea.

FROM a clinical and pathological point of view the coronary arteries are chiefly of interest in connection with angina pectoris and fibroid disease of the heart. In discussing the influence of the coronary circulation on the heart, Cohnheim¹ says that cases are on record where the sole cause of death was found to be embolism of the coronary arteries. That this result is rare is shown by the fact that there is no example recorded in the *Transactions* of the Pathological Society. Embolism of the terminal twigs of the coronary arteries often occurs in pyæmia, giving rise to pyæmic abscesses. Dr. Hebb² has described embolism of the left coronary artery giving rise to myomalacia cordis in a case of ulcerative endocarditis; this condition of anæmic necrosis, due to blocking of the coronary arteries, Osler³ considers to be a common cause of sudden death in medico-legal cases. It is probably much commoner as a result of endarteritis and thrombosis than of embolism, of which the following case is an example:

A boy, aged 17, was admitted to St. George's Hospital on March 4th, and died April 27th, 1896. He was under the care of my colleague, Dr. Penrose, but in his absence from April 14th, until the patient's death, I took charge of the case. He had had two attacks of rheumatic fever—three and two years ago. On admission he was found to be anæmic, short of breath on exertion, and orthopnoic. The heart was dilated and hypertrophied, the apex beat being in the sixth intercostal space, two inches outside the nipple line, and the impulse somewhat diffused. The cardiac dulness began at the fourth rib. There was epigastric pulsation. The heart sounds were short and very rapid, varying during his stay in the hospital between 130 and 175. There was no cardiac murmur. The pulse was small, irregular, and of rather increased tension.

There were signs pointing to pleural effusion on the right side, but paracentesis was performed without any result. There was a cloud of albumen, varying in amount, in the urine. Œdema of the legs, and subsequently ascites and cedema of the scrotum and penis developed. He was treated with cardiac tonics and rest in bed, but his condition in no way improved, and on April 17th, paracentesis abdominis to thirteen pints was performed with considerable relief.

The cause of the condition of dilation and hypertrophied heart was by no means plain, but on the whole it was thought to be subsequent and secondary to rheumatic fever, and to be due to advanced mitral regurgitation; the absence of any characteristic murmur being explained by the extremely dilated condition of the heart. On the other hand, the tension of the pulse, and the absence of any definite signs of organic valvular disease, made it necessary to consider the possibility of the disease being primarily renal and secondarily cardiac. The varying amount of albuminuria, however, was quite compatible with the view that it was merely the result of chronic venous congestion.

The treatment with digitalis, strophanthus, citrate of caffein, and diuretics, in various combinations, being unsuccessful, and the patient's condition becoming steadily worse, it was determined to give a trial to Schott's method. Dr. Bowles very kindly undertook to direct, and indeed himself performed, the manipulations, which were limited to gentle movements of the arms in various directions for periods of quite short duration. He considered that the very serious

condition of the patient did not warrant any treatment by baths. Schott's treatment was begun on April 21st and continued till the end of that week. The patient expressed himself as feeling better, which indeed he looked; his breathing improved, and his pulse diminished somewhat in rapidity.⁴

On April 27th, two days after the last occasion on which Schott's movements had been practised, he was quite suddenly seized with great shortness of breath, and became cyanosed, and in spite of the administration of oxygen and a subcutaneous injection of ether he died ten minutes after the commencement of the attack, apparently from pulmonary embolism.

As it happened, Schott's manipulations had not been performed for forty-eight hours, Sunday intervening, and there does not appear to be any connection between them and the fatal result from coronary embolism. Had the patient lived a few hours longer they would, in the natural course of events, have been repeated, and if embolism had then occurred it might have been thought, in the absence of an necropsy, that they had, by exhausting the already-dilated heart, led to cardiac failure. It is important, therefore, that a careful *post-mortem* examination should be made in any fatal case previously treated by Schott's method, so as to prove whether or not there is any relation between the two.

The *post-mortem* examination performed by Dr. Ogle showed that the pericardium was natural. The heart weighed 28 ounces, and showed hypertrophy and dilatation of both ventricles—especially the left. The auricles were slightly dilated. All the cavities of the heart contained *post-mortem* clots. The valve segments themselves were healthy, but the mitral and tricuspid orifices were enlarged as a result of dilatation of the ventricles, and more especially of the rings from which the auriculo-ventricular valves arise. There was no anæmic necrosis or appearance of an infarct, and both to the naked eye and microscopically the myocardium was healthy. There was an adherent thrombosis near the apex of the left ventricle. The orifices of the coronary arteries and the aorta were healthy, but on slitting open the coronary arteries an *ante-mortem* clot, similar in appearance to that seen in the left ventricle, was found impacted in the left coronary artery, about three-quarters of an inch from its origin, just before the septal branch came off. The vessel wall appeared healthy, except for a few specks of atheroma, and both the distal and proximal parts of the artery were free from any *ante-mortem* clot. This, then, was a case of coronary embolism, occurring—as is said to be most frequently the case—in the left or anterior coronary artery. The heart was shown as a card specimen at the Pathological Society on Tuesday, May 5th, 1896, by Dr. Cyril Ogle.

The right lung was everywhere covered by pleural adhesions, and over the lower lobe, where the visceral layer of the pleura was markedly thickened, there was a localised pleural effusion of small size. The lower lobe showed considerable fibrosis, with some dilatation of the bronchial tubes. There was some obsolete tubercle at the apex. The left lung showed to a slight degree the effects of chronic venous congestion, and contained no tubercle. The lungs were not cedematous. Cohnheim, after experimental ligation of the coronary arteries, never found cedema of the lungs, but Michaelis did, both in dogs and in rabbits.⁵

The left kidney weighed 1 ounce, was occupied by tubercle in a quiescent condition, and contained no secreting substance.

The right kidney weighed 8 ounces, showing considerable compensatory hypertrophy. There were scars of former infarcts and two recent ones. To the naked eye the kidney in other respects appeared healthy. Microscopically there was some increase in fibrous tissue along the lines of the vasa recta and around the glomeruli, but not sufficient in amount to explain the hypertrophy of the left ventricle. The spleen was cardiac, but did not contain any infarcts. The brain and vessels at the base were quite healthy.

Remarks.—That embolism of the coronary arteries has long been recognised as a cause of sudden death, is due rather to

¹ Lectures on General Pathology, vol. i, p. 86, New Sydenham Society.

² Trans. Path. Soc., vol. XXXV, p. 140.

³ Principles and Practice of Medicine, 1895, 2nd edition, p. 676.

⁴ Compare Dr. Bowles, *Practitioner*, vol. lvii, p. 30, July, 1896.

⁵ *Zeitsch. f. klin. Med.*, xxiv, s. 270, quoted in *Twentieth Century Practice of Medicine*, vol. iv, p. 438 (article, Angina Pectoris, by Dr. Whitaker).

the remarkable results of experimental ligature of these vessels in animals than to extended clinical observation. Sir J. E. Erichsen was one of the first to ligature the coronary arteries in animals (*The London Medical Gazette*, 1842, p. 561) taking for his text a speculation of Dr. Marshall Hall's that interruption of the coronary circulation would rapidly cause death. Since then the experiment has been repeated by a number of observers.⁶ The results obtained were usually disturbance of the rhythm, slowing and finally standstill of the heart. They were, however, far from constant, differences being recorded in the time which elapsed before the heart came to a standstill, even when the same species of animal was employed, and as might be expected different animals reacted differently.

Panum⁷ experimentally injected tallow into the coronary arteries, and noticed that the left ventricle ceased to beat 75 minutes and the right ventricle 90 minutes from the time of the injection. According to Cohnheim (*Lectures on General Pathology*, vol. i, p. 35, New Sydenham Society) ligature of any large branch of either coronary artery in a dog produces, after a quiescent interval of 90 seconds, irregularity and slowing of the heart; this is followed not later than two minutes from the ligature of the vessel by sudden stoppage of both ventricles in diastole. After this the heart cannot be roused to renewed contraction. This sudden stoppage he regarded as probably due to the effect produced on the cardiac ganglia by the occlusion of the vessel, but he admits that the distribution of the ganglion cells within the muscular mass of the organ makes it impossible to determine with certainty whether the arterial anæmia affects the nervous or muscular elements. Apart from the view that the cardiac failure is due to failure of nutrition of the cardiac ganglia or muscular tissue, there is the possibility that it is not the mere absence of blood and nutritive material, but the presence of toxic substances generated by the active tissue metabolism which is responsible for the fatal termination. Under normal conditions such toxic bodies are carried away as soon as manufactured, but with arrest of the circulation they might accumulate *in situ* and paralyse the neuro-muscular apparatus of the heart.

The rapid and sudden failure certainly suggests that the cause, whether it be anæmia or a poison, affects primarily the nervous regulation rather than the musculature of the heart, inasmuch as the latter is less sensitive to external influences and has a reserve power which lasts longer than that of the former.

It is noteworthy that while experimental ligature of the coronary arteries in animals or embolism in the human subject may rapidly cause death, gradual occlusion of the vessels as in endarteritis need not have this effect. The gradual diminution in the supply of blood when the trunks or orifices of both coronaries are obstructed would, it may be supposed, allow of some degree of acclimatisation to the altered conditions taking place. When, as is usually the case, one coronary artery only is markedly affected by endarteritis, compensation may be effected by a dilatation of the anastomoses, which Wickham Legg and S. West have shown to take place normally between the two vessels.

A difference between Cohnheim's experiments and this case is the duration of life after the occlusion of the vessel. In Cohnheim's observations on dogs death occurred in two minutes; in this case about ten minutes elapsed. That this is not a point of any great importance or difficulty is shown by the fact that experimentally the results are not quite the same in rabbits as in dogs. If there be a difference between animals in the way they react to experimental ligature or compression of the coronary arteries, a similar variation may well be expected between animals and man.

It does not seem certain what interval elapses between embolism and death in man. Osler (*Principles and Practice of Medicine*, edition ii, p. 676) says, "sudden death not infrequently follows blocking [he is speaking of embolism as well as thrombosis] of one of the branches of the coronary artery and the production of anæmic necrosis." But anæmic necrosis would take some little time to develop, and although in such cases death might be quite sudden it would not be an immediate result of coronary embolism.

⁶ For a summary, vide Porter, on "Ligation of the Coronary Arteries," *Journal of Pathology*, vol. xv, p. 121.

⁷ Virchow's *Archiv.* 1862. Bd. xxv. S. 312.

In a case of embolism of the coronary artery referred to by Dr. Cutler (*Pepper's System of Practical Medicine*, vol. ii, p. 831) death occurred 20 hours after the commencement of symptoms thought to be due to the embolus. It thus differed considerably from my case, in which death followed the onset of symptoms in ten minutes, and where from the absence of any anæmic necrosis in the distribution of the blocked vessel there was no reason to believe that the interval between embolism and death was longer than it appeared to be from clinical evidence.

THE LEFT AURICLE IN MITRAL STENOSIS: HYPERTROPHY AND DILATATION.

By D. W. SAMWAYS, M.D., D.Sc., M.A., B.C.,
Late Fellow of St. John's College, Cambridge; Mentone, France.

PROFESSOR OSLER, in his *Principles and Practice of Medicine*, refers to the changes which take place in the left auricle in cases of mitral stenosis in these words:

"The left auricle discharges its blood with greater difficulty, and in consequence dilates, and its walls reach three or four times their normal thickness." "Eventually the tension is increased in the pulmonary circulation owing to impeded outflow from the veins. To overcome this the right ventricle undergoes dilatation and hypertrophy, and upon this chamber falls the work of equalising the circulation."

A somewhat similar statement occurs in Dr. Fagge's *Medicine*,¹ where one reads, "The effect of mitral stenosis on the heart will naturally be to produce dilatation and hypertrophy of the left auricle."

These statements I believe to be misleading and erroneous, especially as given by Professor Osler, who seems to regard dilatation and hypertrophy as being both useful factors in maintaining the circulation. In the first place, it is probably an error to suppose that dilatation of the left auricle exists to any marked extent in compensated mitral stenosis, and in the second place, if it exists, it is a mechanical error to suppose that such dilatation aids the cardiac chamber to overcome resistance in front. The characteristic alteration in the left auricle in mitral stenosis is not dilatation but hypertrophy, as is proved by the following statistics.

During the four years 1888-91 inclusive there were 70 cases of mitral stenosis examined and reported on in the *post-mortem* room of Guy's Hospital. Hypertrophy of the left auricle is stated to have occurred in 36 of these cases. The 70 cases are divisible into two classes almost numerically equal: those in which the stenosis was severe, one finger (55 millimetres) or less, and those in which the stenosis was less severe; 36 cases belonged to the former class and 34 to the latter.

In 15 of the 36 cases of severe stenosis, the left auricle was hypertrophied without dilatation, the hypertrophy having generally been considerable. In 11 other cases the auricle was both dilated and hypertrophied. In only 3 cases, and in those the orifice measured 5 centimetres at least, was the auricle dilated without being hypertrophied, thus showing a marked contrast to the tendency to hypertrophy. No note was made of the condition of the remaining 7 cases, except in one instance where the auricle was somewhat vaguely stated to have been "a little large."

There was, therefore, hypertrophy of the left auricle observed in 26 of the 36 cases of severe mitral stenosis, dilatation existed only in 14 cases, and was associated with hypertrophy in 11 of these cases.

It must not be forgotten that in *post-mortem* records we have the register, not merely of the cardiac condition in compensated mitral stenosis, but also of the changes associated with the breakdown of that compensation. The latter enormously increases the apparent frequency and importance of dilatation of the auricle in mitral stenosis cases, while the dilatation largely masks the signs of preceding hypertrophy. That dilatation is chiefly a breakdown phenomenon is indicated by the following consideration. The 70 cases to which I have referred included 18 which came from the surgical wards. In only one of these cases was dilatation of the auricle observed. In this case the stenosis was consider-

¹ Fagge and Pye-Smith, *Principles and Practice of Medicine*, Third Edition p. 74.