These favourable changes in the symptoms were accompanied by a gradual diminution in the number of tubercle bacilli in the sputum.

FORM OF INHALER.

With regard to the form of inhaler used, I found that those in ordinary use served the purpose very well. If the patient were in an early stage of the disease I endeavoured to persuade him to use an inhaler that would cover both the nose and the mouth, but in the later stages, when there was shortness of breath, an inhaler covering the mouth only was used. I found that in the early stages of the disease particularly the patients liked the smell and taste of the cinnamon; they could use it most of the day and also at night with very little discomfort. It was also interesting to note that the continued inhalation of the vapour caused no irritation of the buccal cavity or of the air passages, and that no constitutional effects were produced.

Of course, my experience with the oil is comparatively limited, but so far as it goes I have found the oil of cinnamon to have more points in its favour when used as an inhalation in the early stages of consumption, than the drugs that hitherto have been administered in this way and for similar objects.

LANOLIN VACCINE.

By SURGEON-LIEUT.-COL. W. G. KING, D.P.H., I.M.S.,
Sanitary Commissioner for Madras.

IN his interesting contribution to the Jenner Centenary Number of the *BRITISH MEDICAL JOURNAL*, Dr. Hime, in referring to my introduction of lanolin as a preservative of animal vaccine pulp, asserts that, beyond its use by its originator, few have adopted it. This statement may represent facts correctly as to Great Britain, where the use of animal vaccination is very trivial, but it certainly does not do so when applied to India. Indeed, although—owing to official circumstances entered into by you in an editorial some four years back—lanolin vaccine has not (till within the last year) been used since 1890-91 in the Madras Presidency, when 214,876 cases were vaccinated with this preparation, its use in other parts of India may be fairly described as extensive; so that it would really be more correct were Dr. Hime to have stated that the employment of lanolin vaccine is yearly increasing at the hands of others than the originator.

As giving some idea of the extent of this employment, I may state that this is the preservative officially used in the Central Provinces, Hyderabad, and assigned districts, in Bengal, and in Mysore, and that its introduction is now contemplated in the North-West Provinces and Assam. In Mysore all vaccinations are conducted with animal vaccine pulp preserved with lanolin to the exclusion of all other modes, the advanced and independent Government of His Highness the Maharajah having possessed since 1892 the Mysore Government Vaccine Institute at Bangalore, whence it not only supplies its own territories, but much of Upper and Lower Burmah. In 1894-95, in Bengal, 60,613 were vaccinated with this preparation. During 1895-96, by means of issues from the Mysore Government Vaccine Institute, 122,258 cases were vaccinated within the Province, and 79,579 in various parts of India and Burmah. In the former case the percentage of successful to the total vaccinated was 94.20; in the latter the percentage was 94.13. In the Madras Presidency, in 1895, on its partial reintroduction in two districts, 27,540 cases were vaccinated with 94.3 per cent. success, against 89.4 per cent. for glycerinised paste in other districts.

As regards the superiority of this method of preservation in a tropical climate when compared with glycerine, there cannot be the slightest doubt, as amply proved by my own experiments and the evidence of other observers. I see no reason why this cheap and simple method should not have received attention in Great Britain by those concerned with public vaccination beyond the want of familiarity with it. Copeman has shown that, in common with glycerine, anhydrous lanolin, which is the form advised by me, has the power of inhibiting the growth of saprophytic microbes, a fact that Dr. Hime ignores when he calls attention to this special virtue of glycerine, whilst Gottstein has testified to

the protection of lanolin from penetration by microbes introduced on the exterior of the mass. So that it has been found in practice that the mere careful removal with a cleanly instrument of an upper exposed layer of lanolin vaccine secures safety, notwithstanding free exposure of the mass to the air at intervals in a manner that must be used by the busy practitioner in Great Britain, or by the public vaccinator in this country, where single cases may be vaccinated in villages many miles apart. I believe, if ever vaccination with animal lymph be widely employed by British local authorities, that this cheap and simple method of preservation will be found best to suit their requirements.

It is usually understood that anhydrous lanolin is used for the preservation of vaccine lymph. That it will act in this manner is true; but, beyond a few experiments, I have no experience of it and do not recommend it. I have advised it only for the preservation of vaccine pulp. Of course it does not follow that because lanolin is used preservation should result; it is not infallible. As in the case of the glycerine method, perfect cleanliness in the collection and during the process of preservation and storage is requisite; but if this condition be observed, lanolin vaccine, when subjected to tropical heat, will give results at the end of a period when glycerine lymph or pulp would be absolutely inert. I do not recommend its employment in the heat of India on human beings beyond forty days after preservation, although it has been successfully employed by others at the end of three months—for example, Surgeon-Lieutenant-Colonel Hendley, C.I.E., late civil surgeon, Peshawar; Brigade-Surgeon-Lieutenant-Colonel Grigg, late Sanitary Commissioner, Bengal. Nevertheless, I have seen specimens of lanolin vaccine that have been subjected to heat in long transport by runners on Indian roads in post bags arrive completely melted, thrown aside as useless without protection from dirt, and yet, when covered with fungus, still assert their power on inoculation of calves, after the lapse of four and five months. Surgeon-Lieutenant-Colonel Thomson, I.M.S., when a member of the Leprosy Commission, reported activity of a specimen of the ordinary character issued to vaccinators at the end of six months, during which time it had been exposed by opening for vaccination, and had travelled with him throughout Southern India and Burmah. Instances of activity as exhibited by inoculation of calves at the end of nine months of exposed specimens have been common.

The Mysore Government Vaccine Institute was started with vaccine descended from stock that had been preserved by me in July, 1891, on closure of my own experimental depot; this was started for experimental purposes at Vizagapatam in January, 1892, and was preserved and taken by me to Burmah till August of the same year, when it was sent to Bangalore for the purpose stated. One specimen was sent to England by me and was left there for twelve months, was then sent to me at Mandalay, and was brought by me to Madras, whence it was forwarded to Bangalore, and then again started a vigorous stock sixteen months and twenty days after preservation, and after undergoing various changes of climates and frequent opening to ascertain its condition. The longest known preservation has amounted to two years and eight months, as certified by the Superintendent, Mysore Government Vaccine Institute.

It may not be irrelevant to refer to a further matter mentioned by Dr. Hime. I am glad to find he gives his testimony to the fact that the "regeneration of calf lymph" by insertion of human lymph from time to time, as taught a few years back, is not necessary. On the opposite, I think this procedure leads to endless trouble and risk of "dying out" during cultivation. By carefully selecting typical vesicles, and by taking care that the exact stage of perfection in development of these is seized, my variola vaccine stock has been maintained in an unbroken line from calf to calf since March, 1891; at first by myself, but since 1892 under the skilled care of one of my late superintendents, Assistant-Surgeon P. Palgni, now in charge (subject to the Sanitary Commissioner, Mysore) of the Mysore Government Vaccine Institute. My experience has taught me the making of experiments whilst holding an official appointment is surrounded by difficulties, and I therefore intend to carefully avoid them; but it may be worth while my stating, for the information of such observers as would wish to push further investigations

on the subject of "regeneration," that before I ceased my work I found there exists a method that should be of value when, from accidental causes, degeneration is threatened.

It is acknowledged, from causes not yet explained, that it is only in a small percentage of cases that inoculation of the calf with small-pox virus is followed by a crop of vesicles. Nevertheless, in certain instances, whilst no local eruption is found, constitutional disturbance occurs, and is exhibited by fever and various skin conditions that I need not here detail. If an animal that has undergone this modified form of variolation be inoculated with animal vaccine, there result vesicles abnormally slow in their progress, but of perfectly typical character. Lymph so gained, when transmitted, is found to be fully regenerated, and is highly active. In these two facts—which I verified repeatedly—lies, I think, the possibility of some explanation of the true nature of vaccine at the hands of the bacteriologist.

ON THE NIPPLE REFLEX.

By JOHN B. HELLIER, M.D. LOND.,

Lecturer on Diseases of Women and Children in the Yorkshire College,
Surgeon to the Hospital for Women and Children at Leeds.

ERECTION of the nipple was formerly ascribed to vasomotor action, and to the engorgement of erectile tissue with blood. At present physiologists assign a very minor part to vascular engorgement and a prominent part to the contraction of the unstriated muscle of the nipple. To quote from the most recent edition of *Quain's Anatomy* (vol. iii, part 4, p. 286):—"The tissue of the nipple contains a large number of vessels, together with much plain muscular tissue, and its papillæ are highly sensitive. It becomes firmer and more projecting from mechanical excitement—a change caused by contraction of the muscular fibres, which form concentric circles round the base of the nipple, and radiating bands running from base to apex." Playfair writes:—"When the nipple is irritated it contracts and hardens, and by some this is attributed to its erectile properties. The vascularity, however, is not great, and it contains no true erectile tissue; the hardening is, therefore, due to muscular contraction."

I wish to draw attention to a simple method by which this reflex action of the nipple muscle may be plainly demonstrated. A marked reflex contraction of the subcutaneous muscular layer of the nipple and areola can be excited in the following way. With the finger and thumb of one—or, better, of both hands—grasp the nipple and areola, and compress them for a few seconds as firmly as is consistent with the avoidance of pain. Contraction will take place during the following minute. It will be noticed (1) that the areolar tissue becomes wrinkled, corrugated, and reduced in area; (2) that the areolar glands become more prominent; (3) that the nipple becomes more erect and prominent; (4) that the tissues are harder; (5) that there is often puckering around the nipple in the secondary areola and in the adjacent skin of the breast. The contraction soon passes off, and the reflex can be repeated. I have observed this reflex slightly marked in girls before puberty, and even in males. I have found it well marked in young women after puberty, in pregnancy, in lactation, in non-pregnant multiparæ, in women who have passed the menopause, also in patients who are unconscious through the administration of anæsthetics. It is specially obvious when the mammæ are functionally active, but I have never failed to elicit some degree of reflex after puberty. The shape of the contracted areola varies partly with the direction in which the nipple has been compressed. The phenomenon is specially well seen in women who have had several children, and who have large flaccid areolas with scarcely prominent nipples. On exciting the reflex the areola quickly contracts, showing all its glands, the nipple rises up well defined and erect, and the skin of the mammæ all round the nipple is slightly wrinkled.

When lactation is active milk may be ejected from the ducts through the simultaneous contraction of the muscle which surrounds them. During the act of suckling the infant's lips and gums, by compressing the nipple, excite nipple reflex. At the same time during suckling there is increased activity of the gland secretion and increased vascu-

larity in the gland, for it is at this time that the vasomotor element comes more into action.² Some degree of nipple reflex is also excited by cold. No doubt many medical men, as well as many mothers and nurses, are familiar with the contraction thus caused by squeezing the nipples, but it does not seem to be fully described in textbooks. For instance, in Osler's admirable *Principles and Practice of Medicine*, published last year, there is a very full table of the various spinal reflexes, but nipple reflex is omitted. Yet if anyone will make observation on a dozen female patients he will see that this nipple reflex is a phenomenon as constant and as marked as almost any reflex in the body. It will be interesting to study its pathological relations in diseases of the nervous system and in local mammary disease.

My attention was drawn to the subject in clinical gynecological work, where one has so frequently to examine the mammæ in relation to the diagnosis of early pregnancy, when the fact that fluid can be obtained from the nipples is, within certain limits, a valuable clinical sign. Conversation and correspondence with various authorities lead me to think that it is worth while to make this communication to the profession, and I hope I may thus obtain information from other observers, which, with any references to the literature of the subject, will be most welcome. Professor Michael Foster has been kind enough to refer me to articles by Eckhard³ and Röhrig.⁴ These refer incidentally to nipple reflex, though not very fully. These authors seem to take it for granted that erection is due to the contraction of plain muscular fibres, and little part is assigned to vascular engorgement.

DERMATITIS CAUSED BY ROENTGEN X RAYS.

By HENRY C. DRURY, M.D., F.R.C.P.I.,

Assistant Physician, Sir Patrick Dun's Hospital, Dublin.

A GENTLEMAN, aged about 35, having symptoms of renal calculus, was exposed to x rays to determine, if possible, the presence or absence of a stone. On May 22nd, 1896, the first exposure took place; it occupied one hour. To produce the current five cells were used at first, and later five more cells were added. The patient had on an inside thin woollen vest and a linen shirt: a penny was placed for localising purposes on the photographic film, which was applied to the back in the kidney region. The tube was in front of the middle of the abdomen, distant from one to two inches, a celluloid plate intervening between it and the patient's clothing. About three hours after the exposure was over the patient felt some nausea, but set it down to the stuffy smell of the laboratory, and thought little of it as he occasionally suffered from attacks of so-called biliousness. The plate, on being developed, not proving very satisfactory, he was again exposed, six days after—namely, May 28th, this time for an hour and a-half, the details mentioned above being the same on this occasion, except that ten cells were used from the beginning. The current also was occasionally momentarily reversed. After the patient left the laboratory he again felt great nausea, and soon felt so weak that he had to get brandy on his way home. Next evening he noticed the abdomen slightly red, as if sun-burned; a patch about 1½ inch square, just below the umbilicus, being redder than the rest. This part was that exactly opposite the platinum plate in the tube. There was no itching nor pain. The third day the redness was more intense; on the fourth some very small vesicles appeared; these very slowly increased in number and size during the next few days. This was looked upon as a simple eczema and treated accordingly. The vesicles increased to large "blebs" and became ruptured; there was, however, no pain or itching, the patient going about his usual occupation and riding a bicycle.

On June 14th, eighteen days after the second exposure, I saw the sore for the first time. It had begun to cause a great deal of discomfort from the abundant discharge pouring from it. I found a patch 7½ inches across by 8¼ inches long, extending from the ensiform cartilage to about two inches below the umbilicus. It had all the appearance of an acute

² Landois and Stirling, *Physiology*, p. 345.
Beiträge z. Anat. u. Phys., i, 12 and viii, 117.

⁴ *Archiv f. path. Anat.*, 1876.