that prolapse of the uterus may occur secondarily to hypertrophy of the cervix, and independently of any preceding prolapse of the vaginal walls. The relations of the vaginal walls and fornices to the cervix show there has not been a prolapse of the uterus secondarily to prolapse of the vaginal walls, which have dragged upon the cervix and so produced a subsequent hypertrophy, but that there has been a primary hypertrophy of the cervix in all its parts, altering the relations of the vaginal fornices, together with subsequent or secondary prolapse from increase in weight of the whole organ. The microscopic changes I must leave to another occasion.

The treatment adopted was carried out with a twofold object.

1. Removal of the hypertrophic tissues—to reduce the uterus to its normal length.

2. Cutting off of the blood supply to the cervix to prevent recurrence of over-growth—that is, dissection of the broad ligaments and ligature of the uterine arteries.

It seemed to me that simple amputation of the cervix would not be sufficient: the tendency to over-growth would probably reassert itself, and to counteract this the best method would be to ligature and cut off the greater part of the blood supply of the cervical region.

I may shortly indicate the steps of the operation as follows:—

1. Circular incision of the vaginal covering of cervix at the level of the vesical attachment.

2. Prolongation of this incision along the lateral fornices.

3. Separation of cervix from its attachments in front and behind.

4. Dissection of broad ligaments and ligature of both uterine arteries and their cervical branches.

5. Drawing down of the cervix within the vaginal tube and its antero-posterior transfixion by silver wire.

6. Circular amputation of cervix.

7. Bilateral division of cervix.

8. Formation of anterior and posterior cervical flaps in front and behind.

9. Union of vaginal and cervical mucous membranes.

10. Apposition of lateral cut surfaces of cervix and vaginal covering.

The operation was a complete success, the cervix healing by first intention. The stitches were removed on the ninth day, and the patient went home on the twentieth day. Three weeks later I examined the patient again, and found the vaginal roof healthy, the cervical canal freely patent, and the fundus

uteri at its normal level. The sound passed forwards $2\frac{3}{4}$ inches. (This patient has since menstruated normally, the flow, which was unaccompanied by any pain, lasting four days.)

CASE OF HYDATIDIFORM DEGENERATION OF THE PLACENTA.

By ARCHIBALD F. HEPWORTH, M.A. CANTAB., L.R.C.P.
AND S.E.
Steiglitz, Victoria.

ON July 26th, 1896, I was called to see Mrs. T. B., aged 27, who had been married 5 years and had three children, the youngest born December 5th, 1895; she had to discontinue suckling this child when one week old, owing to deficiency of breast milk. She had had no miscarriages. She stated that she had been "flooding" for hours, and believed she was about ten weeks pregnant. The menses had returned about three months after the birth of the last child and had reappeared for one day only two months afterwards. Since then she had had about five slight "floodings" at varying intervals, the present one being the worst.

I found a large hard tumour in the centre and lower part of the abdomen, reaching to within an inch of the umbilicus. The cervix was thick and unyielding, its anterior lip being thickened and elongated. The os uteri was undilated, and there was no shortening of the cervix. Some recent blood was present in the vagina, but no clots. *Ballotement* was absent. The temperature and pulse were normal and her general condition good. I ordered her small doses of opium, and enjoined absolute rest in bed.

During the next fortnight she had three slight attacks of hæmorrhage, each at night time, but otherwise her condition continued fairly satisfactory. The uterus was now level with the umbilicus.

On August 9th she had another flooding and complained of "labour pains." The os was found sufficiently dilated to admit the point of the finger. I therefore decided to remove the contents of the uterus, which procedure was, however, rendered somewhat difficult, owing to some considerable anteversion, the great enlargement and elongation of the anterior lip of the cervix, and the neurotic temperament of the patient. On dilating sufficiently to admit two fingers no solid portions of a fetus could be made out, but a large number of broken down blood clots were removed. "Pains" came

on regularly, being assisted by a hypodermic injection of ergotinin. During the next three hours 24 ounces of clot came away, followed by a considerable quantity of small pieces of semi-fleshy *débris* which measured 20 ounces more. No fragments of bone, cartilage, skin, or foetal flesh came away. As there was evidently a mass of adherent material still remaining, I obtained the assistance of Dr. F. J. Newman, of Geelong, and under chloroform thoroughly dilated the cervix, and removed with fingers and curette the remaining contents of the uterus. These were extremely friable, and broke down under very slight pressure. The whole inner surface of the uterus was then thoroughly curetted, and afterwards irrigated with hot carbolised water. Iodised phenol was then injected by Baum's intrauterine syringe, and the uterus left well contracted.

The patient had a rigor immediately afterwards, but subsequently had a quiet night; she was then put on a mixture of quinine, ergot, and digitalis. The next morning her temperature being 103° F., and pulse 160, the uterus was again irrigated, and a small quantity of *débris* cleared away. The same evening her temperature was 100.6° F., and pulse 130. Next morning her temperature was normal, and pulse 100. Afterwards her recovery was uninter-

A.—Outline of normal uterus and vagina. (Half-size.) B.—Outline of uterus, etc., in present case showing position when patient has been for some time recumbent. C.—Ditto, when patient has been walking or straining. 11—Vaginal portion of cervix. 22—Intermediate portion. 33—Supravaginal portion.

rupted, and she was allowed up on the tenth day. The material removed showed hundreds of cyst-like formations, some about the size of peas or currants, others very small, like pinheads; each contained a clear jelly-like fluid, giving the appearance of small white berries.

The case was thus, undoubtedly, one of vesicular or hydatiform degeneration of the placenta. It will be noticed that the cardinal symptoms of the disease were all present:—

1. The more rapid development of the uterus than occurs in ordinary pregnancy.
2. The repeated attacks of uterine hæmorrhage.
3. The disappearance of the embryo.
4. In this case the absence of "ballotement" was somewhat suggestive.

A CASE OF TRUE HÆMATOSALPINX : LAPAROTOMY : RECOVERY.

By JOSEPH GRIFFITHS, M.A.CANTAB., M.D.EDIN.,
F.R.C.S.ENG.,

Additional Examiner in Surgery in the University of Cambridge, and
Surgeon to Addenbrooke's Hospital.

UNTIL recent years the term "hæmatosalpinx" has been used by writers to denote two distinct morbid conditions: the one arising from the retention of menses, as in cases of atresia vaginæ; and the other a local disease characterised by hæmorrhage into the lumen of the tube. Since Dr. Martin, of Berlin, in 1889 discovered that in a case of hæmatosalpinx under his care there were present in the wall of the tube chorionic villi, though no trace of a fœtus could be found, it has been assumed that all cases of the latter variety arise from abortive extrauterine gestation.¹ This view has in the more recent works overshadowed the older, and until November 4th, 1891, when Dr. Herman showed a case which he termed hydro-hæmatosalpinx before the Obstetrical Society of London, there was no example illustrating the manner in which a simple hydrosalpinx may develop into a true hæmatosalpinx. The compound name given to this unique case sufficiently explains its nature. It occurred in a young woman, the tube was distended and in its cavity was found blood containing serous fluid. There was, however, no evidence found on microscopic examination of the wall to show the manner in which the extravasation of blood took place. In short, this proved to be an example of a mixture between hydrosalpinx and hæmatosalpinx. No definite clinical symptoms appear to have been associated with this morbid state.

In the case herein described one tube—the left—was found in the state described by Dr. Herman as hydro-hæmatosalpinx, and the other—the right—which gave rise to the symptoms, proved to be, I think, an example of true hæmatosalpinx, that is, a hæmatosalpinx in which no trace of ovum structures could be detected. The suggestion to which the case of hydro-hæmatosalpinx above alluded to gives rise—namely, that a hæmatosalpinx may arise from a pre-existing hydrosalpinx—receives, it seems to me, definite confirmation from the following case.

HISTORY.

M. A. K., aged 32, was admitted into Addenbrooke's Hospital, Cambridge, on August 29th, 1896; married, six children, the eldest 16 and the youngest 2½ years; pale, depressed, but healthy-looking; pulse 110, regular and moderately strong; tongue slightly furred and tremulous; very thirsty and bowels somewhat constipated; no cough; urine normal. There is present a more or less constant bloody vaginal discharge. In the inner and lower part of the right inguinal region, just above Poupart's ligament, there is a lobulated slightly tender swelling situated immediately beneath the anterior abdominal wall, and therefore in the pelvis. This was found by vaginal examination to be in the right and anterior part of the true pelvis, lying on the bladder, freely movable and distinct from the uterus, which was not enlarged but displaced backwards and to the left. The cervix uteri was enlarged and patulous, and the vaginal mucous membrane swollen and rather hot.

History.—In November, 1895, she suffered from an abortion after three months' amenorrhœa; a membranous bag came away. After this she got about, but suffered from periodical bleedings, and in between times from a blood-stained discharge. A week or two after this abortion she discovered the lump that can now be felt in the right inguinal region, and she says that it was then of smaller size. This state continued with

¹ Mr. Bland Sutton, in the last edition of his *Surgical Diseases of the Ovary, etc.*, emphasises the view that many of the cases hitherto regarded as hæmatosalpinx belonging to the second variety are in reality examples of abortive tubal gestation, and he makes reference to several instances in which this mistake arose, and especially to a specimen in the Pathological Museum of St. Bartholomew's Hospital. He does not even consider the subject of what I herein describe as true hæmatosalpinx.

hardly any change until July 2nd, 1896, when, while walking about, she was suddenly struck with lancinating pains in the lower part of the body, and simultaneously suffered a great loss of blood from the vagina; no clots. Mr. Ord, of Mildenhall, who was called in to see her, found the lump already referred to in the pelvis, and came to the conclusion that probably it was of the nature of an extrauterine pregnancy. All the symptoms soon quieted down. On July 29th she went to Bury, and while walking about was again attacked with similar pains, and a blood discharge from the vagina. Again all symptoms subsided with rest in bed. On August 23rd, while returning home from Bury, she was again attacked in a similar way. Six days later she was conveyed to the hospital.

On September 1st, 1896, I performed laparotomy, and found that the swelling on the right side involved the Fallopian tube which was removed, together with an enlarged ovary, by transfixing the pedicle and tying a Staffordshire knot. On examination of the left side of the pelvis, the left Fallopian tube was found distended and occluded at its fimbriated end, and, in consequence, it was deemed advisable to remove it. Recovery uneventful.

DESCRIPTION OF THE SPECIMENS.

The Right Fallopian Tube and Ovary.—The outer end of the Fallopian tube is distended into a lobulated congested fluctuating cystic mass with the omentum attached to the lower part of the anterior surface. The wall of the sac is very thin near the upper part. Behind is the enlarged cystic ovary. On manipulating this swelling and endeavouring to deliver it whole, the thin part of the wall ruptured, and much dark coagulated blood escaped, leaving the mass a good deal smaller. The inner part of the tube appears to be natural. In a longitudinal section through the tube and ovary it is seen (as in a diagrammatic representation Fig. 1) that the free

Fig. 1.—Coronal section through right Fallopian tube and ovary (natural size).

end of the tube consists of a thick-walled cyst, 1½ inch in diameter, containing clotted blood, with a portion of omentum adhering to its lower part. The wall is in most places one-sixth of an inch in thickness, and appears to consist in the main of fibrous tissue. The inner surface of the cyst is everywhere uneven and rough. The part of the tube next to this is distended with blood clot, so that it is about half an inch in diameter, but its walls are merely thinned from distension, and not otherwise altered. From this part the tube rapidly diminishes to its normal size, and becomes of normal appearance. The ovary is enlarged, and in its substance there are three cysts, one about 1 inch in diameter and the others each about one-sixth of an inch. The wall of the cystic end of the Fallopian tube is found under the microscope to consist of a thick fibrous wall, containing unstriated muscular fibres, lined by a layer of granulation tissue in many parts of which there are degenerative changes, and into which many small hæmorrhages have taken place. The muscle fibres, which are in the outer part of the wall, are of small size, and are widely separated by much fibrous connective tissue. The bleeding no doubt arose from the lining of granulation tissue. The part of the tube next to the cystic outer end, though distended with coagulated blood, is thinned only, and not otherwise altered; and the tube near the uterine end is