brush the teeth after every meal, and by rinsing with cold water you get an early intimation of danger.

Turning out of Bed at Night.—No amount of precaution can make this otherwise than dangerous. But if an arrangement can be made by which the clothes can be kept aired and warm, and a cup of hot milk with a teaspoonful of brandy can be procured, the risks can be reduced to a minimum.

One word about holidays. Every medical man, if possible, should have an outdoor sport of some kind. Golf and cycling are good, but perhaps the best of all is fly-fishing. It takes one usually into beautiful country, the exercise is gentle and varied, the interest absorbing, and to my mind is far better for the jaded practitioner than scampering half over Europe in a hurry in a second-class railway carriage in charge of a party of tourists.

Gentlemen, I might go on for ever, but I will bring my remarks to a close. I can fancy some of you saying, "These suggestions are all very well, but do you carry them out yourself?" Well, honestly, I do not always, but when I forget them I regret it, and I have reaped health and happiness by carrying them out in the main. I feel it is presumptuous of me to address you on a subject of which you all know as much or more than I do myself, but if any suggestion of mine should have the effect of making any one of you take more care of yourselves in any one particular than you do at present, I shall be amply rewarded.

SOME POINTS ON SACRO-ILIAC DISEASE, WITH A REPORT OF A CASE.¹

By F. B. JUDGE-BALDWIN, M.R.C.S.ENG.,
Honorary Surgeon to the Rotherham Hospital.

THE literature on the subject of disease of the sacro-iliac joint and its treatment is scanty. In Holmes's *System of Surgery* mention is made of the disease, but the results of treatment are stated to be anything but satisfactory, and the same remarks apply to Erichsen's *Surgery* (Edition 1888). Nélaton described the disease, and in 1821 Boyer called it "a chronic scrofulous affection of the sacro-iliac joint." Velpeau in 1824 drew attention to suppuration in the joint during the puerperium, and some years later he described cases not due to the puerperal state.

In 1833 the *Dictionnaire Médical*, by Laugier and Hahn, gave a summary of what was then known of the disease, directing attention more especially to some points about lengthening and shortening of the limb on the affected side. Gwilt, in 1843, in a treatise on joint affections, gave some valuable facts about this disease. For the Faculty of Paris six theses on this subject were written between 1838 and 1847.

Erichsen, in the *Lancet*, 1859, p. 25, gave the first clinical lecture, giving a lucid description of the disease, its course and treatment. He there says that operative treatment is inadvisable. In 1875 Christopher Heath published a series of cases with a view to operative treatment.

The year 1879 gave us a valuable series of cases from Professor Lewis Sayre, and also an interesting case of scraping the joint with the only instrument he had with him—a pruning knife. Gambier, in 1887, operated on two cases successfully.

Van Hook, in the *Annals of Surgery*, 1888, gives the best summary of the disease up to that date, collecting 51 cases from the records of England, America, and the Continent. In 1889 Mr. Mayo Collier in the *Lancet*, vol. ii, p. 787, described a case in which he trephined the joint before suppuration with excellent results. The patient had a clinical history of tubercle, and had had hæmoptysis, but the supposed immediate cause was the lifting of a heavy box.

In 1893 Makins, in the *Clinical Society's Transactions*, had an article on sacro-iliac disease, and in the early part of 1895 Mr. Golding-Bird gave a "clinical" in Guy's on the subject, published in the *Guy's Hospital Gazette*. Mr. Golding-Bird has done several cases, trephining the joint, in one instance freeing a psoas abscess from within. He has also a paper in the *Clinical Society's Reports*, 1895, with cases.

The disease is obviously divided into two stages—non-

suppurative and suppurative. Of the 51 cases collected by Van Hook, 21 were non-suppurative, 30 suppurative; of the former, 94 per cent. recovered; of the latter, 7.9 per cent. only recovered without operation.

The etiology of the disease is obscure. Sayre says that trauma is always the determining cause, but the puerperium has its cases. I think perhaps that, given a tuberculous patient, either cause, trauma or the puerperium, may determine the onset of the disease. It is worthy of remark that in the three cases I have seen the characteristic tuberculous aspect has been present.

SYMPTOMS.

The symptoms of the disease are pain, lameness, tumefaction, and abscess. The pain is insidious, becomes rapidly severe, is gnawing or rheumatic in character, extending to the anus, and down to the knee at times, and aggravated by all movements of the lower half of the body. The pain is also increased by pressing the ilia together or by pulling them apart, and there is intense pain on single-finger pressure over the joint. The line of the joint is given by Mr. Golding-Bird as follows: a line is drawn from a point 3 inches behind the anterior superior iliac spine to the posterior inferior spine. The centre of the joint is on this line $\frac{1}{2}$ inch from the posterior inferior spine. Lying on the affected side is impossible; this symptom is supposed to be pathognomonic.

The lameness is such that the gait is curious and striking, the step being shortened. The attitude changes, the patient trying to throw the whole weight of the body on the other joint; hence occurs the tilted pelvis, which, I think, gave rise to Hahn's erroneous belief that the limb was shortened. Tumefaction is not a constant symptom, but it is in the line of the joint when present, and is easily seen because the buttock is flattened from atrophy of the muscles due to the patient's desire to avoid the use of the side. If abscesses form, these may be extrapelvic or intrapelvic, and, if extrapelvic, the abscess appears over the glutei or at the crest of the ilium, bulging upwards. Intrapelvic abscesses occur as psoas abscesses, at the edge of the gluteus maximus, down into the rectum, and into the spinal canal across to the other joint.

The diagnosis must be made from gout, syphilis, gonorrhoeal rheumatism, affections of the muscles, viscera, or bones, tumours, and neuralgia. As aids to diagnosis I may mention the absence of internal rotation of the leg and of spinal curvature, rectal touch of the anterior surface of the joint, inability to lie on the affected side, and point pain on pressure over the joint.

I have said nothing of the pathology of this disease, since it is common to all joints of this class. One point may be of interest, namely, that the cartilage is thicker on the sacrum, and that in the early period of life, occasionally in the adult, and always in the female during pregnancy, the cartilages are smooth and lined by delicate synovial membranes, with cavities in the cartilages.

TREATMENT.

Erichsen, in his *Science and Art of Surgery* (edition 1888), says: "I have never seen a patient recover after the full development of the disease," and after suppuration has set in he recommends treatment on general principles, fixation, etc. This I think comprises the substance of the literature of the subject with one or two personal observations.

The following case is, I believe, unique owing to its severity and the success following a bold treatment which was justified by the result:—

Mrs. M., aged 26. Family history: Father aged 59, mother aged 59, grandfather aged 80, all healthy; eleven in family, nine living, patient the eighth child; first children living, all healthy; no suspicion of tubercle except in the case of one brother, a labourer, who died aged 29 (after heavy drinking) of some lung trouble (? pneumonia); the other brother died aged 2 years from some unknown cause in twenty-four hours. Patient's past history: She had good health until her marriage three years ago in December, 1892, and was well during her pregnancy. The child was born in September, 1893; instruments were used, and she did well until the fourteenth day, when an abscess of the breast commenced. She was five months recovering from this. She then felt an aching at the bottom of the back, which seemed to affect the left leg in walking; the pain was severe, but not continuous, like a broken bone in character. She remained in bed without active treatment for a week, and then felt better. Some short time after this she felt pain in the groin and lower half of the back. This was just before Christmas, 1894

¹ Read before the Sheffield Medico-Chirurgical Society.

In January, 1895, she noticed a swelling in the left groin, growing at first slowly, and then rapidly. Her general health was now giving way. She had much difficulty in sitting down from pain in the back, which was much increased when rising from a chair; no pain when seated. During this time she had had no treatment.

She came to my out-patients in April, 1895, complaining of a large swelling, occupying the position of a psoas abscess. There was no spinal angularity, no pain over the lumbar region, and no abnormality about the hip-joint. She further complained of losing weight, of pain of a gnawing, aching character over the left buttock, close to the spine. This was increased on pressure of the ilia together or apart. The swelling in the thigh did not alter its shape or position on flexion of the leg; it had no impulse on coughing, but had a tense, deep sense of fluctuation. The circumference of the leg was 2 inches greater than that of its fellow. Patient walked slightly lame, with short steps. On point pressure over the sacro-iliac joint she complained of sickening pain. She could not lie on the affected side. Urine healthy; all other organs healthy. The patient had a tuberculous aspect, and was pale and haggard. The risks of an operation were explained to her, and also the probable termination of the disease if left alone. She consented to an operation, was taken in to Rotherham Hospital, and kept at rest for fourteen days.

OPERATION

On May 1st, 1896, chloroform was administered, and the abscess in the front of the thigh was opened first, and about a pint of thin pus was evacuated. The sac was very thick, and separated the muscles of the thigh deeply. The patient was then turned on the right side with her left leg flexed on the abdomen, and supported on a pillow. The incision was made, according to Mr. Golding-Bird's directions, as follows: The line of the joint as stated above is found by drawing a line from a point 3 inches behind the anterior superior iliac spine to the posterior inferior spine. A point on this line $\frac{1}{2}$ inch from the posterior inferior spine marks the centre of the joint. A horse-shoe-shaped incision about 2 inches wide, having this point as a centre and having its convexity downwards, was made, and the flap turned up and stitched or pinned to the skin above. The muscles were divided down to the periosteum and held well away.

After freely clearing the surface over the joint I saw that it looked spotty and greyish in tint. I had intended to trephine, but on pressing a chisel on this surface I found it so rotten that chiselling was commenced at once. The whole of the joint was removed right through into the pelvis, and soft bone was also found and followed up over the surface of the sacrum between the foramina; two fingers could easily be entered at this stage. The bone was well curetted for fear of any spot being overlooked. The cavity was well washed out with hot water, and afterwards filled with an emulsion of sulphur. A large drainage tube was inserted, and the flap stitched round the tube. The abscess cavity in front was now again dealt with; a long tube was passed up into the pelvis from the opening in the thigh, and a syringe of zinc chloride solution (1 in 20) was injected, the abscess cavity itself being freely irrigated with the same solution. A dressing of double cyanide gauze was applied, a long side splint put on, and the patient was sent back to bed.

AFTER-HISTORY.

Considerable pain followed in the evening, but no rise of temperature. The dressing was changed three or four times during the next 18 hours. Much retching and faintness followed that night and the following morning, and there was a temporary collapse early in the morning, but that was met by a brandy and milk enema.

Forty-eight hours after the operation the sacral wound was irrigated with corrosive sublimate solution (1 in 2,000). The sulphur emulsion caused considerable sloughing of tissue, and I should not use it in a future case.

There is little more to say except that during recovery there were occasional rises of temperature, never excessive except once, when it reached 104.4° . It was relieved by quinine. The temperature was never over 100° after July 5th, only once above 99° after July 22nd, and was practically normal after August 9th. Creolin (1 in 60) was used for irrigating after June 6th.

The patient was allowed up on a spinal carriage on July 24th, and on crutches with a Thomas's splint applied on August 13th; the crutches were thrown away on September 10th. The weight of the patient increased at about the rate of 1 lb. a week during convalescence. The sinus which remained was injected with iodoform emulsion once a week, and once or twice wiped with chloride of zinc solution. It remained open until March 7th, 1896, when it closed. No dead

bone was discharged, although the probe passed on to the surface of doubtful bone.

When last seen on October 31st, 1896, the patient was perfectly well. She had had no ill-health since the operation, and walked so well that it would be difficult to detect that she had ever had any disease of the joint.

COMMENTS.

Six years ago a similar case came under my care, and I must acknowledge that, although I dealt with the psoas abscess, I could not discover its origin to my satisfaction. However, the case left for home, but ultimately ended in hospital, with extension of disease to the other joint. She died from prolonged suppuration.

A necropsy I made in 1879 on a man aged 27 is interesting. He had psoas and gluteal abscesses, caseous material in both sacro-iliac joints, and one caseous spot in the spinal canal between the dura mater and the arachnoid in its tubular contraction about opposite the second sacral vertebra. In the lungs were scattered tubercles, the liver and spleen were lardaceous, there were spots of ragged ulceration in the intestines, and the mesenteric glands were caseating, as was also the prostate. The kidneys were in various parts caseating, cystic, and lardaceous.

I cannot conclude without thanking Dr. Reid, the house surgeon, for his constant and valuable attention to the case; to his care I think much of the success was due.

THE ARTIFICIAL FEEDING OF CHILDREN.

By ANDREW MACPHAIL, M.D., M.R.C.S., L.R.C.P.,

Montreal.

Professor of Pathology and Diseases of Children, University of Bishop's College, Montreal, Canada.

It is only in recent years, since economy has come to be at once the test and aim of civilisation, that waste of child life has excited much interest. Now the problem of infant mortality is one desirable to be faced, and it is being faced, if one can judge from the activity displayed in every country in this branch of research.

It cannot be claimed that we have attained even a reasonable degree of success in dealing with infants if the death-rate can be taken as a guide. In the BRITISH MEDICAL JOURNAL of January 25th are the returns from sixty-five English towns, from which it appears that, of all children born, 19 per cent. die within the first year of life. The Registrar-General's return for the first quarter of 1896 shows the infant mortality for thirty-two provincial towns to have been 15.5 per cent., which is very close to that of the second quarter, and may be regarded as fairly constant. In Paris, the death-rate is the same; in Berlin, 26; in Austria, 25; in Russia, 26; and in Munich, 32 per 100 births. In America, mortality statistics fall under local control and are fragmentary. One reads, for example, "In Baltimore 39 per cent. of the deaths last year occurred in children under 5 years of age, and in Chicago the rate was rather higher." If such things occur in fairly-civilised countries, the conditions elsewhere must be much worse, as we know they are.

Now it is manifestly impossible to isolate the mortality due to improper feeding, but anyone who has to do with the care of infants will appreciate its full significance. The profoundness of this appreciation is revealed in the earnestness of the attempts to procure some efficient substitute for the natural food of the child; but all experience in the use of this or that food is fallacious, since personal interest and professional zeal do so modify the other conditions of the child's existence. Indeed it is probable that too much stress is being laid upon the nature of the food and not enough upon the necessity for general maternal care.

Much admirable work is being done upon the sterilisation of milk and upon the bacteria infesting the intestine, but the results have not yet had their full clinical realisation since the domestic kitchen cannot be transformed into a bacteriological laboratory.

The more recent discussion on this subject in the JOURNAL was commenced on December 7th, 1895, in a "fragment" by Sir William Priestley, in which he refers to the work of Budin and Chavane in Paris. He there says: "Sir Dyce