There remains the question whether anything can be done to improve the stability of the nervous centres by such remedies as are known to be of service when the nervous tone has been impaired, or when the functional activity of the brain has been disturbed by conditions of the circulation.

I have already stated that in epilepsy generally the pulse is small and the arterial tension low. Exceptions occur, and in my experience, when a young epileptic has a pulse of unduly high tension, the prognosis is favourable. The tension can be brought down by care in diet, and by eliminants. In post-hemiplegic epilepsy, again, when the pressure within the arterial system, which has resulted in the rupture of a cerebral vessel, persists, the frequency of the fits can be materially reduced—I think I have seen them kept off altogether—by keeping down the pulse tension, which can best be effected by mild mercurial aperients.

For the generally lowered vitality of epileptics I have given phosphorus and the hypophosphites, or arsenic with strychnine or quinine, and sometimes iron. I am disposed to think that the arsenic, which is often given at the same time with the bromides to prevent the rash, may sometimes have a considerable share in the production of good results. I cannot say definitely that I have seen cures, because in consulting-room practice we cannot follow up patients who recover, but I have seen many instances of remarkable improvement, the fits becoming less frequent and severe month by month. A satisfactory feature of such treatment has been the increased sense of well-being and enjoyment of life which has attended it, the improvement of the memory, and generally increased intellectual efficiency. This has been particularly noteworthy when bromides which had been taken for some time were left off.

Epilepsy will repay further study. Jacksonian convulsions have been differentiated from true epilepsy; the recurrent convulsions caused by local or general thickening of the cranial bones of traumatic or idiopathic origin ought to form a distinct category, and I think that further distinctions will be established, so that we shall speak of the epilepsies rather than of epilepsy. The different forms will then have each its own treatment. It is because my thoughts have moved in this direction that I have brought the subject before you.

A BRIEF HISTORY OF THE OPERATIONS PRACTISED FOR CANCER OF THE BREAST.

By SIR WILLIAM M. BANKS, LL.D., F.R.C.S.,

Surgeon to the Liverpool Royal Infirmary.

In this present age we consider that we live very fast, but, if we turn to the books and newspapers of the beginning of last century, we shall find that our ancestors a hundred years ago believed that they lived at a very rapid pace, while our descendants at the end of this century (when there will be Saturday to Monday cheap excursions to Mars) will look with pitying amazement on the poor old things who vegetated their lives away in 1901. But it must be admitted that the acute rush of life of the present day has come with rather startling suddenness upon us. It is essentially the product of the railway, the telegraph, and the telephone, and, as such, it has occurred during the lifetime of men who are not so far past middle age. So that plenty of us are capable of judging whether this very rapid advance has in every respect been a salutary one. For it must always be remembered that in all Nature the principle of slow growth and long life, of rapid life and speedy extinction prevails. This rule holds good with regard to a man's reputation. Formerly a surgeon took time over his work and thought about it and made sure that it was sound and would stand the test of trial by his brethren. After that fashion worked Syme and Simpson, Christison and Bennett, Graves, Stokes, Paget, Lister, Murchison, Wilks and all those men who have been the real makers of modern medicine. For they and their fellows of the same stamp

planned the structure and laid its foundations, being ignorant of the meaning of self-advertisement and solely intent upon the production of sound and true work. It is a question whether the same fine spirit is animating us now, for praise seems accorded only to the most recent and most obtrusive discoverer, whose discoveries are at the end of six months too often relegated to limbo.

It is due to this state of feeling that there has arisen a complete revulsion from the condition of matters which prevailed in our profession down to the beginning of the 18th century. Till that period a blind and unreasoning worship of the doctrines of the great men of the past was universal. This veneration, moreover, was of the most bigoted and intolerant kind. One cannot obtain a better glimpse of it than by reading the life of Ambroise Paré, and seeing how he was tormented by Gournmelen, the Dean of the Faculty of Medicine. As Mr. Stephen Paget in his life of Paré says, referring to a book on surgery published by Gournmelen, "This book is an idiotic appeal to authority and tradition; the very thing that Paracelsus would have liked to burn. The rest of Paré's answer to Gournmelen is no less emphatic, showing the wonderful vigour and spirit of his old age. He appeals once and for ever from tradition to experience; he will show this *petit maître*, this ass, what it really means to be a surgeon." The unfortunate thing is that the pendulum has swung so far in the opposite direction that it is the man who appeals to experience who is now considered the ass. The idea is that there is no time to attend to persons who mildly plead long experience as a reason why they should be heard. Nothing is worthy of consideration if it is more than eighteen months old. To use the vulgar phrase of the day it is not "up to date," which phrase is too often synonymous with crude and hurried and superficial. When a man takes to critical moralising people say at once that he is a grumbler. I admit the soft impeachment, and what I venture very gently to grumble at is a leading article in the BRITISH MEDICAL JOURNAL for November 16th, 1901, entitled Operations for Gastric Ulcer. A paragraph runs thus:

Twenty years ago, and even more recently, it was considered, in operating upon the breast, for example, that the whole malignant growth should be removed, the incision being made at a short distance from the margin of the tumour in supposed healthy tissue. The researches of Stiles, Heidenhain, and Rotter, and the operation of Halsted founded upon them, have entirely revolutionised that procedure. It is now accepted that, however small the primary growth may be, it is essential that the whole breast, the lymphatic vessels draining the breast, the chains of glands into which the vessels pour, should all be taken away. So the modern and the ancient operations may be contrasted. In the latter the tumour itself was excised, and, as an extreme measure, the glands, if felt to be enlarged, were removed through a separate incision; in the former the tumour, the whole of the breast in which it lies, the whole of the lymphatic system, vessels, and glands accessible, are mercilessly cleared away by patient dissection. Of the improved results that have followed this wide removal all surgeons are, we believe, convinced.

It must be obvious that there can be only one interpretation of this paragraph—namely, that until Dr. Halsted brought out his operation, founded upon the microscopic researches of the three other gentlemen mentioned, there had been no improvement whatever upon the old method of removing the growth by incisions made at a short distance from its margin through supposedly healthy tissue. I cannot but notice the perseverance with which this statement is being constantly reiterated, for in the BRITISH MEDICAL JOURNAL (April, 1900) I thought I had produced sufficient evidence of its extreme unfairness. But here it crops up again, Stiles and Heidenhain being now reinforced by one Rotter. So I am a second time compelled, greatly against my own inclination, to review the position.

It has occurred to me, however, that to look back and see what the surgeons of old thought and did in the matter of cancer of the breast might have some little interest from the antiquarian point of view. Hildanus, in 1646, in his *Observationes Chirurgicae*, writes a letter to his friend Dr. Andreas Weickius upon the Excision of Tumours of the Axillæ and the Removal of the Breasts. Under the former head he makes some most sensible remarks:

But before everything we must carefully make out whether the tumour can be shifted and moved from one point to another, and whether it can be radically excised. For the operation will be fruitless if any part of the tumour, however minute—nay, even the membranes in which tumours of this sort are usually enveloped—be left behind. For the disease sprouts up again, and becomes more malignant than ever, while there is no hope that we can remove what remains behind by cauterising.

Here is reproduced a drawing of the peculiar forceps and knife figured in Hildanus's work.

Fig. 1.—Hildanus. *Figura forcipis et cultelli separatorii.*

In the year 1706 the well-known Conrad Sprengell translated into English the *Chirurgica Curiosa* of M. G. Permannus, who was Chief Chirurgion of the City of Breslau, in Germany, and of the Hospitals of St. Job and All Saints. He dedicated his book to the Emperor Leopold, and it is really worth a minute's reading to notice what a splendid thing a dedication was in those days. After about a page of titles of the Emperor it runs thus:

Most Illustrious Sir: I humbly implore your Imperial Majesty's most gracious pardon, that a person of no greater figure in your vast dominions than myself, though one of your most dutiful and loyal subjects, should presume to prostrate myself at the feet of your Imperial Majesty, and humbly present this book to your Royal Hands; but since all lesser things seek the protection of greater; that rivulets run with that design to the ocean, and small fires ascending the Superior Regions, direct their heat and flame towards the sun: Why should I fear in this little Offering to displease the greatest of Monarchs, the Sun of our State Meridian; whose Magnific and Generous Soul never despises the more minute endeavours of his meanest subjects for the public good: under which encouragement I humbly beg your Imperial Majesty's gracious acceptance of this work, which with great labour and industry I have compiled for the common benefit of the Universe, and I shall esteem that signal condescension in your Imperial Majesty as the greatest honour that could be conferred upon me.

To some of the junior members of the profession, who may not have had time to look into old books, it may be interesting to see how a distinguished surgeon addressed a great monarch in times of old. But it must be remembered that the elaborately submissive manner in which he does so did not at that time imply anything degrading; it was the accepted manner of addressing the great ones of the earth. Surgeon Permannus writes as follows:

Now we will proceed to the cure of a cancer by manual operation, in which divers methods are prescribed for the performance of it. Selingem recommends the old way of thrusting two crooked needles and silk through it crosswise, and leaving the ends hanging out of the cancer, to pull up the breast with them. For my part I never used such a tedious method, but have had more commiseration for my patients, as this city of Breslau can testify, where I have cut nine cancers out of women's breasts from the foundation very successfully, and yet never used this way, for before the needles could be thrust through the breast, which is a prodigious pain to the patient, and the ends of the strings ty'd together to put the hands into I have cut off the breast in half the time only with my hand. Some there are that use hooks instead of needles, and others a particular kind of a bag, in which they enclose the whole breast, and tie it about it where the place is sound, thinking they have done an extraordinary matter, but I still approving the way that has constantly succeeded in my practice, shall proceed to direct you in it. When your patient has been sufficiently prepared for this operation by the advice of an able physician, and you have all things necessary on a table by you, place your patient on a chair towards the light, tie his arms and hands close and fast to his body about the middle; and, if you find it necessary, tie a silk string twice round the breast towards the ribs as hard as the patient can suffer it; to which string you may fasten two other strings, tying them together on a knot, to draw the breast towards by them; but this is only to be done when you think you cannot hold the breast firm enough with your hand. Then take a crooked knife in your right hand, and cut towards the right arm, if you are to cut off the right breast; and, on the contrary, if it be the left; and with all the care and caution imaginable cut out all you find hard or knotty, and as quick as possibly you can, while your assistant in the meantime wip'es away all the blood, that you may see the bottom and feel anything that is left behind it in order to excoriate it, or touch it two or three times with a hot iron.

Let us now pass on to 1740, when that splendid old surgeon, Samuel Sharpe of Guy's Hospital, wrote his *Treatise on the Operations of Surgery*. He held that in extirpating the scirrhus, if it were small, a longitudinal incision would dilate

sufficiently for the operation, but, if too large to be dissected out in that manner, an oval piece of skin was to be cut through first, the size of which was to be proportioned to that of the tumours; for example, if the swelling were 5 inches long and 3 broad, the oval piece of skin cut away was to be nearly of the same length, and about an inch and a-half in breadth. In taking off the whole breast, the skin might be very much preserved by making the wound of it a great deal less than the basis of the breast, which should be carefully cleared away from the pectoral muscle. When the disease was attended with knots in the armpit, no service could be done by amputation unless the knots were taken away, for there was no sort of dependence to be laid on them subsiding by the discharge of the wound of the breast. The possibility of extirpating these knots without wounding the great vessels was very much questioned by surgeons, but he had done it when they had not laid backwards and deep.

Coming to 1745, to Heister's wonderful book on surgery, he mentions the method of raising the breast by cords passed crucially across its base, and I have reproduced his two pictures illustrating the method. He mentions also the prac-

Fig. 2.—Heister. Method of removing breast.

Fig. 3.—Heister. Introduction of cords to pull up breast.

tice of some surgeons who used a thing closely resembling a large carving fork to pull up the breast, while Bidlow thrust through it an instrument very like a small sword. Helvetius, he says, used an instrument like a gigantic tuning fork, with movable sides, which was shoved up on the breast, and the sides screwed together. All these were for the purpose of dragging the breast well away from the thoracic wall. In 1778 William Hunter was lecturing, and I possess a beautiful copy of manuscript notes of his lectures by a certain T. Cowley. Nothing better could be written than this:

Amputation of the breast is an easy operation but a doubtful remedy.

The little success attending it deters the patient, wherefore it ought not to be undertaken without great probability of cure, as we run the risk of a general prejudice against the operation, which may so affright others that they will not run the hazard, tho' their disease may be really curable.

Hunter directs the surgeon to sit down before the patient, to take the breast into one hand and pull it down to make the skin tense that it may be cut more easily, and then, if the tumour be small, he need only make a longitudinal incision and dissect it out, but otherwise he is to make a semicircular one on the upper part, and dissect away all the indurated part from above downwards, finishing the operation below. If any indurated part be found left he must then dig it out, and, if the hæmorrhage be violent, take up the vessels with a needle. In the next place, if there be any indurated glands in the axilla, an incision must be carried upwards from the other wound, and they must be extirpated either with the knife or ligature.¹

When I graduated in 1864 matters seem to have come to their worst point, for the teaching was that a narrow, elliptical bit of skin, which should include the nipple, was all that was to be removed, wherever the tumour might be situated. The breast and tumour being then removed the infected skin was replaced. It was a simple job, easily done and easily recovered from, but so absolutely hopeless from a curative point of view that a large number of men recommended their patients not to submit to it, and I believe that they were right. People lived longer when they were not touched than when they were operated upon imperfectly. If the axillary glands were affected then that was a bar to all operation. Even the removal of a mass of tuberculous glands from the axilla I have seen denounced by one of the best surgeons of the day. A few years afterwards in 1867 Mr. Moore of the Middlesex Hospital, whose experience of cancer in that charity was great, wrote what is now an almost historic paper on "Inadequate Operations in Cancer," which set a great many minds thinking, inasmuch as it was a revolt against constituted authority. It made me think too, so that I became deeply interested in the question. Moore's paper was one of the proper stamp, written out of the fulness of experience, and not an advertisement for practice, from which experience should ultimately come. During the following ten years after Moore's protest great progress was made. Lister and Bryant and Annandale and Heath attacked secondary glandular growths in a manner at once persevering and bold, and demonstrated that all that was required for their safe removal was careful, patient dissection. Among English-speaking surgeons Samuel Gross of Philadelphia was the next who made a notable move in favour of the true removal of breast cancers², and to him much is due for the earnestness and enthusiasm with which he pushed his cause, as will be remembered by all those who, like myself, had the privilege of being his friend.

Coming to my personal share in this matter of the surgery of the breast, I feel it a very painful thing that, on my own part and for the credit of those who have worked on the same lines, I should be compelled a second time in this JOURNAL to defend the work we have done, and to expose the very persistent attempts that have recently been made to ignore our work and to hand over to Dr. Halsted the entire credit thereof. I believe I must have now removed the mamma freely and cleared out the axilla about 300 times, and in the JOURNAL for April 2nd, 1900, will be found the history of about 175 of the patients, so far as I could trace them.

There was also given a list of the last 60 cases I had done with reference to the mortality. I can supplement that by 20 more to-day. So that I can point to 80 consecutive cases of extensive operation, and there has been only one death among them. My first communication was to the Lancashire and Cheshire Branch of the British Medical Association in 1877. It was made about a patient on whom I had performed what may be termed the complete operation, and she was well after about three years. That woman lived for about twenty years, and died, when near 70, without recurrence of cancer anywhere. This paper was termed A Plea for a more free Removal of Cancerous Growths. In it I said:

I can point to several cases of hard scirrhus of the breast in which the whole mamma has been removed which have now remained well for years. There is no mystery about this; all that is required is to take the precaution to cut freely through sound textures. But this is exactly what surgeons will not do. I have over and over again watched a surgeon remove a breast, and have seen him make his incisions so a to

include a little elliptical bit of skin containing the nipple, after which he would proceed to scoop out the breast so as to get nice flaps, which fell easily together and covered in the wound. He never considered that in the brawny dotted skin over the tumour were already implanted the germs of the disease, and that the most dangerous thing he could do was to leave plenty of skin; one of the tissues above all others along which cancer cells make their way, I can safely affirm that in the great majority of operations for cancer which I have seen—and they are not a few—the cancer has not been removed root and branch. When the tumour has been carefully looked at after the excitement of the operation was over, there has generally been found some spot where everyone wished that a bit more had been taken away. Not long ago I was looking over some old illuminated manuscripts. Among other pictures of the tortures inflicted upon the faithful there was one in which a horrid wretch was represented with a great crooked knife mangling the body of a devout female. He had slashed off one breast, the former site of which was represented by a great bleeding chasm, while with one hand he was clutching the remaining organ, and with the other was preparing with one sweep to lop it off also. I could not help wishing that I had this picture to hang up in the students' room at our school; not in *terrorem*, but as my idea of what most amputations of the breast ought to be: not a niggling half-and-half digging out of the tumour, but a radical sweeping away of the whole mamma, skin and all. I have a very strong notion that it would not be a bad plan, when removing a breast, always to clear out the axillary glands, whether we can find anything wrong with them or not, and I mean to try it. At present we only remove the glands when we can distinctly feel them enlarged, and that frequently through a great deal of fat and a thick skin. How do we know they are not already infected? And then the patient disappears, and does not return till the axilla is full, and the great vessels are so mixed up with the disease that we despair about being able to clear everything out, and so leave her alone..... By abusing the lay mind of the unreasonable dread of operation the public will come to regard an operation for cancer in the same light as they do an operation for stone or for cataract—a bad job at the best, they will say, but still with a prospect of ultimate cure. They will come to us when their cancers are small and request us to remove them, and, if we could only extirpate them freely, what might we not expect? Have you ever imagined what the results would be if all cancers were thoroughly excised when they were no bigger than peas?

These words were written nearly a quarter of a century ago, and I assert that they contain the principles upon which the present operations for cancer are based—principles which existed, therefore, long before Dr. Halsted wrote a word on the subject. I proceeded at once to put my views as to the axillary glands into practice, and in August, 1882, I read a paper before the British Medical Association at Worcester on Free Removal of Mammary Cancer with Extirpation of the Axillary Glands as a Necessary Accompaniment, and I gave the results of 46 cases. I said:

In the present paper a principal object is to advocate the removal of the axillary glands as well as the breast in all cases, whether we can feel them enlarged or not; in fact, to make the clearing out of the axilla a necessary part of the operation for removal of the breast. I have been quietly practising this for three or four years, having been driven to the conclusion that it was the right thing to do by discovering that even in those cases where certain glands could be distinctly felt to be enlarged when the axilla was opened small ones also were discovered which, although most palpably affected, were quite incapable of being made out from the outside.

Five years passed on, during which I worked on the lines mentioned, and in 1887 I read a paper to the Harveian Society of London, entitled Extirpation of the Axillary Glands, a necessary accompaniment of Removal of the Breast for Cancer, with details of 82 cases. In addition to the points I had urged in my previous papers, I added some others which had struck me.

Having cleared out the breast, I do not cut it adrift, but leave it suspended by its upper and outer end. I next open the axilla, and in so doing carry away a continuous mass of tissue upwards from the mamma, with the object of including in it all the lymphatic vessels going from the heart to the glands. This is an important step, and should never be neglected. Is it necessary to divide the pectoralis major? Very seldom, I think. I imagine that out of my 82 recorded cases it was not done more than two or three times. If the floor of the axilla has been widely opened, and if the arm is properly drawn up, you can feel with your finger quite well up to the clavicle and pinch or pull down glands from the very top of the cavity. Dividing the pectoral muscle adds somewhat to the tediousness of the operation, but that is all.

At the meeting a number of distinguished London surgeons honoured me with their presence, at which I was very much pleased, but, when the discussion on my paper came on, I found to my grievous disappointment that, with a single exception, they denounced it as one man. I can prove this by a reference to the *International Journal of Medical Sciences* (1888) in which is a very important paper by Mr. Samuel Cross, in which he said:

In the remarks on Banks's paper at the Harveian Society in the spring of 1887, in which that surgeon reported 82 cases of extirpation of the breast along with the axillary contents, Bryant and Butlin questioned whether the complete operation afforded better results than the incomplete; and Butlin, Pick, and Cripps held that clearing out the axilla in every case was neither necessary or judicious, although Pick asserted that the skin should be freely removed. Owen was the only member who gave cordial support to the procedure urged by Banks, the prevailing

opinion being that the operation was attended with an excessive mortality, and did not lessen the risk of recurrence. Mr. Thomas Smith, by the way, thought that the patients lived longest who had never been operated upon, and that it was a question for consideration whether a local excretion of cancer did not render patients less liable to constitutional disease.

It need hardly be said that after this very disappointing reception I returned to Liverpool much downcast. But I felt certain that the principles I had been advocating were sound, and so, far from giving them up, I pushed them still further. I had noticed the great liability of the cancerous material to invade the skin and subcutaneous tissues, coming in small nodules (which I have long demonstrated to students under the name of "small shot") at variable distances from the cicatrix. Both Gross and myself had long previously urged the removal of the fascia covering the pectoralis major, so I now proceeded to do this more carefully than ever, and to remove a wider area of skin. To gain some covering, however, for the great bare area, I began to loosen the skin and fat in the immediate neighbourhood by the process of undercutting it. Finally Thiersch's skin grafting came in, and has proved very valuable.

I wrote nothing about the matter for thirteen years, but continued operating upon the same lines until 1900, when in the Lettsomian Lectures I published my experiences and views regarding the whole question of cancer of the breast.

Concerning the proceeding which I have all along advocated, there is nothing mysterious, nor for the moment do I claim any originality for it, for others have done it as well as I. All the merit I claim is that I have been consistent in its use, and have shown, by a large number of collected cases, what its value as a means of cure really is. The first point is to abandon all formal incisions—angular, triangular, or multiangular. Draw a circular "ring fence" with a knife outside the apparent limits of the mamma if the tumour is central, and if it is lateral the "ring fence" should bulge well beyond the tumour. Then slip the blade on the flat beneath the skin for about 2 inches and undercut it all round. Let the skin be well pulled back at successive points by an assistant, and where the undercutting stops there the removal begins, so that one gets a long way beyond the rough limits of the mamma proper. Then remove the mamma and its contained tumour. One great advantage of the undercutting (which I began on an extensive scale in 1877 to get skin to replace the large areas thereof, which I began to remove) is that one can get the fascia off the muscle as far inwards as the sternum, down to the ribs and almost up to the clavicle. This careful removal of the pectoral fascia was one of the points dwelt upon by Dr. Gross and myself very many years ago, and, if properly done, does away with all necessity for removing the great pectoral muscle. The axilla should now be opened by an incision just under the edge of the great pectoral, and the upper end of the mamma with the whole contents of the cavity (as far as they can be held together) car-

ried away in one mass. The importance of this lies in the fact that recurrence in the axilla takes place, for the most part, not in glands that are left behind, but in fragments of lymphatic vessels. I am convinced of this. The clearing out of the axilla requires patience. A dissecting tool, two long strong pairs of dissecting forceps, and a pair of curved probe-pointed scissors are all that is wanted. We are recommended, in the directions for Halsted's operation, to use the "sharpest possible knife" for this purpose. My recommendation is not to put any knives at all into the axilla; neither the sharpest nor the bluntest possible. With the tools mentioned, by a combined process of teasing out and snipping, the space can be cleared of everything it contains and the digitations of the serratus magnus stripped of their covering. If there should be felt any suspicion that the very top of the axilla has not been reached, then the pectoralis major can be divided as far as is necessary to enable this to be done. Its fibres can be pulled together by buried sutures of catgut. As for the great area denuded of cover, it is wonderful how a covering can be drawn together by strong

wire sutures, if only the undercutting has been sufficiently free. Thiersch's grafting, either done at once or subsequently, will fill up the rest of the gap.

In the JOURNAL of December 8th, 1898, is a very elaborate description of Halsted's operation. It contains fifteen paragraphs, with very minute details, down to the grasping of the tumour with the left hand and the scrupulous dissection of axilla with "the sharpest possible knife." The only points of difference between this operation and that which I have just described which are in any way fundamental consist in the removal of the sternal part of the pectoralis major and the division of the pectoralis minor with reflection of that muscle. This is described as necessary for the complete removal of the glands in the

Fig. 4.—Clearing the axilla.

axilla. Now, from this view I beg to express my absolute dissent. It is not necessary. If the arm be well manipulated and the great pectoral well pulled upwards and inwards, the very topmost point of the axilla can be cleaned out; and if the surgeon feels any doubt on the matter, division and subsequent suture of the great pectoral is all that is needed. But Dr. Halsted goes further than this, and in the *Transactions of the American Surgical Association* relates that he now almost invariably cleans out the supraclavicular region, and that our rule ought to be, "Operate on the neck in every case." For this purpose, however, he does not divide the clavicle as he did five or six years ago; but if it were very desirable to remove the supraclavicular contents in one piece with the axillary contents and breast, one need not hesitate. He considers the neck involvement is no more serious than the axillary, and that the neck can be cleaned out just as thoroughly as the axilla. He believes that in the near future we shall remove the mediastinal contents at some of our primary operations. At this rate we shall no doubt

after a few years have from him a proposition to remove the bronchial glands as well. As Mr. Edmund Owen once said, parodying the French general's criticism on the charge of the Light Brigade "*C'est magnifique, mais ce n'est pas la chirurgie.*" Now, concerning the statement that clearing out the neck glands runs on all-fours with clearing out the axillary ones, I quite deny that it does so, inasmuch as I have not a single case in which a patient has lived any time after having the neck glands removed. No man in this country has had more experience of the pathology of cancer than Mr. Plimmer, and he stated to me that he had never known anybody live long after distinct invasion of the supraclavicular glands. If the neck is to be cleared out as a routine performance, let us consider what will happen to a big-breasted, fat woman, with a weak heart. There will be a serious job to start with in the mere removal of the breast and tumour; then comes the digging out of the axilla, next the division of the clavicle in a bad case, and finally a delicate dissection among the great vessels at the root of the neck. I venture to prophesy that, if this procedure is adopted, there will be a great many more people die from it than will ever be saved from cancer—only their deaths will never be heard of in journals. The powers of endurance of the human body have a limit after all.

With regard to the work of Stiles and Heidenhain (that of Rotter I do not know), I desire to record my respect for these gentlemen, and to express my admiration for their work. That work, however, relates chiefly to the microscopic anatomy of the breast and neighbouring parts, and as anatomical research is worthy of all praise. But I do not wish to detract, and I do not detract, from the value of that work, when I deny that it has any bearing upon the practical doings of the surgeon. We do not operate with microscopes. Let anyone examine the wonderful collection of dissections of the lymphatics in the Musée

Orfila at Paris, and he will see at once that everything that can aid him in practical surgery is there displayed. That the intervals between the macroscopic lymphatic vessels there shown were filled with microscopic ones was very evident, and it remained for Stiles and Heidenhain to demonstrate their presence. But that made no difference in the manner in which the neck and axilla had to be cleaned out surgically. It is the same procedure now that it was before they wrote a word about the microscopic lymphatics. Then where does this work come in practically? It might do so if it were asserted that after free removal of the skin, fat, and super-pectoral fascia, the disease returned frequently in the great pectoral muscle or the tissues between it and the small one. But here, while willingly acknowledging the labours of Stiles and Heidenhain in their own line, I must assert my right, on the ground of a long practical experience, to insist emphatically that it does not return in these sites; and I again call to my aid the unrivalled experience of Mr. Plimmer at the Cancer Hospital, who quite coincides in this view.

Having now made several strong statements it behoves me

to show that at all events these statements contain the views of other persons besides myself. The best way will be to cite the opinions of a man whose practical and literary knowledge of the subject of cancer will not, I think, be questioned by any surgeon, namely, Dr. Coley, of New York. He has written the articles on cancer in the *Twentieth Century Practice of Medicine* and in the *Reference Handbook of the Medical Sciences*, and I take the liberty of quoting from them:

While many who advocate the extensive operation prefer to credit their conversion to the histological researches of Heidenhain, we must in fairness acknowledge that this great change has been largely due to the earlier clinical works and the superior results of Moore, Banks, and Gross. Banks's statistics are the largest and most complete yet published.

After giving a very careful analysis of the statistics, he says:

This shows the great importance of following up the patients for long periods of time, and makes the statistics of Banks by far the most important data we now possess upon which to base a true estimate of the curative value of surgical treatment of cancer of the breast. They certainly justify his conclusion that the plan of taking percentages in small numbers of cases ought to be abolished, and, furthermore, that the three-

year-cure system, too, must be abolished, if people are to be instructed fairly.....I have not yet been convinced of the wisdom or value of removal of the supraclavicular glands in operations for cancer of the breast. Statistics.....which equal those of Halsted would go to show that as good results can be obtained without this step.....In order to obtain better results I think it far more important to remove wider areas of skin and more of the adjacent fat, for we find that recurrence takes place in the skin or fat a dozen times to one in the pectoral muscles.

To conclude I personally am strongly of opinion that for the most part operations for cancer generally have gone about as far as they are likely to do good. As yet the breast is the organ where the best results have been obtained. The operations on the upper jaw, the testicle, and the stomach have hitherto been utterly disastrous as regards preventing the return of the disease, and to say that there are entirely "new lines" still to

Fig. 5.—The denuded area covered.

be discovered upon which new operations are to be founded is what I will not believe. The rectum and the great intestine have yielded, next to the breast, the best hopes of cure, but if the true histories of all the cases of cancer of the tongue of any size and moment, were capable of being told, they would constitute a very sad volume in surgery. I believe that the real advance is to be obtained by surgical teachers everywhere impressing upon their pupils the paramount necessity for recognising the earliest possible appearances of cancer, in the hope that this view may ultimately reach the public and induce them to seek medical help early in place of waiting, as they too often do now, lest their malady is past praying for. I repeat again the words I said in 1877: "Have you ever imagined what the results would be if all cancers were thoroughly excised when they were no bigger than peas?" But if this happy consummation is to be reached it will not be by performing tremendous operations upon practically hopeless cases. I have heard it argued that such cases should be done, even if there was a hundred-to-one chance for the patient. I entirely refuse

to subscribe to this doctrine. It is obvious that, for one who recovers, ninety-nine perish from the operation or from speedy return of the disease. The learning of the disasters to these ninety-nine probably affects the minds of several hundreds of friends and relations who hear about their unhappy fate. When a woman who knows all this discovers a lump in her breast, the unfortunate ends of persons whom she has known or heard about occur to her, and so frighten her that she conceals her complaint until she cannot do so any longer, and when she sees a surgeon it is too late for him to do any good. The operation is prejudiced in the eyes of the public, and for the one person in a hundred who escapes by some desperate operation, there are many who lose their lives by being terrified out of an early operation, which alone could have saved them. So I venture to recall once more the wise words of William Hunter, that the operation ought not to be undertaken without great probability of cure, as we run the risk of a general prejudice against the operation, which may so affright others that they will not run the hazard, though their disease may be really curable.

NOTE AND REFERENCE.

¹ As illustrating William Hunter's quaint ways of speech, the following is his valedictory address at the end of the session: "Now, gentlemen, I have finished ye course and wish you all the success in ye world. It has given me a great deal of pleasure to attend to a number of gentlemen, who have made it their study from ye beginning to apply with assiduity to what was going on. Whatever you may think of it, I am always studying your good, and it is this alone which induces me to give lectures. I have now finished twenty years of lectures, and however, as I presume I am still approved of, I propose twenty years more to begin next October, and, after this is over, I propose to settle in ye world and take me a wife."

² *Tumours of the Breast*, 1880.

A Clinical Lecture

ON

OÖPHORECTOMY IN THE TREATMENT

OF

CANCER OF THE BREAST.

Delivered at St. Bartholomew's Hospital, Wednesday, November 20th, 1901.

By H. T. BUTLIN, D.C.L., F.R.C.S.,
Surgeon to St. Bartholomew's Hospital.

GENTLEMEN,—I have for some time past had it in my mind to lecture on the removal of the ovaries for malignant disease of the breast. I thought I would do so somewhat later in this year or in 1902, but one or two circumstances which have arisen lead me to lecture on it almost before I am prepared to do so. However, the only want in the preparation is that I shall not be able to tell you the story quite so concisely as I might have done if I had had a longer time to put it together. As to the facts and opinions they are not likely to be altered in another three months.

In the first place, some of you may remember a woman whom I showed in consultation some months ago, who was suffering from slight recurrence, not very severe, after amputation of the breast and removal of the glands. The recurrence was scattered, and was not convenient for local operation. She was about 40 years of age, still menstruating, and she had no apparent disease of bone or viscera. I brought her down to consultation, and placed before my colleagues the question of removal of the ovaries and tubes in accordance with the treatment which has been employed from time to time for several years past. Various opinions were expressed, but nobody was hopeful of the success of the operation. Still, many of my colleagues thought it was a proper case on which to try the experiment, and, generally speaking, we agreed that the operation should be practised in that case. My colleague, Mr. Lockwood, removed the ovaries and tubes. I have just heard the end of that case, and I will tell it to you presently.

Then a lady came to me last week. I operated upon her and her sister within a few months of each other for cancer of the breast, and she came to ask me (not for herself, but for some

other lady whom I have never seen, and who had been operated on by another surgeon), whether I thought it would be a good thing that her ovaries should be removed in order to prevent the recurrence of the disease, of which there is, as yet, no sign. I told her I was going to lecture on the question shortly. She wished to know whether the lecture would be published, and, if so, where, as she would like to read it, because she would like to know my opinion on the subject. Regarding her friend. I advised her for the moment not to have anything done. I tell you that because it shows that the matter has now become lay property, and people are beginning to talk of removal of the ovaries in certain cases of cancer of the breast.

At the time I showed my case at consultation I suppose everybody had in mind that the female organs of generation are related to each other, related, that is, in such a way that, for instance, when a woman is menstruating, a tumour of the breast, such as a cyst or a mammary glandular tumour, is said, and I daresay with truth, to get larger for the time being. And it is well known that during menstruation the breasts are apt to become painful and enlarged. So the relation between the ovaries and the breasts is a matter which is admitted by medical people, and is believed by women in all classes of society. But in May, 1896, Dr. George Beatson, of Glasgow, who is attached to the Cancer Hospital there, read a paper before the Edinburgh Medico-Chirurgical Society on the Treatment of Inoperable Cases of Carcinoma of the Mamma: Suggestions for a New Method of Treatment: with Illustrative Cases. Dr. Beatson gave an account of two women to the Society, both of whom he showed. They had been treated by his new method, and eleven months after operation one of them appeared to be perfectly well of an absolutely inoperable condition of cancer.

Before I speak of the cases and of other cases in which the operation and thyroid treatment have been employed, I would like to draw your attention for a moment to his theory. The theory was not a mere general idea that the breast exercises a sort of influence on the generative organs, but he had certain definite views with regard to the nature of cancer of the breast which differed very considerably from those which hold the field at the present time. (I speak of the parasitic origin of cancer, which many people believe in, although it has not yet been proved.) Dr. Beatson considered that cancer of the breast is due to proliferation of the epithelium of the breast. It is a "local proliferation of epithelium," and he said when the cells had proliferated and increased in number, and could not escape from the acini or make their way from the ducts, they gradually swarmed out into the surrounding tissues, and produced the microscopic structure of cancer. He said there was no parasitic character about the disorder, and no special parasite had been seen, because there is no parasite in cancer of the breast. Taking this view of the nature of cancer, and believing it to be a mere proliferation and overflow, so to speak, of the epithelium of the breast, he began to consider what might control or alter this proliferation. I may tell you that this matter was before his mind for many years before he performed his operation. He remembered that in ordinary lactation there is a similar proliferation of epithelium, and the epithelium, as soon as it is formed, instead of filling up the acini and then overflowing into the surrounding connective tissue, undergoes rapid fatty degeneration and is passed away as milk, and so no harm comes to the woman, and he remembered that at such times women do not menstruate. So he thought, if a woman has local proliferation of epithelium of the breast not during the period of lactation, and we suppress the action of the ovaries, it seems possible that we might induce a fatty degeneration in this proliferating and swarming epithelium which might spoil it, so that it would be no longer harmful to the individual. So he proposed to himself that he would, some time or other, if he could find a suitable case, take out the ovaries and tubes, so as to be quite sure that no ovarian tissue was left. That is, in effect, Dr. Beatson's theory; and it is well, before you perform or talk about this operation, that you should know what was in his mind. I wondered, until I read his paper—which is a very good one indeed—what had first led him to perform the operation of oöphorectomy for cancer of the breast. I found it was the result of long study and of speculative thought, and not a mere matter of chance. He considered the matter for a number of years, and worked up a good deal of information in