HYDROA GESTATIONIS DUE TO STAPHYLOCOCCUS ALBUS.

BY

NATHANIEL W. HOLMES, and W. BULLOCH, M.D.,
L.R.C.P., L.R.C.S., L.M. Edin., Bacteriologist to the London
late Government Medical Officer, Hospital.
Australia.

THE following case is of interest owing to its occurrence at three consecutive confinements, and to the infrequency that one comes in contact with such cases, also the etiology as revealed by bacteriological examination by Dr. Bulloch.

Mrs. O., aged 26 years, was confined of her third child on October 28th, 1901, the following days she complained of severe itching in the palms of the hands and soles of the feet, extending up the forearms and generally over the whole body. There was redness of the palms of the hands and soles of the feet, and a thick elevated papular and confluent rash the size of peas with hardened bases on the backs of the hands, wrists, and on the flexor and extensor parts of the forearms, also on the knees. The rash on the thighs, chest, back, and other parts of the body was either isolated, confluent and semicircular, or clustered in round patches the size of a florin. Temperature normal. She informed me (N. W. H.) that she had suffered from the same thing at each of her two previous confinements and was most anxious to know what could be the cause of it.

The following day, October 29th, bullæ formed upon the wrists and forearms from the size of a pea to that of walnuts, fresh ones occurring daily here and there over the body and legs. Upon inquiry I could obtain no evidence pointing to syphilis in the family, and as the case resembled pemphigus (for which she was treated at a skin hospital for three or four months after the first confinement) some interesting cases of which were reported by Mr. Buxton Shillitoe,¹ and Drs. Hadley and Bulloch,² in which a diplococcus was found, I sought to investigate it further. At the age of 13 years she had a very bad poisoned foot and leg due to an injury to the sole of the right foot by a brass rivet from her boot, which has left a scar on the anterior surface of the leg, and about the same time she had diphtheria.

November 6th.—The pimples appeared on the chin and sides of the face, followed by numerous bullæ daily.

November 8th.—By the kind co-operation of Dr. Bulloch a bacteriological examination was made of the fluid from some recent bullæ, and a pure culture of staphylococcus albus was found.

November 12th.—Fresh bullæ appeared on the forearms, and on the 14th some became pustular, brown scars being generally left behind.

The patient improved daily, and the infant, which continued taking the mother's milk from birth, was quite free from any rash. The treatment consisted in the local application of lot. picis liq., internally a saline mixture with liq. arsenicalis, and latterly sulphur tablets only were prescribed. There has been no return of the disease for the past six months, and it will be interesting to note if there should be any recurrence.

REFERENCES.

¹ *Lancet*, November 26th, 1898, p. 1397. ² *Lancet*, May 6th, 1899, p. 1219.

NOTE ON A METHOD OF QUANTITATIVELY ESTIMATING THE PHAGOCYTTIC POWER OF THE LEUCOCYTES OF THE BLOOD.

By MAJOR W. B. LEISHMAN, M.B., C.M., R.A.M.C.,
Assistant Professor of Pathology, Army Medical School, Netley.
(From the Pathological Laboratory, Netley.)

THE question as to the relative value of the parts played by the fluid constituents and the cellular elements of the blood respectively in combating an attempted invasion of the system by pathogenic micro-organisms is still the subject of much experiment and discussion. I have thought therefore that the following preliminary note on a method of approximately estimating *in vitro* the part played by the leucocytes in this struggle for mastery may be of interest.

In the recent work which Professor Wright and others in this laboratory have been doing in the estimation of the bactericidal power of the fluids of the blood, a striking feature has been the extremely feeble power that these fluid constituents possess for the destruction of the common staphylococcus pyogenes, the spirillum of cholera, the micrococcus of Malta fever, and the bacilli of anthrax and plague, in comparison with that displayed for the bacillus of typhoid. Even when imitating the conditions of life as closely as possible *in vitro* with regard to temperature, anaërobiosis, the use of fresh blood serum as a diluting medium for the culture em-

ployed, etc., the results have proved almost completely negative; in other words, the blood fluids alone cannot be shown by experiments *in vitro* to have more than a trace of bactericidal power against these organisms. In the methods referred to the leucocytes have mostly been excluded by clotting or centrifugalisation and the clear serum or plasma used, so that the part played by these corpuscles, if any, has not been taken into account. The well-known phagocytic power of these white cells, though universally acknowledged, is allotted a very different value by the French and German schools respectively, and the method described below seems to promise a possible means of elucidating some of the many difficulties in determining the share taken by these phagocytes, especially if used as a supplement to an estimation of the bactericidal power of the blood fluids such as that of Professor Wright.¹ In any case the method has, I think, a certain value in the ease with which, *in vitro*, beautiful examples of phagocytosis may be demonstrated to and carried out by a class without the necessity for employing animals such as the frog, guinea-pig, or rabbit.

While the power of phagocytosis (the ingestion of living or dead germs or of particles of inorganic matter) is possessed in common by all the white cells of the blood except, with very rare exceptions, by the lymphocytes; the polynuclear or fine-grained eosinophilous cells are endowed to the greatest extent, and I have, partly on this account, partly because of their preponderating number, confined myself to determining their absorptive capacity, leaving out of account that possessed by the mononuclears and coarse-grained eosinophiles.

Briefly, the method aims at enumerating the number of bacilli or cocci phagocytized within a definite time by the polynuclear cells of the blood under examination, obtaining an average of these values, and comparing the result with that of a control blood, preferably that of the observer, put up under identical conditions.

An emulsion of a portion of an agar culture of the germ whose action is being investigated is first made in a few drops of normal salt solution in a watch glass. Even distribution of the germs is ensured by assiduous stirring with a platinum loop or needle, at intervals of a few minutes, until the emulsion presents an even turbidity. A well-marked but not too dense opacity is aimed at, and the culture used should be a fresh one. A well-polished microscope slide and a couple of perfectly clean cover glasses are prepared. A measured volume of blood is now to be added to and mixed with an equal volume of the emulsion of germs. This may conveniently be done by means of a small capillary pipette, fitted with an india-rubber teat and having a mark near the point corresponding to the desired volume. A little practice enables one to judge of the volume necessary to spread itself out in a thin even film between cover glass and slide; it is, approximately, 5 c.mm. A small drop of blood is obtained in the usual way by means of a needle prick, and one volume, or 5 c.mm., is at once taken up by the pipette and allowed to run slightly up to the tube so as to admit an air bubble to serve as an index; the pipette is then dipped in the emulsion of germs, and a similar volume (5 c.mm.) admitted. Both volumes are now blown out into a clean watch glass or on to a slide, and the blood and emulsion well mixed by stirring and drawing in and out of the pipette. One volume of the mixture is now taken up in the pipette and gently blown out on one end of the prepared slide and at once covered by a cover glass, very gentle pressure with the forceps being used, if necessary, to ensure the film spreading evenly to the whole circumference of the circle. A precisely similar preparation is then made of the control blood from the observer's finger, the drop being placed and covered at the opposite end of the same glass slide. The slide is now placed in a moist chamber, and left in an incubator at 37° C. for half an hour. As a moist chamber a Petri dish with a few layers of damp filter paper at the bottom may be used, the slide being placed in it, resting on a couple of wooden matches. The vapour condenses on the under surface of the slide and the walls of the dish, but not on the upper surface of the slide, which remains dry. A small amount of drying takes place at the edge of the cover glass, but the blood under it remains liquid for hours, and the phagocytes, by reason of the temperature and the close apposition of the two glass surfaces, have every facility for

The results of the phagocytic estimations may be more readily followed by reference to the accompanying chart, in which the average number of staphylococci phagocytied by the patient's leucocytes in half an hour is contrasted with that of the control blood—my own. The principle of the chart is that the daily figure of the phagocytic power of the control blood is taken as zero, and the difference between

this figure and that of the blood under examination is shown above or below this line, according to the numerical excess or deficiency in the average of germs phagocytied by each leucocyte.

It is too early to be able to say whether the temporary benefit which followed the inoculation is likely to be permanent; but the point which I wish to bring out is that the patient's phagocytic power, as gauged by the control blood, has varied in direct proportion to the amelioration of his symptoms.

Prior to the date of inoculation I made several examinations of his blood, and found its phagocytic power always lower than the control, but these have not been entered in the chart, as no exact numerical count was made on these occasions. For six days after inoculation it will be noted that his phagocytic power remained considerably lower than the control; during this time he was still having fresh crops of small boils and pimples on his neck. From the seventh to the tenth day he was free from boils, for the first time for many weeks, and a corresponding rise is seen in his curve. From the eleventh to the thirteenth day a few fresh pimples appeared, which gave him less annoyance than those he usually suffered from, and a similar outcrop occurred towards the end of the period of observation. On both these occasions a drop will be noted in his curve below the level of the control blood.

Without insisting on the uniformity or importance of the phagocytic estimation in this case, which has only been selected as an example of the method, the results appear sufficiently striking to warrant the hope that it may be of service in the investigation of immunity when combined with an estimation of the bactericidal power of the blood fluids.

Since the above note was written I have had the opportunity of examining the phagocytic reaction in another case which was treated eight months ago by Dr. Wright with injections of staphylococcus vaccine for boils from which the patient had suffered for eight years, and in whose case a permanent cure appears to have been effected. The phagocytic power of the polynuclears was found to be markedly greater than that of the control blood when tested with the strain of staphylococcus, which was originally isolated from one of his boils, and from which the vaccine used in his case was prepared, while the reaction was slightly higher than the control when tested with another variety of staphylococcus.

REFERENCES.

¹ *Lancet*, December 1st, 1900, March 2nd, 1901, and June 1st, 1901. ² *BRITISH MEDICAL JOURNAL*, September 21st, 1901. ³ *Lancet*, June 1st, 1901. ⁴ Proceedings of the Pathological Society, *Lancet*, March 23rd, 1901.

EPIDEMIC CATARRHAL JAUNDICE.

By ELIOT CURWEN, M.B., B.C.CANTAB.,
Late Physician to the Pekin Hospital.

In view of the present epidemic outbreak of jaundice in Derbyshire, so well described by Drs. Peck and Kershaw, the following brief notes of the first cases in similar epidemics in Pekin and Tientsin during the winters of 1898-9 and 1899-1900 may not be without interest. I have notes of only 44 of the cases seen during the first two months of the epidemic, which, beginning towards the end of October, 1898, lasted well into the spring of the next year; the following winter the epidemic returned, and presented the same features as in the previous year.

The first case seen was on October 22nd, and for some months after this few days passed without one, two, or three patients coming to the hospital for treatment. Their ages varied from 17 to 60, and seven times as many men as women applied for treatment. The attack from which one of my students suffered was typical of the series.

On October 27th and 28th he told me he felt weak and lethargic. On the evening of the second day he returned to his home feeling sleepy, ate two pears, had pain in the abdomen, and felt sick. At night he was very hot, complained of severe pain in his temples, and his conjunctivæ were, he said, suffused. During the next two days his temperature kept up between 102° and 102.4°. He felt very weak and ill, and the pain in his temples continued. The administration of calomel (gr. ii.) resulted in five dark green motions. On November 1st his temperature was down to 96° in the morning, and the conjunctivæ were slightly yellow for the first time; he was still very weak, and had a pulse of 60; there was no tenderness over the liver, or any abdominal discomfort; the tongue was slightly

furred. The colour of the conjunctivæ became more marked by November 4th, when he began to complain of itching of the skin; from this date his symptoms gradually improved; ten days later, however, the conjunctivæ were still of a pale lemon colour.

Most of the patients complained of great weakness and of some shortness of breath; there was generally anorexia, a feeling of abdominal distension, and nausea, and constipation with pale motions was the rule; in 4 cases definite abdominal pain was complained of, but from only 3 did I hear of actual vomiting; diarrhoea occasionally occurred. Another prominent symptom, noted in two-thirds of the cases, was pain in the temples, and this in some cases was described as very severe; it was often accompanied by drowsiness, and in 2 cases by vertigo. More than two-thirds of the patients told me that for the first few days of illness, varying from one to fourteen, they suffered from some degree of fever with shivering, and that as a rule the icteric tint of the conjunctivæ was first noticed the day, or the day after, the fever disappeared. In 5 cases the colour of the conjunctivæ was noted as deep, but in the majority of cases it was a pale straw or lemon colour, and not dark enough to be noticed in the yellow skins of the Chinese; the length of time it lasted was very variable. Only 4 patients complained of pruritus. In 2 cases the liver was noted as slightly enlarged and tender; no albumen was found in any urine examined, but Martin's and Gmelin's reactions were obtained in each case. The pulse was rapid—from 80 to 128—in each patient, with the exception of the case narrated above. One man, aged 60, whose fever had lasted ten days, told me that about the time his conjunctivæ became yellow he began to suffer from pains in the muscles of the right arm and in his two ulnar bones; the muscles and bones were still tender on deep pressure when I first saw him five days later.

The affection cannot be very infectious, for I never saw two patients from the same house, and one of my servants who was attacked did not communicate the disease to the other men who slept on the same brick bed; only one patient told me that he had neighbours suffering from the complaint. Some of my patients were severely ill for some days, and, on account of their high fever, weakness, and headache, were thought by their friends to be suffering from typhus fever; others, on the other hand, were affected so slightly that the febrile symptoms, if they existed, passed unnoticed, and they did not know there was anything the matter with them till I handed them a looking glass and told them to look at their eyes.

ENTERIC FEVER IN THE INOCULATED.

By MAJOR C. BIRT, R.A.M.C.,
Harrismith, South Africa.

THE mortality of 947 uninoculated individuals who have suffered from enteric fever in Harrismith, South Africa, from September, 1900, to September, 1901, has been 14.25 per cent. There were 135 deaths among them.

During the same period 263 persons who had been inoculated with typhoid vaccine, for the most part 6 to 18 months previously, contracted this disease. Eighteen of these cases proved fatal. This is equivalent to a death-rate of 6.8 per cent.

The tabulated results of an analysis of 317 charts of the not inoculated and of 151 charts of the inoculated are given below.

	Uninoculated.	Inoculated.
Mortality	14.25 per cent.	6.8 per cent.
Average duration of pyrexia	28 days	15 days
Average maximum temperature recorded...	103.7°	102.9°
Percentage of cases in which relapses occurred	24.0 per cent.	6.0 per cent.
Percentage of cases with constipation	19.0 "	33.0 "
In cases with diarrhoea, average maximum number of evacuations per diem	4.4 "	2.6 "