far as I know the "dot" in this case was permanent. The patient asserted that in looking at a line he observed the portion looked at to be thicker (metamorphopsia). This phenomenon has been observed in a few of the recorded cases. It was absent in the case I now relate. Many years ago a case came under my notice of a girl³ who suffered from retinitis in consequence of receiving the sun, on a hot summer's day, full on her eye from a reflector which a boy passing her was using. The eye recovered well.

The instance related here is of interest as both eyes were affected, and the patient, an unusually intelligent man and an excellent observer, has been able to give a good account of his case. Here the ophthalmoscopic signs were practically nil; in some of the cases recorded fine changes have been noted at the macula, whilst in others similar indefinite alterations to those referred to in my present case have been observed; in others again no abnormal appearances have been mentioned. The prognosis appears to be favourable in these cases, vision, for practical purposes, being restored, but in many a scotoma, more or less distinct, has been left permanently. The cases illustrate the utter uselessness of the "glasses" too frequently used in observing eclipses.

REFERENCES.

¹ On Blinding of the Retina by Direct Sunlight. ² *Ophthalmic Review*, 1884, p. 72. ³ *Ibid.*, 1883, p. 141.

A REPORT OF NINE CASES IN WHICH HAAB'S MAGNET WAS USED FOR THE EXTRACTION OF FOREIGN BODIES FROM THE EYE.

BY

W. T. HOLMES SPICER, and A. F. MACCALLAN,
F.R.C.S., F.R.C.S.

Ophthalmic Surgeon, St. Bartholomew's Hospital; Surgeon, Royal London Ophthalmic Hospital, Moorfields. Senior House-Surgeon, Royal London Ophthalmic Hospital, Moorfields.

THE cases reported here were under the care of Mr. Holmes Spicer at the Moorfields Eye Hospital. Our thanks are due to Mr. Mackenzie Davidson for his valuable help in localising the foreign bodies for us before operation.

Two of the cases resulted in recovery with almost perfect vision, $\frac{5}{6}$ with correcting glasses; one with vision of $\frac{5}{6}$, although at the time of admission there was iritis with hypopyon; two others had vision of $\frac{5}{6}$, with a possibility of further improvement by needling of opaque lens capsule; all these had perfect projection. In two other cases the eyes were in a condition of acute inflammation at the time of the operation, and in one case so much destruction had been produced by the large size of the piece of steel that the prospect of saving the eye was from the beginning hopeless. These eyes were lost.

CASE I.—T. T., aged 27. On May 23rd, 1900, the left globe was penetrated by a chip of steel from a lathe. He had had very little pain or inconvenience since the accident. When seen at Moorfields on July 1st he was found to have a corneal scar and a traumatic cataract. The pupil was dilated under atropine, and the eye was fairly quiet. No red reflex was obtained from the fundus. V.L.=hand movements. A foreign body was localised by the aid of x rays in the vitreous, 1.3 cm. behind the centre of the cornea.

On July 4th the extraction of the foreign body was attempted. The patient's head being put on a level with, and at a distance of a few feet from, Haab's magnet, the electric current was switched on gradually. At the same time the patient's eye was approached to the terminal, and he at once experienced pain. The foreign body was seen to come forward through the lens, and to take up a position on the lower pupillary border of the iris.

The patient, who had previously been sitting on a stool in front of the magnet, was transferred to the operating table. At this point the foreign body disappeared from view. An incision was made with a keratome at the lowest part of the corneo-scleral ring. The curved terminal of the horseshoe magnet was introduced into the anterior chamber. The foreign body was drawn over the pupillary border of the iris and removed. Some lens matter was evacuated with a curette; a little inflamed vitreous was also removed.

On the patient's discharge, July 15th, the eye was quite quiet, and the remainder of the lens matter was becoming absorbed.

On December 10th Dr. Edgar Stephenson, of Liverpool, under whose care the patient originally was, most kindly sent us a note as to his condition. The pupil was quite clear; the lens matter was all absorbed, but there were some capsular remains in the periphery. V.C. sph. +11 D.= $\frac{5}{6}$; V.C. sph. +15 D.=J.r. The patient has been back at work for some months, and has had no trouble of any kind with his eye. Size of F.B.=6.5 mm. x 2 mm. x 1 mm.

CASE II.—W. N., aged 16. On March 16th, 1900, a piece of steel flew into the patient's left eye from a chisel. He was found to have a scar at the outer part of the cornea, a hypopyon, and iritis. Both cornea and lens had been wounded; there was a dull red fundus reflex only; projection was good. V.=hand movements. On March 20th the hypopyon had disappeared. The foreign body was localised in the vitreous, 1.2 cm. behind the centre of the cornea and 4 mm. to temporal side. A few days later the patient was introduced to Haab's magnet, and the foreign body was withdrawn—without further wound of the lens—into the anterior chamber, through the lower periphery of the iris. Probably it came through the zonule. A section was made with a keratome, and on inserting a curved terminal of the horseshoe magnet, the foreign body adhered to it and was removed.

When the patient was seen eight months later the condition was as follows: An anterior synechia at the site of the corneal wound, an active pupil, and some anterior and posterior capsular opacity of the lens. After correction of hypermetropia patient could read $\frac{1}{2}$ partly. The patient now follows his trade of carpenter with ease.

CASE III.—W. R., aged 28. A piece of steel penetrated left globe four weeks previous to admission. He had attended another ophthalmic hospital where a small magnet had been tried unsuccessfully and where excision had been recommended. When seen, December 20th, 1899, there was a corneal wound to the inner side, and a nebula just above the centre of the cornea; there were also definite signs of iritis. The lens was clear; there were vitreous opacities, V.L.=fingers at 4 feet. The tension was diminished. On introducing the patient to Haab's magnet, the foreign body was drawn just over the pupillary border of the iris, but fell back again. Owing to failure of the current the small magnet could not be used, and further operation was deferred. Later, the patient was x-rayed, and the foreign body localised at the back of the vitreous. Haab's magnet being applied (January 13th, 1900) the foreign body was drawn into the anterior chamber. After a corneal section had been made with a keratome, the small magnet withdrew the particle. Some lens matter was curetted; it was also necessary to perform a small iridectomy. The patient was seen on January 20th of this year. He was found to have a thick capsular membrane, through which the details of the fundus could not be seen. Projection was perfect. V.L.C. sph. +10 = $\frac{5}{6}$. Field good.

CASE IV.—A. P., aged 23. Came to the out-patient department April 4th, 1900, complaining that a piece of steel had struck his right eye. He thought that nothing remained in it. He said that the fragment was $\frac{1}{2}$ inch long, and had been found on the ground after the accident. On examination there was found to be slight conjunctival injection, a vertical wound in the outer hemisphere of the cornea, and some iritis. The lens was becoming opaque. V.R.=hand movements. The tension was diminished. He was treated as an out-patient until April 7th. In the interval he was x-rayed and the foreign body was localised in the vitreous, probably close to the retina. He was admitted to the hospital, and Haab's magnet being applied, the foreign body was drawn into the anterior chamber. The patient was then anaesthetised; with a Graefe knife a section was made upwards, and the foreign body was extracted with ease by the small magnet. Some lens matter escaped, and some was evacuated with a curette. Iris prolapsing a small iridectomy was done. The patient was seen January 20th, 1901. A shrunken, calcareous capsule was found to be present, which almost entirely blocked the pupil. On either side was seen a faint red fundus reflex. V.R.C. sph. +10 = $\frac{5}{6}$. Size of F.B.=5.5 mm. x 3 mm. x 2 mm.

CASE V.—W. B., aged 42, first came to the hospital on September 9th, 1897, having been struck in his left eye with a piece of steel from a tool ten weeks before. There was then a scar on the cornea, the lower part of the lens was opaque; no details of the fundus of the eye could be seen. The eye remained quiet while he was watched for a period of three months; no operation was thought advisable, as there was no evidence of the presence of a foreign body in the eye, and he ceased to attend. He returned to the hospital in September, 1899. On examination the eye was quiet; the lens was partly opaque; a red reflex was obtained from the upper part of the fundus, but no foreign body was seen. On November 3rd the foreign body was localised by the aid of x rays in the globe, situated on a horizontal plane running 1 mm. below the lower margin of the cornea, and 7 mm. back from the plane of the centre of the cornea, almost in the middle line. On November 5th the condition of the eye was as follows: Some general conjunctivitis; no visible wound; pupil irregular but active; lens opaque; no red fundus reflex.

On introducing the patient to Haab's magnet the foreign body was drawn into the anterior chamber, and removed by the small magnet, after a corneo-scleral section had been made. The patient was discharged at the end of a week, with the eye in a perfectly quiescent condition. On December 19th he was re-admitted. The pupil was irregular, but fairly active. The lens had absorbed, and a dense capsular membrane in the pupillary area remained. No details of the fundus could be seen.

V.L.C. sph. +10 = $\frac{5}{6}$. T.=normal. Projection good. The capsule was needed, and a good central opening made.

The patient continued to attend as an out-patient till February 10th, when his vision C. sph. +13 D.= $\frac{5}{6}$.

CASE VI.—W. M., aged 24, came to the hospital on October 25th, 1899. He had been struck that morning by a piece of steel in the right eye; he did not know how large it was, nor what became of it.

There was general injection of the globe; the cornea was dull all over; there was a wound extending from just above the middle of the cornea downwards and inwards to the edge. There was no anterior chamber, the aqueous still leaking away from the wound, through which the iris was freely prolapsed. The lens was wounded and was becoming opaque. Vision of the wounded eye amounted to perception of light only. The tension was markedly diminished, being about equivalent to T.—3.

On bringing up the patient to Haab's magnet, he complained of intense pain; the cornea became conical, as if pushed forwards by something within. On bringing the eye close to the pole of the magnet, the whole globe was seen to be considerably drawn forwards out of the orbit; the movement was visible to everyone around, and was certainly not less than $\frac{1}{2}$ inch. On examining the eye, a shining piece of metal could be seen in the anterior chamber.

The small magnet was then applied between the lips of the corneal wound, but the piece of metal was too large to extract. The wound was

somewhat enlarged, and the piece of metal removed after some manipulation with the small magnet. Size of F.B. 15×5 mm.

Three days later the anterior chamber was reformed and the blood for the most part absorbed, but the iris was entangled in the wound in its whole length.

On November 11th the eye was still injected and irritable; the iris was entangled in the wound; there was bare perception of light, and the projection was not good. There was no keratitis punctata. The tension of the eye was normal.

Considering the probable extensive damage by the very large piece of metal it was not considered advisable to make any further attempt to save the eye. Accordingly the contents of the sclerotic were eviscerated and a glass globe was inserted. He was discharged on November 21st.

CASE VII.—C.G., aged 25, was admitted two days after the penetration of a piece of steel into his left eye. He had much conjunctival discharge; the cornea was hazy, and there was a small central wound. There was marked iritis. The foreign body was removed on the same day by Haab's magnet by way of the original wound. On the following day, the inflammation having become more acute, the edges of the wound were touched with the galvano-cautery. The eye appeared likely to become a source of danger to its fellow and Mules's operation was accordingly performed a week after the date of injury.

Size of F.B. $= 3.5$ mm. $\times 3$ mm. $\times 1.5$ mm.

CASE VIII.—Charles S., aged 30, was struck in the left eye by a chip of steel from the face of a hammer fourteen days before coming to the hospital. When seen there was much conjunctival injection. In the upper and outer quadrant of the corneo-scleral margin was a wound in which the iris was entangled. The lens was clear. There was much hæmorrhage into the vitreous. V.L. $= \frac{5}{6}$.

At the outer canthus beneath the conjunctiva a small swelling could be seen. The skiagraph which was taken localised the foreign body in the region of the swelling about 6 mm. deep. On the day after his admission, May 12th, 1900, Haab's magnet was applied to the globe without causing any pain; when applied to the outer canthus the patient experienced pain at once, and the conjunctiva bulged forward. The conjunctiva was divided over the region of the swelling, and extraction with the small magnet was attempted without success. On again applying Haab's magnet the foreign body appeared between the lips of the conjunctival wound and was extracted with forceps.

He was discharged from the hospital on May 21st, V.L. $= \frac{5}{6}$, when the condition was: Fundus obscured by vitreous hæmorrhages; iris still somewhat adherent to the corneal wound. The patient was seen on February 23rd, 1901, when V. was found to be $\frac{5}{6}$ partly. There was a large scar in the outermost periphery of the fundus. In this case the foreign body must have passed through the corneo-scleral ring, the ciliary body, the vitreous and the sclerotic, the scar in which is still evident. Size of F.B. $= 8.5$ mm. $\times 5.5$ mm. $\times 3$ mm.

CASE IX.—F.P., aged 20, was struck in the right eye by a piece of steel on April 2nd; he came to the hospital on the following day. There was slight conjunctival injection; the iris was prolapsed through the upper part of a wound in the cornea; the lens was opaque; the pupil was dilated with atropine, and Haab's magnet was applied. Pain was at once experienced, and the foreign body appeared in the middle of the anterior chamber, having apparently travelled up from the lower part of the vitreous. On breaking the circuit the foreign body dropped into the lowest part of the anterior chamber. Haab's magnet was again applied, and the foreign body was drawn out from the original wound. A small prolapse of iris was cut off. Some lens matter and a bead of vitreous escaped during the exit of the foreign body. On the following day a hypopyon was found to be present; this disappeared under treatment by dry heat.

Five days after the operation there was intense chemosis of the conjunctiva; the anterior chamber contained turbid fluid; the tension of the globe was increased. A curette evacuation of broken up lens matter and lymph was performed, and the anterior chamber was washed out with saline solution. In spite of this treatment the dangerous inflammatory condition persisted, and it was considered advisable to remove the eye; excision was accordingly performed.

The above cases may be epitomised as follows:

CASE I.—The lens was opaque previous to operation; it has now entirely absorbed. A correcting sphere brings the vision up to $\frac{5}{6}$.

CASE II.—There are anterior and posterior capsular opacities, which were present previous to operation. They do not appear to be extending. V. $= \frac{5}{6}$. It is to be noted that there was considerable inflammation, iritis, and hypopyon at the time of operation. The foreign body was drawn through the zonule without further injury to the lens.

CASE III.—The lens, which was wounded in the withdrawal of the foreign body, has absorbed, leaving a capsular membrane. V. after correction $= \frac{5}{6}$. The projection is good.

CASE IV.—The lens, which was wounded previous to operation, has become absorbed, leaving a thickened capsule. V. after correction $= \frac{5}{6}$. Projection perfect.

In the two last-mentioned cases it will be a matter for future consideration as to whether or not dissection of the capsule is attempted. In one of them, at any rate, this does not appear to be contraindicated. In the other case the capsule is thick and calcareous, and is adherent to the iris. Here we should not advise further operation.

CASE V.—The lens was opaque previous to operation. It has absorbed leaving a dense capsular membrane. V. with a correcting glass $= \frac{5}{6}$. The projection was perfect. A good central opening in the capsule was obtained by needling

some weeks afterwards, and vision with a correcting glass was eventually $\frac{5}{6}$.

CASES VI AND VII.—In both these cases the eyes were in a state of acute inflammation at the time of operation. The injured eyes were excised a few days after the foreign bodies were extracted.

CASE VIII.—This case differs from the others in that the foreign body was removed from a position external to the globe after having passed through cornea, iris, vitreous, and sclerotic. V. at present time is $\frac{5}{6}$.

CASE IX.—Considerable injury was inflicted upon the eye in this case during the passage of the foreign body through the vitreous and lens into the anterior chamber; and, again, the extraction through the corneal wound caused considerable damage to iris and cornea. It is not to be wondered at, therefore, that the resulting inflammatory condition demanded excision.

A few general conclusions that we have arrived at as to the extraction of foreign bodies may be mentioned as follows:

1. When the patient is brought up to the magnet it is the rule for severe pain to be experienced by him if a magnetisable particle is present in the globe. Haab's magnet is therefore of valuable diagnostic service in determining the presence or absence of a magnetisable particle.

2. Localisation by the x rays is extremely important. By their means it is possible sometimes to avoid injury to a clear lens in the extraction of a foreign body, and entanglement in the iris can be avoided if its precise situation is known. Nevertheless, if some delay is unavoidable before the x rays can be applied, we advise in certain cases immediate extraction of the foreign body by Haab's magnet without precise localisation; for example (1) in which there is considerable inflammation; (2) in which there is a recent traumatic cataract. In each of these cases there is considerable danger in delay.

3. If there is a recent wound we attempt to remove the foreign body through it, using Haab's magnet alone. If this wound has firmly healed it is necessary to make an opening for its exit. This may be either corneal, in which case the foreign body is withdrawn by Haab's magnet; or peripheral, in which case we usually employ the small magnet for the removal of the foreign body from the anterior chamber after it has been brought forward by Haab's magnet. The latter method is the one we generally adopt.

4. In all cases it is most essential that entanglement of the particle in the iris and ciliary body should be avoided. By bringing the patient up to the magnet gradually from a distance, and by increasing the strength of the current up to the maximum slowly, this is to a certain extent guarded against. The idea of this manoeuvre is to prevent the particle from rushing forwards from its posterior position and burying itself in the iris.

THE RELATION OF GLAUCOMA TO THROMBOSIS OF RETINAL VEINS, WITH NOTES OF CASES.

By E. HARRIES JONES, M.B., C.M. EDIN., M.R.C.S. ENG.,
L.R.C.P. LOND.,

Honorary Ophthalmic Surgeon, Northampton General Infirmary; late
Resident Surgical Officer, Birmingham and Midland Eye Hospital.

THE relation of glaucoma to diseases of the retinal vessels has long been a subject of discussion to ophthalmologists. Several papers have appeared from time to time recording cases showing this relation, notably that of Wehrli,¹ which deals almost entirely with glaucoma following albuminuric retinitis. The following notes illustrate the fact that glaucoma may occur as a sequel to unilateral thrombosis of the central retinal vein, associated with arterio-sclerosis, but without albuminuric retinitis. Curiously enough, Michel, who was one of the first to describe the appearances due to thrombosis, stated "that the intraocular tension during the course of this affection is never increased." Glaucoma, however, is a well-known, though rare, complication of so-called hæmorrhagic retinitis.

CASE I.—G.W., aged 50, copper dresser, admitted as out-patient to Birmingham Eye Hospital, August 27th, 1900. Complaint: Gradual loss of sight of left eye since first week in August, 1900. No history of sudden onset. On examination fundus presented typical picture of thrombosis of