existing compression, and could have waited without much anxiety until further skiagraphs localising the bullet had been prepared, when I should have performed a second operation for its removal.

Operation.—The whole area of damage was exposed by turning down a large flap, beginning in front at the margin of the hair and ending posteriorly about 2 inches above and behind the mastoid. The fracture proved to be of the gutter variety—glancing impact; a number of fragments of the outer table were adherent to the flap, and a small quantity of blood clot and broken-down brain substance was found beneath the flap; on removing this, the outer table was seen to be splintered at the anterior part, while posteriorly the whole thickness of the bone had been ploughed up. The splintered fragments were removed and a large rent in the dura through which protruded brain substance was exposed, in which splinters were embedded. It was during the removal of the splinters that I touched metal, and on seizing this I withdrew the bullet; the projectile, which was embedded vertically, was split open from end to end, and a portion of the mantle was turned backwards and projected near the surface of the brain, and it was this projecting piece which I had touched. After careful exploration to ensure that all embedded fragments of bone had been removed, the wound was closely sutured and dressing applied. There was considerable shock during the operation, but this rapidly passed off.

After-History.—The convalescence was most satisfactory; at no time were there any symptoms to cause anxiety. The headache subsided within twenty-four hours, as also did the facial paralysis. On the fourth day the power of flexion at the elbow had returned; three days later the wrist could be extended, and on the eighth day flexion at the wrist was possible, then followed extension and flexion of the fingers, the movements of the thumb being the last to return. On the fourteenth day the stitches were removed and the wound was healed. The patient was kept in bed for three weeks; at the end of this period all but the finer movements of the hand could be performed but they required considerable effort. When I last saw the patient early in October the improvement was marked, the grip of the left hand was strong, and the execution of the more delicate and complex movements of the hand had to a great extent recovered.

EXCISION OF SPINA BIFIDA AND ENCEPHALOCELE.

By JOHN LITHGOW, M.D.,
Cleland, Lanarkshire.

WITHIN the last few years the old method of treatment of spina bifida and encephalocele by injection of Morton's fluid has been almost discarded for the newer, and certainly more surgical, treatment by excision. There have now been no doubt a considerable number excised, but the few cases published, and the meagre information given as to the details of the operation, suggest that it has only met with a limited degree of success. Even the newer textbook dismiss the discussion of the treatment of these cysts by excision with the statement that such has been attempted and the bare suggestion that it may be feasible.

The following cases occurring and treated in general practice may therefore be of some interest:

CASE I.

In the month of November, 1893, my attention was directed by my assistant to a tumour on the back of a child which he had delivered a few days previous. The child was a male, well developed, and except for the tumour seemed in perfect health. The tumour, situated in the lumbar region, was about the size of a small hen's egg; the skin over it was healthy though thin, and it was sparsely covered with hairs at the base, while the attachment to the spine was narrow. The tumour was diagnosed as spina bifida of the meningocele variety and as there appeared to be no urgency it was supported with collodion and wadding.

When two and a-half months old he had, as described by the mother, slight convulsive seizures which passed off after administration of calomel; there was never any trouble with the tumour, the skin covering it retaining its healthy conditions. I advised the parents to take him to the Glasgow Royal Infirmary in order to have something done to it in the way of operative treatment, and when three and a-half months old he was taken to that institution when the surgeon offered to inject Morton's fluid. Owing to the unfavourable prognosis given the parents refused to have this done, and the child was brought home with them. Shortly after this he had a sharp attack of whooping cough, and so great was the tension on the sac during the spasms that there was great danger of it giving way. The parents then gave their consent to me to try anything I choose in the way of operation, but it was not till the child was fully ten months old that I considered him free enough from the whooping-cough to attempt the removal of the tumour by excision.

Operation.—Dr. Witherspoon gave chloroform. After disinfecting the skin well over the tumour I marked out and dissected two flaps as follows: Commencing the incision from a point over the spine, about half-an-inch above the insertion of the tumour, it was carried over the side of the tumour near its base to a similar point over the spine, about half an inch below it. This was done on both sides, and the flaps dissected well clear of the base. By this means I was able to thoroughly lay bare all round the insertion of the tumour into the spine. Owing to the almost membranous state of the more prominent part of the cyst, great care had to be taken not to open into the cavity. When the base was thoroughly laid bare a pair of phimoses forceps, the jaws of which had been previously guarded, was passed with a blade on either side. The tumour was now pierced at its apex, and while it was slowly emptying itself the screw of the forceps was sent home, keeping the blades pressed

towards the spine as much as possible. Further escape of the cerebro-spinal fluid was thus prevented by the phimoses forceps, which now acted as an effectual clamp. Using catgut I stitched with continuous suture and small catches that part of the pedicle which was between the clamp and the spine; when satisfied it was watertight I cut with curved scissors close to the clamp all the membranes which lay on the top of it. The clamp was now removed, the stump washed, and the flaps brought over and stitched. In bringing the flaps together the stitches were made to include the stump lying underneath them, thus forming a much stronger buttress than would otherwise have been the case. During the operation no trace of any nerve substance was seen in the sac, which was composed of skin and membranes distended with fluid, which communicated with the spinal canal by an opening in the vertebrae, into which the tip of the finger easily fitted. There were no after-effects from the operation, and no after-treatment in connection with it, the child in a few days making a perfect recovery. The operation was done in a collier's house of one apartment, with no assistance save that of the chloroformist. A short time after the operation this family left the district, returning at the beginning of this year, when I found the boy had talipes equinovarus. My partner, Dr. Robertson, cut the tendo-Achillis, with a satisfactory result.

Since operating on the above case I have delivered three children (two of them cousins) afflicted with spina bifida. These three were almost identical; the opening was in the lower lumbar region, involving more than one vertebra; the covering of the tumour, which had a broad base, was like mucous membrane, and was eroded from birth. Owing to the soft nature of the covering, and the fact of the most of it being involved in the erosion, I thought they could stand nothing in the way of operation. All three died of convulsions within a few days of birth.

CASE II.

On August 27th, 1901, my *locum tenens* asked me to see a child he had delivered during that day. I found the child had an enormous tumour arising from the back of the head. The tumour was almost the size of the child's head, was slightly translucent at the apex, had a narrow attachment at the occipital region. This attachment was felt to contain between the skin something more than fluid. The whole tumour was covered with healthy skin. Diagnosis of encephalocele was made, and permission obtained to excise it.

Operation.—On September 8th, when the child was 12 days old, Dr. Robertson administered chloroform, and I proceeded to remove the growth. The details of the operation were the same as in excising the spina bifida. Flaps were dissected from the sides of the tumour; these flaps were well retracted all round, phimoses forceps clamping the attachment. I unfortunately removed the tumour before I stitched the pedicle; this made it much more difficult to do, as the brain substance on the pedicle collapsing between the jaws of the forceps, tended to throw them off; they could not therefore be utilised in steadying the part. After satisfying myself that there was no leakage, I stitched the flaps together, including the stump. The child was so much collapsed after the operation that we did not expect it to revive; however, on a rectal injection of brandy being given, the collapsed condition passed off. There was no trouble with the wound afterwards, healing took place by first intention, and all dressings were removed on the twelfth day, the scar and adjacent tissue being firm and normal. On the fourteenth day after the operation, when all danger was considered to be past, the child unfortunately had some convulsive seizures, and died.

In this case, owing to the difficulty in handling the child, and the fear of ulceration or accidental rupture, the operation had to be performed at an exceedingly tender age, and no hope was entertained that it would survive the initial stage of collapse. On the third and fourth days the mother reported that it had slight convulsions; after this, till the fatal seizure, there was no indication of brain disturbance. Owing to the rapidity with which the operation was performed, no time was spent in minutely examining the relations of the parts at the base of the skull; but an opening about the size of the point of the finger communicated with the interior. The tumour was lined throughout with brain substance. The whole was distended with fluid. This operation was also carried out in a collier's house of one apartment, and with no assistance save the chloroformist.

A POSSIBLE PREDISPOSING CAUSE OF CANCER.

By HAROLD MASON, M.B. LOND.,
Chairman of the Health Committee of the Borough of Leamington.

THAT cancer and malignant diseases generally have been for some time past on the increase is, I believe, an undisputed fact, the reason for it being one which has not yet been conclusively determined, and the investigation is naturally surrounded by a great many difficulties. The increase has become more marked during about the last sixty years.

I have collected from the death returns of the registrar upwards of 400 cases occurring in the Leamington district,

spreading over a period of fifteen years, and have analysed them in several ways. In the first table they are arranged according to the systems and organs which were primarily first attacked:

	Male.	Female.	Total.
Alimentary:			
Lips ...	—	—	—
Jaws and Face... ..	5	1	6
Tongue... ..	6	2	8
Esophagus	12	6	18
Stomach	28	33	61
Liver	18	36	54
Pancreas	5	3	8
Intestines	17	28	45
Rectum	19	18	37
Totals ...	110	127	237
Circulatory:			
Heart	1	1	2
Respiratory:			
Pleura	0	1	1
Lung	1	1	2
Larynx	2	1	3
Totals ...	3	3	6
Glandular:			
Mammary	0	57	57
Lymphatic	4	7	11
Spleen	1	0	1
Totals ...	5	64	69
Urinary:			
Kidney... ..	5	6	11
Bladder	7	3	10
Totals ...	12	9	21
Generative:			
Prostate	1	0	1
Testicle	1	0	1
Vulva	0	1	1
Ovary	0	5	5
Uterus	0	51	51
Totals ...	2	57	59
Organs not stated ...	8	17	25
Tegumentary, etc.:			
Leg	4	2	6
Pelvis	0	2	2
Eye	0	1	1
Totals ...	4	5	9
Grand Total, 427			

From this it is evident that the alimentary system accounts for 55½ per cent., the glandular for 16 per cent., the generative organs for 13½ per cent.; also that if those cases affecting only the mammary gland and the uterus are deducted, the liability of male and female is about the same.

The second table shows the ages in periods of ten years at which death has occurred:

	Male.	Female.	Total.
From birth to 10 years	1	1	2
From 10 years to 20 years	2	1	3
From 20 years to 30 years	4	5	9
From 30 years to 40 years	3	14	17
From 40 years to 50 years	15	44	59
From 50 years to 60 years	41	69	110
From 60 years to 70 years	41	80	121
From 70 years to 80 years	29	48	77
From 80 years and upwards	8	21	29
	144	283	427

This table shows: (1) That there is no decade in which cancer does not occur; (2) that the greatest liability occurs ten years earlier, and persists for ten years longer, in the female than in the male.

The next analysis refers to the residences of those dying

from cancer, which is too lengthy for this paper. The following facts, however, may be gathered from it:

1. That out of these 427 cases deaths from cancer occurring twice in the same house is noted but seven times, and in one of these the patient only moved into that house a few days prior to her death.

2. It is, however, very common for it to occur in houses in the same row and on the same side of the street. The more striking examples I have tabulated, using a letter and figure for a street index, giving the actual numbers in the street:

Street Index.	Numbers of Houses in which Cancer has occurred.
A ₁ ...	14. 16.
B ₅ ...	16. 18.
B ₁₃ ...	39. —. 43. also 17. 19.
B ₁₅ ...	40. —. 44.
C ₂ ...	2. 4. 6.
C ₇ ...	3. 5. —. —. 11. —. 23
C ₈ ...	50. —. 54.
C ₁₄ ...	12. 14. —. —. 22. 24. —. —. 30.
C ₁₆ ...	6. 8.
C ₁₈ ...	33. 35.
C ₁₉ ...	29. 30. —. 32.
D ₁ ...	2. —. 6. —. —. 42. —. 46.
D ₄ ...	1. 3. —. 7. —. 11.
G ₃ ...	5. —. 9.
G ₆ ...	48. —. 50.
H ₂ ...	18. —. —. 24. —. 28. 30.
L ₂ ...	14. 16.
N ₂ ...	21. —. —. 27. —. 31. —. —. 39.
O ₃ ...	14. 16.
P ₆ ...	4. 6. 8.
Q ₂ ...	36. —. —. 42.
R ₂ ...	31. —. —. —. 39.
R ₆ ...	74. —. 76.
R ₁₀ ...	38. 40.
R ₁₃ ...	55. —. 59.
T ₇ ...	5. 6.
U ₁ ...	3. —. 7.
U ₂ ...	3. —. 5.

The dashes represent houses in which deaths from cancer have not occurred during the last fifteen years.

These are only a few of the most striking, but sufficient for showing that the frequency with which two or more houses adjacent to each other are cancer houses, is probably not accidental or mere coincidence, but that there is some determining cause in common.

Out of the 427 cases 44 died in public institutions. Their previous residences I have not been able to obtain, and deducting 7 in which cancer has occurred twice, there remain 376 dwellings in which it has occurred, and of these I now give an analysis:

1. I find that 66, or 17.5 per cent., of these houses are the end houses of rows, or corner houses of streets.

2. That 178 are above the annual rental of £20, and 198 are below that value.

Now in this town there are approximately 2,263 houses of the former, and 3,793 of the latter value. This gives a percentage of 7.8 of the better class to 5.2 of the poorer.

There are also a good many streets in which deaths from cancer are conspicuous by their absence, as the next table will show:

A ₂ has 40 houses in which cancer has not occurred during the last 15 years.	
B ₂ " 12 " " " " " " " " " "	
B ₄ " 28 " " " " " " " " " "	
C ₆ " 35 " " " " " " " " " "	
C ₂₁ " 62* " " " " " " " " " "	
C ₂₂ " 40 " " " " " " " " " "	
D ₃ " 8 " " " " " " " " " "	
E ₁ " 47 " " " " " " " " " "	
E ₃ " 26 " " " " " " " " " "	
G ₂ " 40 " " " " " " " " " "	
H ₇ " 21 " " " " " " " " " "	
L ₉ " 21 " " " " " " " " " "	
N ₈ " 18 " " " " " " " " " "	
P ₃ " 20 " " " " " " " " " "	
S ₁₆ " 22 " " " " " " " " " "	
V ₆ " 26 " " " " " " " " " "	

* Including 2 courts.

With the exception of C₂₁ and L₉ all the houses are under £20 per annum.

The distribution of cancer is singularly patchy, some districts having very few, while in others it appears in large groups. There is one particular district with a population of 930 or thereabouts, in which cancer has occurred in four houses in these fifteen years. All these houses are well built, above the annual rental of £50, and situated on a stiff clay ground.

As regards the nature of the ground upon which the cancer-houses abound the following table will show that:

Gravel...	...	...	is the foundation or subsoil of about	125
Sandstone	...	...	"	57
Loose soil, made up, etc.,	"	"	"	3
Clay	...	...	"	86
Unascertained	...	...	"	105

As regards the drainage of the various houses in which cancer has terminated fatally, I regret I have been unable to obtain returns for the first half of the period (of these 427 cases), but I have collected drainage statistics for the last 175 cases, and I find that defective drainage has been reported at the office of the sanitary authority in 45, which gives a percentage of 25.7; a very large proportion, considering that it is the exception rather than the rule to call upon the services of the sanitary inspector, particularly in the larger and better class houses, as these occupants usually call in their own sanitary engineer, and so no record comes to the notice of the Health Department. Out of these 175 cases, deducting 1 in which a case has occurred twice, I find that 117 of these houses are upwards of thirty years old and 57 are less than thirty years old, that is 67 per cent. of these cases occurred in the older houses. All these observations seem to point to the conclusion that whatever the ultimate cause may be, cancer will be found to be due to some germ whose habitat is a sewage-contaminated subsoil, and that its entrance into the organism is through one of the mucous-membrane-lined apertures of the body. If this be admitted then all the following statements will be readily accounted for:

1. That the cancer death-rate has increased during the last sixty years—a time synchronous with the development and extension of the watercloset system of drainage, bearing in mind that the first style of the w.c. system was simply unglazed drain-pipes, laid end to end, sometimes joined with clay and sometimes without. I have even seen in old cancer houses drains made up of one brick laid flat, with two placed edgewise, and covered with slate running beneath the dwelling-house.

2. That the infection of the alimentary system is as high as 55 per cent. For in those houses in which the drains are in such a faulty condition as to contaminate the subsoil, the food, which is often kept in the cellars, would be the first to become infected by any germ that might be existing in the subsoil, considering there is always an upward current of air out of the subsoil into any building upon which it is built.

3. That people over 40 years of age are more subject to it, as they become more confined to their dwelling as age advances.

4. That females are more subject to it than males, possibly on account of their domestic duties keeping them longer exposed to any poisonous influences that may thus arise. (Even the shape of the female garments would favour infection of the generative organs.)

5. That consecutive houses in the same street are found often to be cancer houses. As many of these houses were built at the same time and have a common drain and if it has been badly laid or for any reason it becomes faulty the subsoil of several houses may become sewage contaminated. (A great many of the cancer houses enumerated in this paper have been found to have old disused cesspools, old contaminated wells, or decayed brick sewers lying forgotten in the soil beneath the house).

6. That a large percentage of these cancer houses (17 per cent.) are end houses of rows, corner houses of streets, or houses on either side of court entrances, as these are places where any defect in a drain would be most likely to be severely affected by sewage.

7. That the older houses are more subject to become cancerous than the modern ones, as the drains in the earlier days were not laid with the care they now are, that glazed pipes were not known and brick drains were often used and the drains were often run under the dwellings, which would after a time decay, become leaky, and thus contaminate the subsoil.

8. That the houses which are built on a porous subsoil, such as sandstone, gravel, etc., are more likely to become cancerous than those built on clay, though the latter in time will become saturated with sewage matter.

For a great amount of the statistical information I have to thank the registrar, the medical officer of health, and sanitary inspectors for their courtesy in giving me every facility in the above investigations.

TWO CASES IN WHICH THE ACTIVITY OF PHTHISIS WAS EXCITED BY OPERATIONS FOR LOCAL TUBERCULOSIS..

By A. J. RODOCANACHI, F.R.C.S.Eng.,
Honorary Surgeon, Ancoats Hospital, Manchester.

THAT phthisis may be rendered active, or general tuberculosis excited by an operation upon strictly localised tuberculous disease, such, for example, as fistula in ano, is a fact well known but not, perhaps, sufficiently insisted upon. The two cases here reported illustrate this danger in a striking fashion.

The first case was that of a man, aged 40, who presented himself with extensive disease of the metatarso-phalangeal joint. He had previously suffered from some strumous disease about the lower jaw but this was now healed. He also suffered from cough but there were no physical signs of phthisis in the chest. He was admitted an in-patient, and treatment was directed towards improving his general and local condition. He was then sent to a convalescent hospital, and on his return, being then in much better health but still unable to walk owing to the painful condition of the toe, an operation was undertaken on May 21st, and the diseased bone freely gouged away. Subsequently the toe became gangrenous and separated. The resulting wound granulated healthily. In spite of the local improvement the cough became much worse, signs of rapidly spreading phthisis appeared, and he died exhausted on July 15th, 1901.

The second case was that of a man, aged 22. He had a large mass of glands in the right groin and a smaller mass in the left. He also suffered from cough, but the sputum was examined for tubercle bacilli with negative results. There were, however, present physical signs of apical phthisis. An attempt was made in the first place to reduce the glands by potassium iodide liniment, together with rest and general measures of increasing nutrition. As, however, it became evident that suppuration had commenced in the glands, it was decided to excise them. This was done on September 13th, 1901. The smaller mass on the left side was easily dissected out, and the wound healed subsequently by first intention. On the right side operation was rendered difficult by the old adhesions present, and in the course of the dissection several large veins were wounded. Three days later the temperature rose, and subsequently remained at 102° to 103° F. A temporary improvement was brought about by the injection of antistreptococcus serum, and the wound appeared to granulate healthily. On the other hand, the disease in the chest made rapid progress, and vomiting also setting in, the patient died exhausted on October 18th. Shortly before death an examination of the sputum showed the presence of tubercle bacilli.

The question naturally arises whether the treatment adopted was the best for the patients, and whether the operations were justifiable. It was with some reluctance and at the special desire of the patients that I was induced to operate. Had they been in a station of life which would have permitted them to spend several months in idleness at a healthy seaside place no operation would have been undertaken; but as each patient was the breadwinner of a poor family I felt justified in attempting a cure by radical means, even at some risk, knowing that the local condition prevented him from attempting any work, and that a refusal would have condemned the patient to a lingering death in either a poor cottage or a city workhouse.

PROPOSED MEDICAL INFORMATION BUREAU IN PARIS.—The Council of the Paris Faculty of Medicine has under consideration a proposal to establish within the precincts of the Faculty an information bureau for the benefit of medical visitors, native and foreign, to the French capital. All necessary information as to lectures and courses of instruction, public and private, hours of hospital attendance, etc., will be supplied by an official on the premises, and the foreign visitor will even be able to obtain a list of any fellow countrymen who may be resident in Paris. The suggestion is due to Dr. Blondel, Secretary-General of the International Association of the Medical Press.