and that there was a long ulcer running across the lesser curvature, with a second perforation at its posterior end, which opened into the sac of the omentum.

CONCLUSION.

These 9 cases occurring before you within a short interval offer a good deal of food for reflection. The first related were comparatively simple, and under careful treatment ought to recover without operation. But it is impossible to say in any case of gastric ulcer whether some of the severe complications will not arise, and when they may be dangerous. Some of the most terrible, such as haemorrhage, come "like a bolt from the blue," without any warning; others, such as cicatricial contraction, are preceded by a long train of troublesome symptoms. All this must make us reflect whether surgical treatment ought not to be employed in a far larger proportion of cases where proper medical measures are not leading to a cure, than is at present the case. Operation, as you have seen, is only a desperate last resource, where perforation or haemorrhage is surely killing a patient, whereas, as you have also seen, recovery from operations for the other non-malignant conditions is the rule, if done before the patient is quite worn out.

So far, I am glad to say, all the operations I have done for non-malignant disease (without perforation) have been successful, and you have seen in some of the cases just related how completely they have relieved suffering and enabled the patients to blossom out into useful, active life again. The relief is perfectly marvellous often after years of suffering and inability to follow any avocation. I have recently had another in private, where a gentleman of 44 had been an invalid for a long time, and a great sufferer. A posterior gastro-jejunostomy has enabled him not only to return to his hard work after only a fortnight in bed, but also to enjoy social life, from which he had long been cut off.

I could have emphasised these points by a much larger number of cases operated on, but have preferred to confine myself to those which you have had an opportunity of studying for yourselves quite recently.

AN ADDRESS ON SOME POINTS IN CONNEXION WITH ULCERATION OF THE STOMACH AND DUODENUM.*

By C. R. BOX, M.D. LOND.,

Assistant Physician, St. Thomas's Hospital: Assistant Physician, London Fever Hospital.

My choice of ulceration of the stomach and duodenum as a subject on which to address you has been influenced by the great frequency with which such ulcerations occur; by the fact that the diagnosis of these ulcers is by no means so easy a matter as is commonly supposed; and by the serious complications which may occur in these affections, rendering their recognition and prompt treatment all-important.

It is impossible within the limits of a short paper to treat the subject of gastric and duodenal ulceration with exhaustive detail, so, with your permission, I propose to draw your attention to a few points which appear to be of practical importance.

First, with regard to the probable causation of these ulcers. In the current textbooks on medicine one finds a bewildering number of theories with but little actual evidence to support them. The ulceration is variously ascribed to arterial thrombosis, to embolism, and to vascular obstruction caused either by endarteritis or by simple spasm, all leading presumably to the formation of an infarct or of a haemorrhagic area, which is in turn digested by the gastric juice. Mechanical injuries are also invoked, such as a continued pressure due to occupation, as in shoemakers, or the constant wearing of tight belts or corsets, or actual blows upon the epigastrium. Or, again, a diminished alkalinity or the blood is supposed to lower the power of resistance of the gastric wall, and so lead to its digestion.

* Read at a meeting of the South London District of the Metropolitan Branch of the British Medical Association.

Of late years one finds advocates of the theory of microbic infection of the lymphoid follicles of the stomach wall. This is certainly the mode of production to which I incline myself, and for the following reasons:

1. For any part of the intestinal tube below the duodenum the theory of the infective nature of ulceration is at once accepted. Typhoid, tuberculous, and dysenteric ulcerations are unanimously attributed to an infective cause. The same holds for ulcerative colitis and the destructive process in appendicitis.

2. The fact that it is not uncommon to find that ulcers both in the stomach and in the duodenum may be multiple, and not only multiple, but so placed—either exactly opposite or quite close to each other—that infection from one to the other may have easily occurred. When two ulcers are placed opposite to each other, it is said that close examination will often show that one is an older ulcer than the other. I have repeatedly met with these opposed ulcers *post mortem*, not only in the stomach but also in the duodenum.

3. The appearance of a recent acute ulcer at the time of operation for perforation is itself highly suggestive, and it was this appearance which first seriously directed my own attention to the microbic theory. Such an ulcer is always surrounded by a zone of highly oedematous and softened tissue of considerable extent. Stitches easily tear through the oedematous wall, and so considerable difficulty may be experienced in applying the sutures. Every surgeon who has had much experience of the suture of acute perforated ulcers recognises this oedematous condition.

Surrounding oedema is not present in every perforated ulcer, for there is a form of perforation which is conveniently termed "secondary" in contradistinction to the "primary" perforation of an acute ulcer.

By "secondary" perforation, we understand a perforation in the floor of an ulcer, due either to the rent of a thin peritoneal base, or the tearing of some old adhesions uniting such an ulcer to the surrounding structures. The perforation of a stomach in this way by disturbance of its adhesions or by tear of its peritoneal floor may be brought about by overdistension of the stomach by food, by taking copious draughts of fluid which weigh down the organ, or by some unwonted muscular exertion when the stomach is so distended.

An ulcer which has perforated in this manner, and not by active spread of an inflammatory process, need not present an area of surrounding oedema at all.

We must then somewhat modify our conceptions of the acute ulcer. Surrounding the round or oval characteristically punched-out ulcer, containing, perhaps, a small necrotic slough, we must picture a large surrounding zone of oedema and softening, which can hardly be of other than inflammatory origin. Sometimes the recent ulcer so far departs from the textbook description that it actually has undermined edges like those of a recently-formed typhoid ulcer, an additional evidence of its formation by bacterial necrosis and sloughing.

What is true for the gastric ulcer, probably also holds for the common ulcer of the duodenum, nor do I believe it is necessary to assume that there is any essential difference in the origin of acute and chronic ulcers in these localities.

With regard to the embolic causation of gastric ulceration, the following points may be noted. It is difficult to produce ulceration this way experimentally; in those diseases in which embolisms of other viscera are most frequent, gastric ulceration is exceedingly rare; and when embolism of the stomach does take place, the embolus usually is produced by disease of one of the large arterial trunks in the neighbourhood of the stomach, such as, for instance, aneurysm of the coeliac axis. In the ordinary case of gastric ulcer one may search in vain for either the embolus or evidence of disease such as might give rise to an embolic process.

Superficial gastric ulceration is present in some cases of pyaemia, and gastric ulcer is sometimes met with in cases of mitral disease. In these cases the ulcers may possibly be of vascular origin.

The ordinary signs and symptoms of gastric and duodenal ulceration are too familiar to you to need any recapitulation on my part, so we may proceed at once to the consideration of two of the main signs, haematemesis and vomiting.

Haematemesis.

It is of often stated that gastric ulceration can only be positively diagnosed by the occurrence of haemorrhage in conjunction with other gastric symptoms. This is to a certain extent true, but there are many gastric ulcers which never make themselves known in this way, and in going over the accounts of cases of perforated ulcers of the stomach, it is very instructive to note in what a large percentage haematemesis never occurs. Indeed, beyond vague dyspeptic pains, the first obtrusive symptoms and signs are often those of the perforation itself.

Again in many cases of duodenal ulcer melaena is never noticed, and sometimes when it does occur, unassociated with haematemesis, the ulcer proves to be gastric after all.

My object, however, in referring to haematemesis is to draw your attention to the converse of the picture. There are cases in which haematemesis is associated with gastric pain and with vomiting, in which there is, nevertheless, no gastric ulcer at all. It is very important to recognise that such cases occur, particularly now that surgeons are claiming a larger share in the treatment of stomach complaints.

In the first place, haematemesis, together with epigastric tenderness and pain aggravated by the ingestion of food, may occur in certain cases of uraemia. The simulation of gastric ulcer by these cases is rendered the more complete by the facts that (a) such uraemic manifestations may occur in young women, and (b) that in some cases presenting such symptoms, albuminuria may be temporarily absent. I recorded two such cases in the *St. Thomas's Hospital Reports* five years ago, and since then have seen several others in which gastric ulcer was mimicked by renal disease, besides examining two similar cases, both in young women, *post mortem*.

A confident diagnosis of gastric ulcer, then, even in the presence of haematemesis, should never be made without careful, and if possible repeated, examinations of the urine, with the view to the detection of renal disease. This has been my own practice for some years past. The examination should not be limited to the detection of albuminuria, but a proper search should also be made for casts. It may be necessary to examine catheter specimens to avoid the fallacies dependent upon the presence of leucorrhoeal discharges.

In the second place—and what I mention now refers more particularly, but not exclusively, to male patients—it is necessary to bear in mind that the vomit which is brought up during gastric crises sometimes contains blood in traces, and occasionally in some quantity. This is important, for the possible occurrence of haematemesis in these cases is not even mentioned in many of the textbooks in ordinary use, although of course it is referred to by Gowers. I have recently seen two cases of locomotor ataxy in which during the crises haematemesis occurred; and here I may mention another in which, although there had never been haematemesis, yet the patient was so definite in his statement that his abdominal pain always came on two hours after a meal, that I was at first under the impression that he must have a duodenal or pyloric ulcer. The absent knee-jerks and pupils almost inactive to light revealed the true nature of the case in question.

Thirdly, there are those instances in which patients, usually young women with hysterical tendencies, simulate haematemesis. Cases of this sort are, unfortunately, not so very uncommon, and the simulation is often effected with such ingenuity, that, even though suspicion is aroused, it is very difficult to expose the fraud. The blood is often obtained by producing some lesion of a mucous surface, such as the mouth or vulva. Sometimes such patients also mix their urine with their vomit, and then the urinous odour, and the production of effervescence by the addition of sodium hypobromite as a test for urea, may expose the deception. In cases of this sort it is necessary to have the patient under close observation, and particularly to watch the actual act of vomiting, and to critically examine the vomited matters. This thorough observation can only be carried out in a hospital or similar institution, and even under these circumstances I have known a patient to come in for the second and even the third time before the fraud was discovered.

Fourthly, haematemesis occasionally occurs in connexion with mitral disease. Here, too, it may be associated with

epigastric pain and tenderness, and with vomiting after food. The routine examination of the patient should lead to the discovery of the mitral lesion, and possibly some evidences of venous back pressure, such as enlarged and tender liver, some oedema of legs, albuminuria, etc.

Vomiting.

It is important in cases where gastric ulcer is suspected to bear in mind that ulcer is not the only cause of constant vomiting in young people. The causes which appear to be most frequently overlooked are the vomiting of early pregnancy, the vomiting of uraemia, and the vomiting of cerebral tumour which is not manifesting itself by paralyses. It is necessary to bear the first in mind and to be always on one's guard. With regard to the second, the routine examination of the urine, upon which I have already laid stress, should prevent mistake; and in dealing with obscure cases of vomiting from cerebral tumour, in most cases an ophthalmoscopic examination materially helps in arriving at the diagnosis. It will be gathered from what I have already said that even when the vomiting regularly occurs a short time after meals, and is accompanied by pain which is relieved by rejection of the stomach contents, it is not necessarily characteristic of gastric ulceration.

It has been stated that in cases of gastric ulcer, after perforation of the stomach vomiting ceases. I do not know where this statement originated, but I do know, from repeated observations, that it is incorrect. It is not at all uncommon for patients with perforated gastric ulcers to continue to vomit. Nor must we infer, because the normal area of gastric percussion resonance is still present, that perforation has not occurred. The obliteration of the area of stomach resonance would at first sight appear to be a sign of some value in the diagnosis, but the resonance usually persists, and not only may gas be seen bubbling out of the perforation during operative manipulations, but the perforated stomach has been found distended with gas at the necropsy.

I would next direct your attention to the changes which occur in the physical signs at the base of the left lung in some cases of perforated ulcer on the anterior wall of the stomach. After the perforation of an anterior gastric ulcer, the peritoneum of the greater sac is not always at once involved in a general peritonitis. Occasionally the abdominal pain to a certain extent subsides, and the more acute symptoms for a time pass off. In such cases a careful examination of the base of the left lung, in the axilla and posteriorly, may shed considerable light upon the progress of events. Also, after operation for an ulcer of the anterior stomach wall identical signs may be observed, when a residual abscess is forming around the spleen. The abnormal signs discovered in this situation under these circumstances are:

1. Considerable restriction of the respiratory movements of the left side of the chest.
2. Diminution or cessation of the action of the diaphragm on the same side, with immobility of the upper left quadrant of the abdomen.
3. More or less localised pain and tenderness in the lower left axillary region, and sometimes actual oedema and bulging in this situation.
4. The presence of percussion dullness, which extends upwards in the axilla for some distance, and also posteriorly as high or higher than the angle of the scapula.
5. Diminution of the vocal fremitus, with more or less suppression of the breath sounds, or in some cases the appearance of distant bronchial breathing, over the area of diminished percussion resonance.
6. Friction sounds with a respiratory or even a cardiac rhythm over the lower part of the same area.
7. A certain amount of upward displacement of the impulse of the heart.

In addition to these signs there is another which is known as the "diaphragm phenomenon," by which is understood a shallow depression which travels across the intercostal spaces of the left side as the diaphragm ascends and descends. This last sign I have not investigated myself and so cannot speak positively about.

The proper interpretation of these signs is an important matter. They do not necessarily indicate that there is any

collection in the pleural sac, but they are primarily the signs of a localised abdominal collection around the anterior and external surfaces of the spleen, which in some cases also extends between the left lobe of the liver and the diaphragm. These collections are in the greater sac of the peritoneum, and are localised below by inflammatory adhesion between the belly wall and the great omentum or stomach. Indeed, as was pointed out by Greig Smith, a resistant or even indurated line may sometimes be felt through the abdominal wall in the situation of the lower limit of the abscess cavity.

The localisation of these collections of pus in the situation mentioned is considerably aided by the presence of a well developed "costo-colic fold." This fold, you will remember, extends from the diaphragm opposite the tenth and eleventh ribs, to the splenic flexure of the colon, forming a sort of shelf, above which lies the spleen. This fold materially assists in preventing the downward extension of these perisplenic inflammatory exudations, particularly if the patient is, as so often is the case, lying in the supine position. There is then a natural well in this position, containing the spleen and a part of the stomach, and ready for the reception of any fluid which may trickle into it.

Often the collection in this locality consists partly of fluid and partly of gas which has escaped from the rent in the stomach. Then we have the condition which has received the self-contradictory and cumbrous appellation of left subphrenic pyo-pneumothorax. The relative positions of the fluid and the gas will vary with the position of the patient. In the dorsal position the resonant area, which may yield a coin-sound and impart an amphoric quality to the sounds of the heart, will be found anterior, whilst the dull area is posterior. If the patient be turned upon the right side, the resonance makes its appearance at the lower part of the axilla.

In conjunction with Mr. Cuthbert Wallace and Dr. Russell I have paid considerable attention to the diagnosis and treatment of localised collections in this situation. We strongly deprecate the use of the exploring needle, for when this is used one is never certain of the exact locality of the collection. The proper method of exploration is by surgical incision, and resection of rib. By actual experiment we have found that resection of several inches of the sixth or seventh rib in the left axilla allows a thorough examination of the area involved. The pleural sac can thus be first examined, for it frequently in these cases becomes secondarily infected. If the pleural sac is still dry it can be accurately stitched off, before the diaphragm is incised to evacuate the subphrenic collection. Through the incision named a thorough examination of the external and anterior surfaces of the spleen can be made; the liver can be explored on its upper surface as far as the falciform ligament, and the left vault of the diaphragm also. It is also possible to drain by this route a collection of pus in the lesser sac of the peritoneum by tearing through or incising the gastro-splenic or the lienorenal ligament.

Another position in which we may often find a localised collection in connection with gastric ulcer which has perforated is in the pelvis. It is but natural that after perforation, some of the contents should gravitate to the lower part of the abdomen and the pelvis be thus infected. But it is not in every case apparently that the pelvis is directly involved in this manner. I have examined one case in which, apparently owing to a very badly-developed costo-colic fold, the perisplenic abscess has tracked down, within the peritoneum, along the course of the colon and been thus directed to the pelvis. A similar state of affairs is sometimes met with on the right side of the body, in connection with both gastric and duodenal perforation. In these cases the collection which has tracked down is sometimes situated between the lower surface of the liver and the head of the right kidney, this being one of the natural wells of the peritoneum and a favourite site for purulent collections.

There are, so to speak, certain natural highways in the belly for the spread of infection within the peritoneum. The ascending and descending portions of the colon, lying at the sides of the spine are two particularly important routes, and along these lines purulent infection may either ascend or descend. In conjunction with Dr. Russell, I drew attention to this important fact in a paper published in the *St. Thomas's Hospital Reports* for 1897.

It is therefore important in all cases of perforated ulcer, whether these have been operated on or not, to keep a careful watch upon the pelvis as well as upon the upper parts of the abdomen and the lung bases, and particularly to do this should the persistence of fever indicate that all is not going on well.

The localized collections I have mentioned are collections in the greater sac of the peritoneum. A general infection of the lesser sac is uncommon, as ulcers on the posterior wall of the stomach are usually, so to speak, "defended" by the formation of adhesions to the structures behind them and a partial obliteration of the lesser sac in this way. If this were not so the posterior collections should be more frequent than the anterior, as ulcer is more frequent on the posterior wall. As it is, however, perforation is much more frequent with the anterior ulcer. Recently I examined two cases in which, by secondary rupture due to tearing of old adhesions, the lesser sac had been infected in its whole extent. I do not, however, propose to discuss to-day the evidence of lesser sac infection, nor will I delay you with a consideration of the right subphrenic abscess in connection with disease of the stomach or duodenum.

In describing the localized collections I have mentioned, I do not wish it to be inferred that I advocate waiting in the ordinary case of perforation, in the hope that such a collection may form. On the contrary I believe that early operation is the proper method of treating all such cases.

In conclusion then, we should particularly bear in mind the following points in the investigation of supposed gastric ulcer:

1. The patient's statement as to the occurrence of haematemesis should be received with due reserve, and corroborative particulars inquired for.
2. In doubtful cases it is necessary to see the vomited materials oneself if this be possible, and better still to observe or have observed the actual act of vomiting.
3. In all cases the urine examination should be a routine procedure. This examination should be thorough, and include the search for casts as well as for albumen. A single examination is often not enough.
4. The eyes should be always examined, particularly with a view to the presence of the normal light reflex, and the occurrence of changes in the fundus. Optic atrophy may put us on the track of locomotor ataxy; the presence of optic neuritis may reveal an unsuspected cerebral tumour, and the presence of neuro-retinitis may indicate renal disease.
5. An examination of the stomach contents for the presence of excess of free HCl should always where practicable be made. And if the first examination be negative a second should be undertaken before being positive as to its absence. Hyperacidity is a valuable although not absolutely constant sign of gastric ulceration.
6. Every patient in whom gastric ulcer is diagnosed should be looked upon as seriously ill. The disease is commonly treated much too lightly, considering its liability to such serious and fatal complications. I believe that prolonged rest in bed and proper dieting for some time is essential in all early cases.

THE TREATMENT OF INTESTINAL OBSTRUCTION FROM MALIGNANT DISEASE.

By LEONARD A. BIDWELL, F.R.C.S.,
Surgeon to the West London Hospital; Dean of the Post-graduate College.

In this paper I shall only briefly refer to cases produced by a growth in the lower part of the sigmoid or the rectum, since I wish to particularly draw attention to the treatment of obstruction in the ascending and transverse colon. Before doing so I should like to mention a few points which may be of use in diagnosing at what part of the intestinal tract the obstruction is situated. First with regard to the history. When the obstruction is situated in the rectum, or lower part of the sigmoid, the early symptoms have usually consisted in the passage of small tapelike motions, and the patient has suffered from intercurrent attacks of diarrhoea and constipation, together with the presence of blood and mucus in the