SCIENTIFIC GRANTS COMMITTEE OF THE British Medical Association.

REPORT LXVII.

ON THE EFFECTS OF FORCED FEEDING IN CASES OF PULMONARY TUBERCULOSIS AND IN NORMAL INDIVIDUALS.

BY

NOEL D. BARDSWELL, M.D., M.R.C.P.LOND. & EDIN.,
Physician to the Sanatorium, Banchory;

FRANCIS W. GOODBODY, M.D., M.R.C.P.LOND.,
Assistant Professor of Pathological Chemistry, University College,
London;

AND

JOHN E. CHAPMAN, M.R.C.S., L.R.C.P.,
Clinical Pathologist to the Sanatorium, Banchory.
(From the Pathologico-Chemical Department, University College,
London).

THE following work on the metabolism in pulmonary tuberculosis was undertaken in the hope of throwing some light upon the question of how to feed patients suffering from this disease. For the opportunity of carrying out the research we are indebted to the courtesy of the Brompton Hospital Committee, and to Dr. Kingston Fowler, who placed six beds at our disposal for the purpose. The cases were selected as representing fairly typical phases of the disease. We also wish to express our cordial thanks to Professor Vaughan Harley, of University College, for much valuable advice and assistance in carrying out the work.

METHOD OF INVESTIGATION.

Before the commencement of the observations the food of each patient was weighed separately, in order to find out the composition of his ordinary diet; from this a diet was made out for one week. The urine and faeces were collected on the last four days in each period. The following estimations were made throughout the month:

In the Urine: Quantity, specific gravity, amount of total nitrogen, urea, uric acid, chlorides, phosphates, and sulphates (total and aromatic). *In the Faeces:* Quantity, water per cent., amount of nitrogen, and of fat. *Clinically* change of weight and condition as regards anorexia, dyspepsia, etc., were noted.

A.—CASES OF ACUTE PULMONARY TUBERCULOSIS.

Case 1.

H. A., aged 22, admitted July 18th, 1900.

Lesion.—Left lung: Extensive infiltration with commencing excavation in the upper lobe, and extensive infiltration of the lower lobe. Right lung: Infiltration of apices of upper and lower lobe.

Symptoms.—His temperature (rectal) ranged from 99° to 100.5° F.; appetite and digestion were good; night sweats were slight. His weight was 120 lbs., this being 22 lbs. at least below his highest known weight.

Diet.—During the first week under observation he was continued on the hospital diet, which consisted of milk 3 pints, bacon 1 oz., butter 1 oz., meat 3 ozs., bread 8 ozs., sugar 1 oz., vegetables 4 ozs., and milk pudding 5 ozs., roughly equivalent to 120 grams proteid, 120 grams fat, and 300 grams carbohydrate. On this diet he was found to be practically on nitrogenous equilibrium, only 0.32 grams of nitrogen being retained. His absorption of nitrogen and fat were somewhat low, namely, 91 per cent. and 94½ per cent. respectively, these low percentages being probably due to the use of enemata. During the following three periods the original diet was gradually increased, until in the fourth period the patient was taking double the original amount of proteid and fat and 25 per cent. more carbohydrate, the diets being made up of milk 5 pints, meat 4 ozs., eggs 2, bacon 2 ozs., butter 2½ ozs., bread 11 ozs., sugar 3 ozs., maltose 3 ozs., milk pudding 5 ozs., vegetables 6 ozs., beetroot 3 ozs., fruit 8 ozs.

TABLE I.

Period.	Increase of N. Ingested over Last Period.	Increase of N. Excreted in Urine and Faeces over last Period.	Increase of N. Retained over last Period
	Grams.	Grams.	Grams.
1	—	—	0.32
2	3.42	— 3.17	4.59
3	15.07	+ 6.84	9.23
4	6.21	5.27	0.94

Table I shows that on Diet II the patient's retention of nitrogen was very satisfactory. On Diet III the retention of nitrogen was certainly greater, but the extra excretion of nitrogen amounted to almost half of the extra amount ingested, while on Diet IV almost the whole of the extra amount ingested was excreted.

TABLE II.

Period.	N. per Cent. Excreted as Urea.	N. per Cent. Excreted in less Oxidized Forms.
1	85.82	10.74
2	91.74	5.95
3	79.70	17.33
4	79.41	18.12

Table II shows that the largest amount of nitrogen was excreted in the most oxidized form, namely urea, in the second period, the percentage amount excreted in less oxidized forms increasing greatly on the increased diets. Absorption of nitrogen was highest on the second diet, namely, 97 per cent., but fell to 90 per cent. on the fourth diet. Absorption of fat, on the other hand, was very good throughout. Clinically he did best on the second diet. On the large diets of the last two periods he gained very little more weight than on the second, and suffered much from anorexia and dyspepsia, symptoms which were quite absent on the earlier diets.

Case 2.

C. T., aged 36, admitted July 28th, 1900.

Lesion.—Right lung: Extensive disease, with excavation in the upper lobe, infiltration of the middle lobe, and extensive disease of the lower lobe. Left lung: Early disease in the upper lobe.

General condition very grave. High continuous fever, severe night sweats and exhaustion, appetite and digestion bad, sputum averaged 10 ounces daily. Was losing weight in hospital at the rate of 2 lbs. a week.

The diet in this case was very similar to that in Case 1; commencing with milk, 3 pints; meat, 3 ozs.; bacon, 1 oz.; bread, 8 ozs.; butter, 1 oz.; sugar, 1 oz.; vegetables, 4 ozs.; and pudding, 3 ozs., equivalent to proteid 120 grams, fat 120 grams, and carbohydrate 240 grams. The diet was gradually increased, until the patient eventually took double the original quantity of proteid, 70 per cent. more fat, and 20 per cent. more carbohydrate.

TABLE III.

Period.	Increase of N. Ingested over Last Period.	Increase of N. Excreted in Urine and Faeces over last Period.	Increase of N. Retained over last Period.
1	—	—	3.71
2	2.39	0.60	3.00
3	14.45	4.03	10.42
4	4.90	2.40	2.50

Table III shows that, as in the last case, the first increase of diet was followed by a lessened excretion of nitrogen and increased retention. On Diet III the amount of nitrogen

retained was increased, but 28 per cent. of the increased intake was excreted, while on Diet IV 50 per cent. of the increased nitrogen intake was excreted. The least waste then occurred on the second diet. Table IV shows that the

TABLE IV.

Period.	N. per Cent. Excreted as Urea.	N. per Cent. Excreted in less Oxidized Forms.
1	81.90	18.04
2	80.00	20.00
3	75.38	24.63
4	73.16	26.84

percentage of nitrogen excreted as urea was low all through, and, as in Case 1, the percentage diminished on increasing the diet. The absorption of nitrogen was low throughout, but highest on Diet II—namely, 94.4 per cent. The absorption of fat, however, was good on every diet. The excretion of aromatic sulphates increased greatly on the two larger diets, indicating increased intestinal putrefaction. Throughout the whole month under observation his weight was practically stationary, though it fluctuated considerably in each period. The larger diets of the last two weeks gave worse experimental results, and certainly gave rise to more dyspepsia and general discomfort than did those of the first two. The slightly increased diet of the second week was probably the best from every point of view.

B.—CASES OF CHRONIC PROGRESSIVE PULMONARY TUBERCULOSIS VERY MUCH BELOW WEIGHT.

Case 3.

J. T., aged 49, admitted July 16th, 1900.

Duration of illness, 10 months.

Lesion.—Left lung: fairly extensive disease in upper lobe, with some arrest, and infiltration of the lower lobe. Right lung: disease at apices of upper and lower lobes.

General condition rapidly improving. Appetite and digestion very fair. Weight 143 lbs.—still 53 lbs. below his normal weight; practically apyrexial.

Diet during the first period was very similar to that in the previous cases, and was roughly equivalent to proteid 110 grams, fat 120 grams, carbohydrate 245 grams. The diets during the second and third periods were increased greatly in proteid, and to a less extent in fat. The diet of the fourth period was reduced considerably in proteid only.

TABLE V.

Period.	Increase or Decrease of N. Ingested.	Increase of N. Excreted.	Increase of N. Retained.
1	—	—	10.67
2	+ 3.56	—0.46	+ 4.02
3	+15.53	+3.83	+11.70
4	—10.7	+0.11	—10.8

TABLE VI.

Period.	N. per Cent. excreted as Urea.	N. per Cent. Excreted in less Oxidized Forms.
1	79.24	20.76
2	81.03	18.97
3	82.45	17.55
4	79.02	20.98

On Diet I he was found to be retaining 10 grams of nitrogen and to be absorbing 87 per cent. and 93 per cent. of nitrogen and fat respectively, these low absorption percentages being

partly due to the use of enemata. On Diet II the increased nitrogen intake of 3.5 grams was associated with better nitrogen absorption, an increased nitrogen retention of 4 grams, and an increased nitrogen excretion of only 0.46 gram. The percentage of nitrogen excreted as urea also increased. The absorption of nitrogen was highest on this diet—namely, 95 per cent.; and that of fat was satisfactory—namely, 96 per cent. The gain in weight was the same as on the previous diet—namely, 3½ lbs. On Diet III, which was greatly increased in proteid, the patient continued to do very well. The greater part of the increased proteid was retained, and this retention was not associated with diminished nitrogen elaboration and increased intestinal putrefaction, as in Cases 1 and 2. The absorption of nitrogen and fat also showed no deterioration. Clinically the patient remained very well and gained 4½ lbs., so that we must conclude that in this case the large quantity of proteid given was of value. During the last week under observation, the diet as regards carbohydrate and fat was practically the same as in the previous week, but the amount of proteid given was considerably reduced. This reduction in proteid meant a reduction of 10.7 grams of nitrogen ingested compared with the last diet, and it is seen from Table V that the reduction in the amount of nitrogen retained was practically the same—namely, 10.8 grams—the absorption of nitrogen remaining the same. The weight increased 1 lb. This fact supports the opinion expressed as to the value of the large proteid diet in period 3, and we consider that in this case the largest diet given was the best. It is of interest to note that the increase of proteid was given in the form of somatose, and so caused but little increase in the bulk of food given.

Case 4.

E. R., aged 34, admitted July 22nd, 1900. This patient's tuberculous history was of at least five years' duration, and his symptoms and physical signs were characteristic of chronicity; for example, dyspnoea, clubbing of fingers, marked dyspepsia, etc. Unfortunately he failed to take the prescribed diets, and was very untrustworthy, so that the results are not of much value. One point, however, was of interest, namely, that in spite of very pronounced symptoms of disordered alimentary tract his absorption of fat was normal, 97 per cent. His absorption of proteid, on the other hand, was poor, 93 per cent.

C.—CASES OF ARRESTED PULMONARY TUBERCULOSIS: APYREXIAL AND UP TO WEIGHT.

Cases 5 and 6.

Both these cases had been admitted some six months previous to the commencement of observations, and had during that time been treated in the special sanatorium wards of the hospital. The results of this treatment had in both cases been very satisfactory, the disease in each patient being arrested. Both patients were also quite restored in general health, free from fever, and up to their highest known weights.

During the first week under observation they were continued on the ward diet, which they had been taking since admission. This diet consisted of milk, 3 pints; meat, 7 ozs.; bread, 7 ozs.; 1 egg; bacon, 2 ozs.; butter, 3 ozs.; sugar, 1 oz.; milk pudding, 3 ozs.; vegetables, 6 ozs.—roughly equivalent to 150 grams proteid, 210 grams fat, and 230 grams carbohydrate. The diet was very similar during the second week. The following table gives the metabolic results during the first two weeks:

CASE V.			CASE VI.	
	First Week.	Second Week.	First Week.	Second Week.
Proteid	148 grams	145 grams	144 grams	143 grams
Fats	212 "	192 "	212 "	183 "
Carbohydrates...	222 "	217 "	235 "	212 "
Total calories ...	3,494	3,274	3,526	3,265
Total N. in urine	13.2 grams	14.6 grams	15.6 grams	17.6 grams
Urea	24.1	27.0	28.3	30.9
N. absorption...	95.8 %	96.2 %	95.0 %	95.7 %
Fat absorption...	98.5 %	98.9 %	97.3 %	98.2 %
Weight in kilos..	34	34.7	65.3	66.2

It is seen from the above that the diets gave very good metabolic results. The retention of nitrogen was very satisfactory, and the percentage of nitrogen excreted as urea was high, showing that it was used to good advantage. The absorptions of nitrogen and fat were normal, the amounts excreted in the faeces being small. The small amounts of aromatic sulphates excreted in comparison to the alkaline showed that the intestinal tract was in good condition as regards putrefaction. Clinically the patients also did very well, both gaining weight. During the last two weeks under observation it was found impossible to persuade these patients to alter their diets sufficiently to allow of any comparison of results being made.

SUMMARY.

The observations on these cases show the following points of interest:

1. That pushing the amount of proteid in the diets in Cases I and II resulted in:

(a) An increased excretion of nitrogen out of all proportion to the increased amount retained.

(b) A diminution in the percentage of nitrogen excreted as urea, and consequently an increase in the percentage amount excreted in less oxidised forms, indicating diminished nitrogen elaboration.

(c) A diminution in the percentage of nitrogen absorbed.

(d) An increase in the amount of aromatic sulphates excreted, indicating increased intestinal putrefaction.

2. The absorption of fat remained very good in all six cases, even when very large amounts were taken.

3. The amount of nitrogen excreted in the urine on comparable diets was very small in cases much below weight, and was largest in patients approaching or up to their normal weights.

4. Clinically it was observed that the very large diets gave rise to much anorexia, dyspepsia, and in one case to vomiting. It was also noticeable that the onset of dyspeptic symptoms nearly always coincided with experimental figures indicating deranged metabolism.

5. In every case weight was gained, the gain being greatest on the very large diets, but with these large diets the gain was achieved at the expense of the physical well-being of the patient, as was shown both by clinical and experimental results.

6. It was observed in regard to the diets that the original generous hospital diets of Cases I, II, III, and IV were capable of being distinctly increased with advantage to the patients. Still further increase of the diet as already stated was followed by less satisfactory results, for example, in Cases I and II. The original diet of Cases V and VI, which was that of the special sanatorium wards, was found to be satisfactory in every respect, and was not improved upon during the month's observations.

RESULTS OF FORCED FEEDING IN NORMAL INDIVIDUALS.

The effect of forced feeding upon three normal individuals up to weight and in nitrogenous equilibrium were:

1. A marked increase in the amount of nitrogen excreted in the urine, with a proportionate increase in the amount of urea.

2. No diminution in the absorption of nitrogen.

3. A diminution in the absorption of fat.

4. A rapid and large gain in weight, which was in every case associated with marked impairment of general health. The chief symptoms induced were anorexia, nausea, dyspepsia, drowsiness, abdominal discomfort, and diarrhoea.

5. The weight put on was rapidly lost on resumption of ordinary feeding.

CONCLUSIONS.

In conclusion, we venture to make the following suggestions:

I. That since very large diets gave worse results than those of more moderate amount, the indiscriminate stuffing of all tuberculous patients should be replaced by systematic dieting. The diets as regards amounts and constitution should be determined in each case after due consideration has been given to the respective conditions as regards:

(a) The activity and extent of disease.

(b) Amount below weight.

(c) Digestive capability; and

(d) To some extent, personal dietetic likes and dislikes.

II. That in view of the bad effects which over-feeding gave rise to in the normal individuals, great care should be taken in the selection of a diet for patients who, as the result of

treatment, have reached or passed their highest known weights. When this regain of weight is associated with arrested disease, the original diet found suitable for a person very considerably under weight and with active lesions, should be reconstructed more upon the lines of what would be suitable for the same person in perfect health.

The observations were not sufficiently extended to allow of conclusions being drawn as to which diets gave the best results as regards the condition of the lungs; but subsequent sanatorium experience of two of us, with the advantage of being able to carefully observe the course of pulmonary lesions in patients upon weighed diets, leads us to think that the lungs do not improve any more rapidly upon forced feeding than upon generous diets. Further there is no doubt that anorexia, dyspeptic symptoms, and vomiting are much more frequently met with when working with very large diets than when more moderate amounts of food are given.

The following diet summary may be of interest and of practical value:

Diet Table.

	Diet of Cases I and II, which Proved to be too Small.	Diet of Cases I and II, which Proved to be too Large.	Diet of Brompton Hospital Sanatorium Wards, which gave Good Results.
Milk	3 pints	5 pints	3 pints
Meat	3 ozs.	7 ozs.	6 to 7 ozs.
Somatose ...	—	3 ozs.	—
Egg	—	2	1
Bacon	1 oz.	2 ozs.	1½ to 2 ozs.
Bread	11 ozs.	11 ozs.	7 ozs.
Butter	1 oz.	2½ ozs.	3 ozs.
Vegetables ...	4 ozs.	6 ozs.	2 ozs.
Fruit	—	8 ozs.	—
Milk pudding...	5 ozs.	5 ozs.	4 ozs.
Sugar	1 oz.	4 ozs.	1 oz.

Metabolism Tables.

CASE I.

	Period 1.	Period 2.	Period 3.	Period 4.
Proteid	116 grams	137 grams	232 grams	271 grams
Fat	123 "	137 "	184 "	231 "
Carbohydrate ...	296 "	296 "	321 "	392 "
Total calories ...	2,897	3,105	4,126	5,026
N. in urine	16.6 grams	14.5 grams	20.8 grams	24.0 grams
Urea	30.7 "	28.3 "	35.5 "	40.0 "
N. absorbed... ..	91.2 %	97.5 %	94.2 %	89.0 %
Fat absorbed	93.8 %	98.1 %	98.0 %	96.0 %
Weight in kilos. ...	50.7	52.3	54.3	56.0 %

CASE II.

	Period 1.	Period 2.	Period 3.	Period 4.
Proteid	115 grams	130 grams	220 grams	251 grams
Fat	121 "	127 "	180 "	208 "
Carbohydrate ...	240 "	225 "	265 "	297 "
Total calories ...	2,590	2,654	3,661	4,187
N. in urine	13.4 grams	13.0 grams	15.0 grams	18.0 grams
Urea	20.5 "	22.0 "	24.5 "	28.5 "
N. absorbed... ..	93.3 %	94.4 %	91.6 %	94.0 %
Fat absorbed	95.0 %	98.3 %	95.7 %	97.0 %
Weight in kilos ...	53.4	53.0	53.8	53.6

CASE III.

	Period 1.	Period 2.	Period 3.	Period 4.
Proteid	109 grams	131 grams	228 grams	161 grams
Fat	123 "	190 "	197 "	186 "
Carbohydrate ...	245 "	249 "	283 "	267 "
Total calories ...	2,594	3,332	3,936	3,495
N. in urine	6.7 grams	6.8 grams	10.2 grams	11.1 grams
Urea	11.5 "	11.9 "	18.0 "	18.8 "
N. absorbed... ..	87.0 %	95.0 %	94.4 %	94.5 %
Fat absorbed	93.0 %	96.4 %	98.5 %	97.8 %
Weight in kilos ...	62.5	64.0	65.5	66.0