chemical and physiological standpoints, show the valuable nutritive properties of the substance under investigation.

NOTES AND REFERENCES.

¹ London: R. H. Porter, 1900. ² The number so obtained was used for calculating the amount of proteids on the basis of Chittenden's (*Zeitschr. f. Biol.*, vol. xxv, p. 358) analyses of myosin, which showed it to contain 16.86 per cent. of nitrogen. ³ Probably what is taken as glycogen includes also intermediate substances between glycogen and dextrose. ⁴ Dormeyer (*Plüger's Archiv*, vol. lxi, p. 341, lxxv, p. 90) has pointed out the importance of this precaution. If gastric digestion is omitted, all the fat is not extracted.

FOUR CASES OF RODENT ULCER TREATED BY X RAYS.

By JOHN WILLIAMSON PUGH, M.D.LOND.,
Brighton.

THE following cases of rodent ulcer successfully treated by x rays are placed on record, as the number of such cases hitherto published is very small. The treatment is so successful

Fig. 1.—A.

that if applied at an early stage the ravages of this disease ought to be almost completely prevented. The danger of dermatitis is almost absent if the treatment is carried out under proper supervision, including an efficient arrangement of masks. There can be little doubt that the exaggerated fears of dermatitis which prevail have led to this efficacious form of treatment not being utilized to anything like the extent which it deserves, and which is desirable in the best interests of the patients. As a matter of practical detail I do not advise a sitting beyond fifteen minutes for cases of rodent ulcer. I am, however, at present giving longer sittings for a patient suffering from carcinoma of the breast, for which operation has been refused, special precautions being taken in this case against dermatitis. There can be no doubt, in my opinion, that the x rays have a marked destructive effect upon lowly organised and rapidly-growing cells, as seen in the following cases of rodent ulcer and in recorded cases of ulcerating carcinoma. I have applied the same treatment to boils, and in several instances have noticed that the course of the suppuration was much curtailed.

CASE I.—Mr. S., aged 93, gardener, for his time of life enjoys very good health.

History.—Three years ago a small pimple was noticed on his left cheek.

A year later he struck the pimple when chopping wood, and from that date it began to grow and ulcerate, and also to pain him at times.

State on Examination.—At the beginning of December, 1900, the ulcer was nearly circular, about an inch in diameter, the edges considerably raised (one-sixteenth to one-tenth of an inch) and depressed in the centre (crateriform). The discharge was blood stained.

Treatment.—I commenced treating him with the x rays on December 11th, 1900, using a coil which gave a 10-in. spark, and was worked by a 12-volt battery. At first the rays were applied for 5 minutes twice a week, keeping the ulcer 6 in. from the focus tube. The face was covered by a mask of lead foil with a hole opposite the ulcer. The time of exposure was gradually increased to 10 minutes, and the tube was brought within 4 in. of the ulcer.

Progress.—Up to the middle of January, 1901, there was not much encouragement; at that time the ulcer was a little smaller and the edge not quite so raised.

On February 11th, and up to the 18th, I gave him a daily sitting of ten minutes, and by that time the mask had become too small through the edges tearing away, and thus the greater part of the right side of the face was exposed to the rays. The result was that the patient had a sharp attack of dermatitis over the exposed part, but the effect on the ulcer was most marked. The discharge and pain almost ceased, and the edge was considerably reduced. Lotio plumbi c. op. was applied and the dermatitis rapidly disappeared. From February 18th to March 11th the rays were applied every other day, the edge gradually disappeared, the ulcer acquired a healthy appearance, and healed up from the sides

Fig. 1.—B.

The process was complete by March 18th, 1901, and the treatment was then discontinued. The scar at the present time (March 25th, 1902), is perfectly sound, and looks better than it did at first. (See Fig. 1.—A and B.)

The rays were applied about thirty-six times altogether. The hairs were destroyed over the exposed part where he had dermatitis. The same thing happened to my own finger while attempting to intensify the effect of the rays by holding a convex lens between the focus tube and the ulcer. This was a useless proceeding, as the rays, I understand, are very slightly, if at all, refracted by an ordinary convex lens.

CASE II.—Mr. T., aged 83, was treated for a rodent ulcer upon the left temple, 1½ in. by 1 in.

History.—It commenced about thirty years ago, and for many years it had been too tender to allow him to lie upon that side. During the last two years he had suffered very much from severe darting pains. The discharge was very abundant and bloodstained.

Treatment and Progress.—The treatment was commenced on March 18th, 1901. After a few sittings the first improvement noticeable was the disappearance of the pain and the lessening of the quantity of the discharge, which became of much lighter colour. Owing to the infirmities of the patient the rays could not be applied from April 27th to May 11th. Then a slight improvement was noticeable in the ulcer, the edge on one side was less raised, and its circumference was diminished. The tube had hitherto been placed about 4 in. from the ulcer, and the rays applied for seven and a-half minutes. As the ulcer was healing from the sides the distance of the tube was increased to 6 in., and the hole in the mask

lessened so as to protect the new skin. Whenever there was any sign of dermatitis around the edges lead and opium lotion was applied. By the middle of July the ulcer was about $\frac{3}{4}$ in. in diameter. There was no pain, and he was able to lie on his left side. After the above date no rays were applied. (See Fig. II.—A and B.)

On returning from my holidays a month later I went to arrange for further application, but to my great surprise the ulcer had healed up, thus showing that the diseased tissue had been destroyed, and also showing that there is no need to apply the rays until all vestige of such an ulcer has disappeared. The number of sittings was thirty-four.

Fig. II.—A.

CASE III.—Miss P., aged 48.

History and State on Examination.—A little over two years ago a small spot appeared on her forehead. By the time she came to me (April 22nd, 1901) it had gradually increased to $\frac{3}{4}$ in. $\times$ $\frac{3}{4}$ in. and projected considerably beyond the surface. The proliferations were large and the veins showed very much on the surface. There was no ulceration or discharge, but she had darting pains radiating from the growth.

Treatment and Progress.—The rays were applied to the growth for the first time on April 22nd, 1901, the rest of the face was protected by tea lead. A ten minutes sitting was given every day. After the tenth sitting its appearance gradually changed, the veins disappeared, and a scab formed on the surface. The scab came off showing a raw surface with some bleeding, but each time the place was smaller. The sitting was then

reduced in duration to 5 min. and the distance of the tube increased to 6 in. The last application (the 28th) was given on May 24th, and a fortnight later the wound had entirely healed, and on December 14th, 1901, there remained a slight flat scar as if from an old burn.

The growth in this case had a year previously been regarded as a keloid. There was no ulceration until the use of the x rays, but the growth presented the appearance of soft reddish nodules, projecting from the skin. It was probably rodent ulcer prior to the stage of ulceration.

CASE IV.—Mrs. M., aged 70.

History and State on Examination.—This was a very old case of rodent

Fig. II.—B.

ulcer of the nose. Owing to the ravages of the disease, and the use of knife and caustics, a considerable amount of the nose had disappeared. The rodent ulcer recurred in the scar on the nose and a spot of an apple jelly appearance showed on the cheek about $\frac{1}{4}$ in. beyond the scar.

Treatment and Progress.—After a few applications the ulcer on the nose soon began to heal. The spot on the cheek at first broke down into an open ulcer, and then healed up in the usual way. The number of sittings altogether were twelve. The parts were quite healthy, nearly three months after the last application of the rays.

In all these cases I have never used full power of the coil, only sufficient to produce fair penetration as shown by the screen, and as the tube became weaker the sparkage was increased.

FIVE CASES OF MONILIFORM HAIR ASSOCIATED WITH ALOPECIA AREATA.

By DAVID WALSH, M.D.,
Honorary Physician, Western Skin Hospital, London, W.

THE interesting congenital condition of beaded hair known as monilethrix is one of great rarity. It may be of interest, therefore, to publish a note of some cases in which hairs were found of moniliform type, although differing from the sharply-defined condition of monilethrix. The latter is congenital, and runs in families, while those affected have the hair of the head short and scanty, so that they are more or less bald. Lastly, the eyebrows and any part of the hairy body may be affected. That description differs widely from the following cases, in which the hair, although alopecia patches were present, was not short and scanty. The peculiarity was found only in hairs that had turned or were turning white. It did not involve hairs on other parts of the body. Lastly, the nodes were of irregular length, except in a few hairs of

the first case, which showed a fairly uniform beading, and the pigment had no distinct relation to the internodes—unlike monilethrix, in which the pigment is found gathered in the internodes.

CASE I.—A.B., female, aged 42, came to the Western Skin Hospital complaining of hair falling out in patches. There were several rounded spots on the crown and occiput covered with fine downy hairs, evidently a lanugo growth on areata spots. The surface layer of hairs of the head were white, while the hairs beneath retained their original dark colour. Attention was drawn to the refraction of some of the grey hairs, some of which were examined under the microscope, and found to be of moniliform shape with irregularly distributed pigment. The coarse white hairs were all more or less moniliform, but the change was not found in the eyelashes or eyebrows, nor, after repeated examination, in any of the original dark hairs. The pigment had no uniform distribution in relation to the nodes. In that way it differed from monilethrix, in which the pigment is collected at the internodes, and is general on the hairs of the head and other parts of the body. A hair shown at the Dermatological Society of Great Britain and Ireland is figured in illustration (No. 1.) In some of the hairs the pigment was carried right through the nodes and internodes, while in others it was broken up irregularly.

CASE II.—The patient volunteered the statement that her sister had bald patches and similar hairs. Some of the hairs sent for examination showed irregular refraction, as in the foregoing case. In a few of the grey hairs there were some long, slender, moniliform narrowings.