asthma. She has one brother and he is quite healthy. There is no consumption or other disease in the family.

Personal History.—She has always enjoyed very good health, and has never been laid up by a serious illness.

History of Present Condition.—The disease began on the left cheek when she was 16 years of age. She remembers it first as a small red patch which gradually increased in size. Ten years ago it was scraped, and two years ago it was again scraped and cauterized. In spite of the operations the disease has never been completely eradicated, and since the last scraping it has been a good deal worse.

Present Condition.—There is a triangular-shaped patch of lupus on the left cheek, the base of the triangle being uppermost, and represented by a line drawn from the lobe of the ear to the lowest point of the ala of the nose. From this base line the patch gradually tapers almost to a point, terminating at the lower edge of the jaw. Transversely the diseased area measures about $1\frac{1}{2}$ in. and the same vertically. The patch presents a dark-brown, scaly, slightly-projecting surface. It is infiltrated with the disease, which is confluent except at the upper edge, where there are some discrete nodules. There is no visible scarring, but the characters of the disease are partly obscured by large scales. There are no satellite nodules and no disease elsewhere. March 22nd, 1901. Began light treatment, one hour daily. March 26th. Severe reactions have occurred and the part is sore and covered with large blisters. April 23rd. There is less thickening of the affected skin, and it is paler in colour. May 6th. The whole area is inflamed and discharging so that the granulated tissue is hidden. June 15th. She is making a steady improvement. Most of the infiltration has disappeared. July 31st. Has had a fortnight's rest. The patch is rather red, and very little "apple-jelly" can be seen. August 13th. Treatment postponed for summer vacation. September 10th. Still further improvement, but one or two small superficial nodules remain. She has had eighty-eight applications altogether. October 15th. She is now only coming for treatment occasionally having three or four consecutive applications and then an interval of a fortnight or so. December 21st. One or two nodules still crop up occasionally after treatment has been stopped for any considerable period. They appear in the scar, which takes the form of a vertical ridge with lines radiating from it. This scar, which is rather dense, was not apparent at the commencement of treatment, being hidden by the lupoid infiltration. January 14th, 1902. There is now a smooth shiny scar with a small doubtful nodule in one part of it. This was treated to-day. There is also a small scaly spot, brown in colour but leaving no stain on deep pressure in the healthy skin just below the main patch. As this has been present for some time it was thought advisable to give it an application of the light.

Rodent Ulcer.

CASE I.—Man, aged 45. In this case the small rodent situated on the right cheek, which disappeared after seven applications of the light, the last being on July 6th, 1900, relapsed after an interval of about a year. He returned on September 11th, 1901. At the upper and inner margin of the scar there was a small raised growth the size of a pea, the top was flat and slightly depressed in the centre, while the edge was rounded and hard. This was the site of the original trouble, and the only part that had any suspicion of induration when the light treatment was stopped. The remainder of the scar was quite healthy. He had noticed the occurrence for some months, and it was gradually getting larger; it was very slight at first, and he cannot remember when he first noticed it. It was decided to treat the lesion with x rays, and he began with this treatment on September 11th. The face was covered in the usual way with a lead mask, a small hole having been cut out corresponding to the position of the growth. The primary current was one of 4 or $4\frac{1}{2}$ ampères. The tube gave a spark of 8 in. and was placed at a distance of 4 in. (= 6 in. from anticathode) from the patient. Daily exposures of twelve minutes were given. After the third application (September 13th) there was some diminution in the irritation and the growth seemed to be slightly softer. After the fourth (September 16th) the lesion was flatter and less indurated, but there was no redness of the skin. After the fifth a small crust formed on the summit of the wart-like growth. The flattening and softening were more noticeable. After the sixth there was practically no induration, the scab came off easily leaving a glazed slightly red surface. After the seventh (September 20th) another small crust had formed, and remained until treatment was discontinued on September 23rd after he had had ten applications. On September 27th the crust was easily detached and revealed a smooth red surface, with no elevation or depression of the epidermis and no underlying or surrounding induration.

CASE II.—Mrs. B., aged 36. This patient was under treatment from June 1st to June 16th, 1900, for a small rodent of the right cheek in the malar region. When she had had nine applications of the light the infiltration had broken down to form a circular ulcer which healed leaving a healthy scar. On December 10th, 1901, she returned for treatment. There were two points of recurrence, one was a raised, warty-looking growth, the size of a pea, at the upper and outer edge of the scar, and a linear, slightly raised, shiny, translucent infiltration corresponding with the inner border of the scar, the rest of which was to all appearances sound. She says the spots formed very gradually, but she cannot give the exact time of recurrence. Light treatment was resumed on January 13th. The area treated broke down as before to form a punched-out ulcer. She is still under treatment.

Lupus Erythematosus.

Married woman, aged 44. This patient, who had very extensive lupus erythematosus affecting the upper and lower lips, chin, and skin of the infraorbital regions, derived great benefit from the treatment, the diseased portions being replaced by a very fine white scar, resembling the normal skin. Unfortunately, however, she was only able to have treatment intermittently, so that the disease was never entirely eradicated; and in some parts—namely, on the skin of the lips and chin—the disease began to spread again soon after the cessation of treatment. She stopped having treatment after having had about one hundred applications of light, the condition having very much improved.

REFERENCES.

¹ Loc. cit. ² Loc. cit.

THE THERAPEUTIC EMPLOYMENT OF X RAYS.

By G. H. LANCASHIRE, M.R.C.S.Eng., L.R.C.P.Lond.,
M.D.Brux.,

Assistant Physician to the Manchester and Salford Hospital for Skin Diseases.

THE physical treatment of certain skin diseases being a subject which is now arousing much interest, I purpose shortly summarizing here the results I have obtained during the past twelve months with the x-ray tube.

There are many physical phenomena associated with a Crookes's tube in action, but the deeply interesting question as to which of these phenomena is the curative factor will not be discussed in this paper. Be this as it may, the evidence collected up to the present time, both from this country and abroad, may be said to have fairly established the x-ray apparatus as a curative instrument of the first importance in the treatment of certain diseases. The list of these diseases has been as yet by no means determined, but would appear to be extending.

The successful employment of the x-ray apparatus for curative purposes calls for considerable experience in its management. The treatment is not to be indiscriminately resorted to for all cases, or in all stages of disease; moreover, as in most other "treatments," the method of application is half the battle. Passing by the question of the supposed directly bactericidal effects of the x-ray tube—a now fairly disproved hypothesis—can any broad theory be advanced which shall explain the lines on which the cure of disease by this method proceeds? It would appear reasonable to conclude, in the present state of our knowledge, that the good effects of the "raying" are due to the mechanical stimulation of normal cell activity, whereby morbid tissue is either thrown off or absorbed by the healthy. In other words the process partakes of the nature of an inflammation. It is the control of this inflammatory reaction, as far as possible, which calls for the nicest judgement.

Hitherto one of the chief bugbears of x-ray work has been what is known as x-ray dermatitis, meaning by this an intense and long-persisting aggravation of the normal reaction. The avoidance of this calamity in itself demands no little foresight and experience, and even despite the greatest precautions the trouble will occasionally occur. The difficulty is sometimes increased by the long incubation period, so to speak, of the dermatitis, which may develop its full intensity even so late as a fortnight after the last exposure.

In certain individuals there would appear to be some idiosyncrasy which renders them prone to excessive reactions of this type; on the other hand, with others it is remarkable how long they may be exposed to the tube without appreciable reaction. The occurrence of x-ray dermatitis may probably be best avoided by keeping the tube at a fair distance (8 or 9 in.) from the skin, and by ensuring short exposures (ten to fifteen minutes) and periodic rests from treatment. The moment the reaction appears to be too severe (which may be suspected if the redness of the unbroken skin becomes too vivid, or if increasing discomfort be complained of) the treatment should be at once suspended. I have found simple dressings of linimentum calcei relieve the pain in this condition as well as anything.

After the dermatitis has once healed there would appear to be some immunity against further outbreaks of excessive reaction, an immunity not to be explained by the increased precautions one would naturally take to avoid a recurrence. The apparatus I employed was modelled on the plan recommended by Schiff and Freund, whose methods I had the advantage of studying in Vienna in the autumn of 1900. Briefly, this apparatus consisted of a resistance board so arranged as to bring the main supply (at 100 volts) down to 12 volts; a mercury break, which could be regulated so as to give a wide range of alternations. As a rule, the apparatus was worked at the rate of 600 interruptions to the minute; about 4 or 5 ampères were used. The coil was capable of giving a 12" spark in air.

Next as regards the tube. Personally, I prefer this to be fairly "soft," and one giving plenty of green fluorescence.

The "zone" round such a tube in action can be felt by the hand to be in a high state of activity, and even at a distance of several inches a distinct "brushing" sensation can be felt. Tubes which have become too "hard" by prolonged use should be temporarily discarded.

The mask is an important item. It should consist of fairly stout lead sheeting, with an aperture in it so arranged as to allow the rays to play upon the diseased part. Tinfoil is sometimes used for making the mask, but it is not to be relied upon. I have seen dermatitis occur under a mask of this description.

Another important point is the condition of the diseased surface. This should be as clean as possible, and all loose scabs and debris removed. Under no circumstances should the parts be covered by ointments containing metallic bases—these in themselves forming formidable masks. I have found a thin smearing of zinc ointment upon a glass plate give a very decided skiagraph.

Turning now to the question of exposure. As mentioned above the distance from the anti-anode to the skin should be about 3 in., and the correct position of the tube is of course important. During the first week or two exposures may be given daily. Afterwards, on the first suggestion of reaction, the patient will only require to sit three times a week—an obvious advantage. In the intervals between the exposures a dressing of moist boracic lint often adds to the patient's comfort and presents none of the disadvantages of some ointments referred to above. It is not wise to attempt the raying of too large a surface at each sitting, and it is always better to devote alternate days to the treatment of two adjacent surfaces, such as the chin and the cheek, rather than to attempt to cover the whole surface each time.

I will now mention the various kinds of skin trouble, for the relief of which by this treatment I have had personal experience. I am much indebted to my colleague, Dr. Brooke, for permission to utilize some of his private cases, the treatment of which with the x-ray apparatus I had undertaken when working with him, as material for the following observations. Other material was found at the Hospital for Skin Diseases, where both my colleagues, Dr. Brooke and Dr. Wild, have for some time had patients under this treatment, a couple of x-ray apparatus being constantly in use for this purpose.

Hypertrichosis.

Hitherto one has only been able to cope with cases of this affection which were amenable to electrolysis; and indeed, for most cases which present themselves for relief, I believe this still to be the best and speediest method. But there remains a class of cases in which one could obviously not employ the needle, by reason of the unmanageable extent of the trouble. It is in this class, to which formerly one has been compelled to deny relief, that one may look hopefully to the x-ray tube for help. My own experience of this field has been limited, but I can point to at least one case in which the treatment has been highly successful so far, and in which I confidently expect a permanent cure. The patient referred to is a blonde in whose cheeks, upper lip, chin, and infra-maxillary regions a multiplicity of thick reddish hairs were conspicuous, while over all there was spread a coarse undergrowth of shorter hairs, the latter condition being obviously irremediable by means of the needle. After a course of treatment extending over some eight months, but with frequent (and sometimes long) intervals in order to allow reaction to subside—though these were in no way ever excessive—the skin is now absolutely smooth and more delicate in texture than formerly. I am continuing the exposures in this case, but with lessening frequency, and so far there is no vestige of a return of the trouble. Schiff has had many of these cases to deal with, and recommends that the treatment be extended over a period of eighteen months in all. The hairs are of course got rid of in a very much shorter space of time, but Schiff insists upon an "after-treatment" if the cure is to be permanent. Moreover, the actual number of sittings is not so large as might be thought. They are merely distributed through a long period.

I have placed the consideration of this condition first, because one is dealing here with healthy skin and can consequently better trace the sequence of events on exposure to the tube. These events are, according to Freund:

1. Pigmentation.
2. Erythema.
3. Inflammatory exudation.
4. Loosening of hairs.
5. Loss of pigment in dark hairs.
6. Subjective symptoms—itching and feeling of tension.

I can fully corroborate the appearance of these signs and symptoms, though by no means in the order quoted above. I should certainly place the erythema first, the pigmentation appearing at a much later date. This pigmentation varies immensely in different people, and is apt at times to cause some alarm on the patient's part, though I have never seen the discoloration last for long. The case above quoted, for instance, has for a long time past been quite free from all traces of pigment, although of course the exposures have been continued. From what I have seen I would unhesitatingly recommend the judicious employment of this method of treatment for exaggerated forms of hirsuties of the class referred to above.

Coccogenic Sycosis.

I have seen several obstinate cases of this affection clear up remarkably well after repeated exposures to the tube. In each case where benefit has accrued there has been first an obvious reaction—a mild dermatitis. It is possible that this inflammatory reaction has proved inimical to the pus organisms. Be this as it may, one has observed the pustules dry up and form scabs, which in due time separate leaving sound skin. It is not necessary that the exposures be pushed to the stage of epilation in order to secure these effects. I am unable to say as yet from personal experience how far these effects are permanent.

Lupus Vulgaris.

In dealing with this disease my experience has been in the main confined to long-standing and severe cases. It is, I believe, in widespread and extensive cases that the use of the x-ray tube is more especially indicated. Cases of less extent, more especially those in which individual nodules or colonies of nodules stand out in isolated patches, can be, I believe, more speedily dealt with by "Finsen's" apparatus. On the other hand, most cases of ulcerated lupus call for x-ray treatment, at least in the first instance.

In dealing with lupus by this method, it is often especially difficult to foretell the extent of the reaction by reason of the disguise the surface wears. If there be healthy skin included within the area of the mask aperture, its condition, of course, affords a valuable index of the stage reached by the treatment. How far should the treatment be pushed? I believe that one can obtain the best results by keeping the reaction within bounds, although I have seen cases where without much warning a mass of morbid tissue has broken down, and where the pain incident to this accident may conceivably have been atoned for by the vast improvement seen on healing. Nevertheless, I believe the object should be to avoid excessive reaction of this kind where possible, seeing that we can get good results comparatively painlessly. I have observed the naked-eye appearances during the treatment of a patch of lupus to be briefly as follows:

The first visible change is an apparent exaggeration of the disease. Any isolated nodules which may happen to be present stand out more prominently and become of a darker hue. Lesions in which the nodules have become fused together get more lumpy and raised and also darker. There is now a feeling of tension in the part, and some itching or slight burning may be complained of. Later on some desquamation occurs. Presuming that appropriate periods of rest are given when the engorgement of the lesions has reached this stage, and that ulceration has been avoided, the subsequent progress is marked by gradually-increasing pallor and softening of the diseased skin. Absorption of the morbid tissue is evidently slowly proceeding, though it may take months before this is finally effected.

It is often a most important matter to decide in a given case what is the maximum number of exposures which ought to be given, or rather what is the longest interval which can be allowed between the sittings? No rule can be laid down on this point which shall apply to all cases, though I believe one is rather more apt to over-do than to under-do the number of exposures. Sometimes exposures of twice or thrice a week would seem to do as well as daily sittings. Whatever

be the process started by the raying, it is certainly a persistent one, and I have several times been struck by the improvement which has been maintained in patients who for holiday purposes or the like have been several weeks away from treatment.

The softening of the skin referred to above is often the first indication of absorption, and may be the first good sign noticed. This was markedly so in a case of mine—that of a middle-aged married lady whose face had been for long hide-bound in an expressionless mask of lupus non-exedens. This patient told me, long before any visible sign of improvement was present, that the skin felt more flexible round the mouth. Later on, as improvement became manifest, it was interesting to observe a gradual return of the normal lines of the features with increasing command of expression. When I last saw this case, early in the present year, a complete cure seemed to be not far off. The nose, lips, and cheeks were free from disease, and mere remnants of the trouble were left in broken streaks on the forehead and over the angles of the jaws. In this case, be it remarked, the reaction had at no time been excessive, and the removal of the disease had been brought about without any ulceration. The softness and healthy appearance of the skin marking the site of former disease is quite remarkable, and there is no treatment which could better have given such complete absence of scarring.

Cases of lupus of the face and neck are often seen where, as the result of disease and of frequent scraping operations, large hypertrophic scars are left which are almost as unsightly as the remaining disease. Such cases are eminently suitable for this treatment, whereby not only the active disease is overcome but the accompanying scars are softened and much improved in appearance.

The great drawback to the application of this method for lupus is, of course, the time and patience required on the patient's part. The treatment is not to be recommended in all cases, even of extensive disease.

I would summarize the cases of lupus suitable for x-ray treatment as being: (1) Cases of disease in prominent parts, such as the face, where the mischief is too extensive for Finsen's method; (2) some ulcerated cases; (3) cases associated with unsightly scars; (4) some cases of disease of mucous membrane.

Rodent Ulcer.

If it were only for the results achieved in dealing with this distressing disease, the x-ray treatment deserves special recognition. Instances of its successful employment in this field are multiplying on all hands, and my own experience has been equally gratifying. One may roughly divide rodent ulcers into two broad groups clinically—the ulcerated and the non-ulcerated. I believe the former class responds more readily to the treatment, although I have seen excellent results with the latter, which I shall describe later. I shall describe first the course of events in a typical ulcerated rodent as I have almost uniformly observed them in cases where the malady has not covered any very large area or extended deeply to underlying structures.

After two or three weeks treatment one can invariably detect some softening in the hard collar-like crater of the ulcer, and there will probably be slight erythema of the skin around coinciding with the mask aperture. Coincident with these changes one can observe the surface of the ulcer alter its appearance. The characteristic waxy pale-red colour gives place to a deeper tint, and soon becomes dark red. At the same time the smoothness disappears, and one can detect small granulations springing up from the surface. These granulations multiply apace, and after a time raise the base of the ulcer flush with the surrounding skin. In one case, that of an old but vigorous man of over 70, who had a rodent ulcer on the nose the size of a shilling piece, these granulations proliferated with extraordinary rapidity, and soon formed a table-land raised one-sixteenth of an inch above the surface, requiring restraint with copper sulphate. While the granulations are forming the border undergoes still further softening and thinning, and finally appears no thicker than normal skin, while a thin blue epithelial lip is seen to be spreading from it over the surface of the ulcer. We thus come to deal with what is apparently a simple healthy ulcer, from which all the characteristics of malignancy have disappeared, and this ulcer gradually heals in the ordinary way. The resulting scar is marvellously smooth and supple. It

may occasionally happen that during the healing process the reaction becomes too strong. In two recent cases of this kind I have seen the newly-formed tissue break down again with distressing rapidity. The treatment was in both instances immediately suspended, and in one of them, when last seen, there was very evident healing again going on, while there was no suggestion of rodent ulcer to the naked eye. The other case was particularly interesting. There had been for some time a thick scab on the deepest part of the ulcer (which involved the upper lip and nasal orifice), and this I had not disturbed, though now I wish I had done so. It was under this scab that the breaking-down process first began, and on removing it I found a considerable quantity of pent-up discharge. This must have been rich in putrefying organisms, and could not but have proved deleterious to the granulation tissue. I am not without hope that this case, which up to the time of the relapse had healed in fully two-thirds of its extent, will in due time recover lost ground.

I have seen only one case—a very long-standing and extensive ulcer—which showed no apparent response whatever to the treatment. The non-ulcerated class of rodent ulcers are by no means insignificant in numbers, and not infrequently remain unrecognized until they have made considerable headway. They are characterized by the pale worm-like ridges, hard to the touch, which mark the advancing border of the disease, which proceeds in slowly-extending waves—their track shown by pale scar tissue more or less distinct.

Cases of this kind which I have observed have shown the usual reaction after a time. The ridges have first shown softening, sometimes being slowly absorbed without ulceration. Generally, however, they have broken down. The ulceration then marks out peculiarly well the sinuous lines which form the site of the thickened epithelium. Elsewhere the scar tissue becomes softer and redder, and finally the patches and streaks of ulceration heal in the ordinary manner.

It still remains to be seen if the "cures" of rodent ulcer by this method will stand the test of time. The treatment is of too recent introduction for anything definitely to be said on this point. I can only remark that two cases of mine, which were discharged from treatment four and six months ago respectively, are still free from visible recurrence. It would probably be a wise precaution in all cases to continue the exposures for a time after apparent cure.

THE CURATIVE EFFECT OF THE X RAYS ON CALLOUS SINUSES OF THE ABDOMINAL WALL.

By D. BERRY HART, M.D.,

Gynaecologist to the Royal Infirmary, Edinburgh, etc.

PERSISTENT sinuses of the abdominal wall where no foreign body is present are a great annoyance both to operator and patient. In two cases I have found these heal with marvellous rapidity after a few exposures to the x rays, and thus think the experience worth recording for further trial. The fact of this action was stumbled on by accident as follows:

A patient, Mrs. S., was sent me from Singapore by Dr. Leask with a persistent sinus in the abdominal wall, after an incision in the right iliac fossa to evacuate bloody fluid of unknown origin. This sinus was 5 to 6 in. long, and ran from near the anterior iliac spine downwards and inwards to the uterus. It had existed for several months. In order, if possible, to see how it ran, I asked Dr. Dawson Turner, of the Edinburgh Infirmary Electrical Department, to take an x-ray photograph (a probe being placed in the sinus and one in the uterus). This did not help one much topographically as to the sinus, but I was struck with the almost magical cicatrization of the sinus after plugging with iodoform gauze for several weeks had failed.

In a second case of more marked sinus persisting for some months in a case of operations for extraperitoneal gestation where fetal bones had been retained for five years,¹ the patient was sent out well but with a narrow sinus into the shrunk sac. This did not heal as I had hoped and the patient was again sent into Ward XXXVI with a sinus admitting the