first also furnishes a good example of the manner in which they sometimes become disseminated—namely, by implantation, and is, as far as I know, the only example of excision of two separate portions of gut for cancer. The first was done with the bowel in a state of distension, and though successful should not be taken as a precedent, for one of the cardinal rules upon which I wish to lay stress is that wherever possible the gut should be efficiently drained when distension exists before an attempt is made to reunite the ends. It is the disregard of this rule which has made the death-rate so high until recently. For the fluid in the distended coils is practically a cultivation of the *B. coli communis* in a high state of virulence, and it is not reasonable to expect the intestine to resume its functions while filled with this. The first case also illustrates the great extension which is added to our field of operations by the use of intravenous injections of saline fluid, with the addition, at the operator's discretion, of strychnine and brandy, for it is impossible to believe that such an operation as that described could have been borne by an exhausted patient suffering from intestinal obstruction without its aid. It also illustrates what I think is a very important point—namely, the method of giving it. It is no good waiting until the anaesthetist tells you the patient is very bad and his pulse failing, and then in a panic sending assistants and nurses scurrying about to get the necessary apparatus to transfuse him. Everything must be arranged beforehand, and the transfusion started at the commencement of the operation and made to extend over the whole time it is in progress.

The second case is an example of the typical and ideal method of proceeding after discovering a malignant tumour of bowel—namely, its removal in three or more stages:

1. Consisting of the delivery and fixation of the growth and contiguous bowel outside the abdomen.
2. Opening and drainage of the gut above the growth.
3. Removal of growth.
4. Freeing and uniting ends of bowel and repair of abdominal wall.

Some of these four stages may, if necessary, be combined—for example, 1 and 2 may be done at one sitting if the need for relieving distension is urgent, but it is always best to allow, if possible, twelve hours to elapse between them, to give the peritoneal surfaces a chance of erecting a barrier against general infection.

Then, again, 2 and 3 may be combined, as in my second case; and this is a very useful combination, and can generally be carried out, as, if you are going to tie in a Paul's tube, you may just as well cut away the growth; it requires no general anaesthetic—packing swabs wrung out of a solution of cocaine round it is quite enough.

Four must always be a separate and carefully planned operation, but the method chosen for uniting the ends is capable of great variation; 200 plans for effecting it have been described, and about 20 of these survive as reasonable and practicable, but I will not prolong this paper with any description of them.

The usefulness and simplicity of any one plan over any other lies not so much in the superiority or the reverse of the method as in the familiarity of the operator with the details of the plan he is employing. Surgeons A. and B. might each be getting excellent results, the one with simple suturing, and the other with the employment of some mechanical aid, as a bone bobbin or metal button, but let A. suddenly adopt B.'s method and B. take A.'s, and they would both be disappointed for a time, chiefly from the extra time taken, for in this, more than any other operation, every minute saved is an important gain, as the severity of the operation is in an almost exact proportion to its length.

One word on the question of anaesthetics in these cases, and I have done. It has been too long considered essential before opening the abdomen to give a general anaesthetic, and the custom dies hard; but in a simple opening for the relief of distension, or where the delivery and fixation of a loop of gut is contemplated, it is a great advantage to the patient, and quite easy to do, after an injection of eucaine.

Of course, I do not for a moment claim this as in any way original, but if it were more generally adopted, especially in cases which have to be relieved at very short notice without any preparation, a great many lives would be saved which are now registered as dying of exhaustion or from the anaesthetic.

NOTE.—In July, 1902, the second patient was seen, and found to be in good health, and again in November, 1902.

THREE CASES OF ACUTE INTUSSUSCEPTION IN THE SAME FAMILY.

By ROBERT G. RIDDELL, M.B., F.R.C.S. EDIN., D.P.H.
Rotherham.

THE following is the history of 3 cases of intussusception occurring in a family of four children, 1 ending fatally and 2 being relieved by operation. I can find no account of a similar series of cases in any of the works on the subject that I have consulted.

CASE I.—This was not diagnosed as such by the medical man then in attendance, or, if so, the mother was not made acquainted with the fact; but from the accounts of the mother and friends there can be little doubt that it was a well-marked case of intussusception. The child was born in 1895, and was brought up chiefly on the breast, but occasionally was allowed to have other kinds of food, and, no doubt owing to this, suffered from attacks of diarrhoea and abdominal pain. When 5 months old it was one evening seized with sudden pain and screamed violently. It vomited all the food it was given, and passed several motions, which the mother describes as looking like apple jelly. All next day it was in great pain, and vomited a quantity of greenish-brown fluid which smelled "like a motion." It also passed a considerable quantity of blood, and became very collapsed and sunken about the eyes. On the third day it died.

CASE II.—This child, a boy, was born in 1900, and was strong and healthy up to the age of seven months, when he began to have the usual digestive troubles connected with teething. One evening a little blood was passed in the motion, and during the night he cried a good deal. Next day he threw up his food, and more blood was passed by the bowel. In the evening the vomit became brown and offensive. Next day more blood was passed, and the sickness continued. The mother at last became alarmed, and called in the doctor, who at once diagnosed intussusception, and had the child removed to the Barnet Victoria Cottage Hospital. Drs. W. Thyne and D. R. G. Corrigan operated, and found an intussusception, which was reduced without much difficulty. The child made a good recovery, and has not since had any serious abdominal trouble.

CASE III.—This child was born in 1901, and, with the exception of an attack of broncho-pneumonia in the spring, had good health till he began to cut his teeth at 7 months, when there was some trouble with the bowels, occasional attacks of diarrhoea alternating with constipation. On May 6th the child seemed quite well till the evening, when some blood was passed in the motion, and an hour later he began to scream and to vomit his food. More blood was passed, and the mother, who by this time had become an expert in these matters, diagnosed intussusception, and brought the child to the surgery. He was then collapsed, the eyes sunken, and the pulse rapid and feeble. The abdomen was not distended or rigid, and it was easy to palpate an elongated lump in the upper part of the right lumbar and umbilical regions. As soon as the necessary preparations could be made an operation was performed. With the assistance of Mr. W. H. Elam, the anaesthetic being given by Dr. W. Thyne, I had not much trouble in reaching the intussusception by a linear incision above the umbilicus sufficient to admit two fingers. It was easily reduced, and the patient suffered little from shock. Recovery was uninterrupted.

It is difficult to account for the occurrence of these cases in the same family. In all three there was a history of digestive troubles due no doubt to careless feeding, accentuated in the last two by dentition, and this is given by most authorities as a frequent cause. Added to this, one would expect that there must have been some congenital condition common to all three, which in the presence of an exciting cause such as the above would be sufficient to determine the occurrence of intussusception. It has been suggested by Leichtenstern that an abnormally movable caecum predisposes to ileo-caecal invagination, while a caecum firmly fastened down may give rise to the ileo-colic variety. One can also imagine that when, from some cause, one part of the bowel is abnormally fixed, and an adjacent movable segment is thrown into violent peristalsis, a similar condition of things is produced to that brought about in the experiments of Nothnagel, where one segment of the bowel was paralysed and an adjacent one stimulated, an intussusception being thereby produced.

None of these conditions were distinguished at the operations, but only the most limited examination was possible (or desirable) through the small incision. One can, however, only come to the conclusion that the occurrence of a somewhat rare malady in three children out of a family of four is to be considered something more than a mere coincidence.

I am indebted to Dr. Thyne for permission to publish these cases.

CASE OF VOLVULUS OF THE CAECUM.

By F. R. B. BISSHOPP, M.A., M.D., M.R.C.P. LOND.,
Tunbridge Wells.

WITH NOTES ON THE OPERATION BY

ALFRED FRIPP, C.V.O., C.B., M.S., F.R.C.S.

Assistant Surgeon to Guy's Hospital.

SIR FREDERICK TREVES, in his *Intestinal Obstruction*¹ says: "More than two-thirds of all cases of volvulus of the intestine are instances of twist of the sigmoid flexure about its mesenteric axis," and he quotes only one case of volvulus of