stones was then felt. These were easily removed easily, until the neck of the gall bladder was reached. Here was found a large and very dense mass, which seemed to be made up of gall bladder, cystic duct, stones, and dense adhesions. The bladder was therefore stripped as cleanly as possible from the duodenum and colon, to both of which it adhered. A large stone was felt impacted in the neck of the gall bladder and was found almost to have ulcerated through. It was somewhat doubtful whether the dense thickening was due to malignant disease, and as, whether simple or malignant, the gall bladder was useless and harmful, it was determined to remove it. This was done, a small portion of the liver also being taken away, the gall bladder being inseparably welded with it. Several bleeding points in the liver were caught by a suture. The cut end of the cystic duct, from which bile flowed freely, was ligatured and a peritoneal flap stitched over. Drainage was employed. For the first three days the patient's condition was very critical; the pulse ran up to 120, and frequent vomiting and hiccough were very distressing. For forty-eight hours—between twenty-four and seventy-two hours after the operation—there was haematemesis, inky-black fluid being vomited every hour or two in quantities of 2 to 4 oz. The vomit was very acid, the corner of the patient's mouth and his lips and chin being scalded by it. The stomach was washed out, and all fluids withheld. High enemata of hot saline solution were given. The vomiting ceased, and for the next three days great progress was made; the bowels acted, there was no vomiting, and fluid, increasing from 15 oz. to 4 pints, was taken each twenty-four hours. On the seventh day the secretion of urine suddenly stopped. Before this, urine had been passed in quantities of 30 to 42 oz. daily. On the seventh day he did not pass urine for fourteen hours, and I therefore introduced a catheter, but drew off only 1 drachm of urine, almost solid with albumen. For forty-eight hours after this, less than 2 oz. of urine passed, in spite of saline infusion, pilocarpin, and other remedies. During this time the patient complained a little of dull aching in the left lumbar region. All the urine drawn off was highly albuminous. On the ninth day I cut down on the left kidney, incised the capsule, and stripped most of it away, and then incised the kidney through the convex outer border into the pelvis. During the next forty-eight hours, 11 oz. of urine were passed, the earlier discharge being slightly blood-stained. No adequate relief, however, was afforded, and the patient became comatose and died eleven and a-half days after the operation. The abdominal wound was soundly healed. Indeed, on the evening of the sixth day I felt assured of the patient's recovery, and I am confident that, but for the suppression of urine, he would have made a good recovery.

CASE V.—Chronic Sclerosing Cholecystitis: Multiple Perforations: Stone in the Common Duct.

History.—Mrs. M. S., aged 51, sent to me by Dr. W. Graddon Mortimer, of Thorne, had suffered for between three and four years from attacks of acute pain in the epigastric and right hypochondriac regions, lasting three or four hours, and then passing gradually away, leaving a sense of stiffness and soreness for a few days. Though these were the first severe attacks, the patient had for many years experienced sensations of discomfort, and grave uneasiness in the epigastrium, which was called "indigestion." In June of this year an exceptionally severe attack was followed, for the first time, by jaundice, and the jaundice has continued up to the present time. During this period—three months and a half—the depth of the tinge had varied; it is less in the morning, and gets gradually deeper during the day. There have been several attacks of shivering and faintness, followed by sweating, and after all of them a deepening of the jaundice has been observed. The motions have been like putty, and the urine has been high-coloured. A floating stone in the common duct was diagnosed.

Operation. October 3rd, 1902.—The abdomen was opened through the right rectus muscle and the upper end of the incision was prolonged towards the xiphoid cartilage. An immense number of adhesions were found and separated. It was then found that three stones, each equal in size to a Barcelona nut, had ulcerated almost through the walls of the gall bladder. Two of these lay in pockets in the omentum, and one was almost hidden in a cavity in the liver. The gall bladder was shrunken. The stones were removed and the gall bladder cleared. A stone was then found in the common duct, and was removed. The rest of the common duct was explored carefully and found to be clear. The gall bladder, which was shut off from the bile channel by an impervious cystic duct, was then removed. Drainage by a split rubber tube and gauze was adopted for forty-eight hours. The patient is now quite well.

CASE VI.—Acute Cholecystitis: Incipient Gangrene of Gall Bladder: Cholecystectomy.

Mrs. B., aged 33, seen with Dr. Rowling.
History.—The previous history of the patient was unimportant, save for the fact that there had been occasional "indigestion." On November 6th, 1902, she went for a long drive on the top of an electric car and felt chilled. On the following day she felt cold, shivered a little at times, and suffered all day from headache. On November 8th the headache persisted. In the early morning there was a large loose stool; there was some abdominal pain, and later in the day there were several attacks of diarrhoea and vomiting. On November 9th she was seen by Dr. Rowling for the first time. The temperature was 102°, the pulse was 90. An abdominal examination proved negative. On November 10th severe pain was experienced, entirely on the right side of the abdomen, and on examination an indefinite fullness was perceived. On the following day this had resolved itself into a definite mass in the right hypochondriac region. The mass increased in size rapidly, and by November 13th it had reached almost to Poupart's ligament. The general condition of the patient became increasingly serious; the temperature never fell below 100°, a little general distension of the abdomen developed; vomiting persisted, and was most painful and distressing, and the bowels were unrelieved. On the morning of November 13th I saw the patient with Dr. Rowling, and diagnosed the case as one of acute cholecystitis, with peritonitis. On account of the increasingly serious aspect of the case, but chiefly because of a severe rigor in the early morning, I was inclined to suspect incipient gangrene of the gall bladder.

Operation.—On Friday, November 14th, the abdomen was opened through the right rectus muscle and the gall bladder exposed. It was deep maroon in colour, with irregular spotted patches of dark green. The gall bladder measured about 7 in. in length, and was tensely distended. On aspiration about 2 oz. of mucopurulent fluid were removed. The walls of the gall bladder were felt to be stiff, and very much thickened. An incision into the fundus gave exit to some small gall stones. After their removal a stone about equal in size to a walnut was taken away. Beyond it and

near the cystic duct a stone was felt encapsuled in the wall of the gall bladder or imbedded in a diverticulum; it was impossible to say precisely. As the gall bladder was almost gangrenous it was decided to remove it. This was done in the manner already described. On account of the intense local inflammation drainage was adopted for two days. The operation was fairly well borne; the patient rallied from her prostrate condition very quickly, and was out of danger in about three or four days. She is now (January, 1903) in good health.

On examining the gall bladder it was found to be intensely congested, and in parts gangrenous. The points of gangrene corresponded to deep ulcerations seen from the inner surface. A stone equal in size to a nutmeg was found in a separate loculus near the junction of the gall bladder and cystic duct.

GALL STONE IN THE COMMON BILE DUCT: CHOLECYSTOTOMY: RECOVERY.

By S. J. ROSS, M.B., CH.B. VICT.,
Assistant Surgeon to the Bedford County Hospital.

I RECORD this case chiefly on account of the age of the patient, the duration of symptoms, and the absence of complications at the time of operation, notwithstanding the long history.

History.

A woman, whose age was 67 years 10 months upon the day of operation, had her first attack of biliary colic at the age of 40 years, just after the birth of her last child. The attacks recurred at irregular intervals up to four years ago. They then became more frequent, and during the past twelve months still more frequent and more severe. During an attack she experienced severe paroxysmal pain in the region of the gall bladder accompanied by vomiting, shivering and jaundice, and fever which lasted for several days. My partner, Dr. Coombs (to whom I am indebted for the previous history, and also for the great interest he has taken in the case subsequent to the operation) advised operation. On November 6th, 1902, I saw the patient towards the end of an attack. She was then slightly jaundiced. In the gall-bladder region was a globular swelling, the lower rounded border of which could be easily defined. It was evidently the distended gall bladder. The patient was anxious for surgical relief. Calcium chloride gr. x was administered for three days thrice daily.

Operation.

On November 10th, the patient having been prepared for operation, Dr. Coombs administered chloroform, and assisted by Mr. Nash I made an incision 4 in. in length in the right semilunar line. Upon opening the peritoneal cavity the distended gall bladder was readily seen, drawn into the wound, and packed round with sterilized flat cloths. The contained bile was then withdrawn by means of a cannula and the gall bladder opened. The stone, which was lying in the common bile duct, was easily felt, pushed onwards into the gall bladder, and delivered. The stone was oval in shape, and weighed 5½ gr. x. The ducts were examined and found free from further obstruction. The edges of the wound in the gall bladder were sutured to the parietal peritoneum. The remainder of the abdominal wound was closed in three layers, a rubber tube was inserted into the gall bladder, and removed at the end of forty-eight hours. Very little blood was lost during the operation.

After-history.

The wound healed by primary union, and I removed all the sutures upon the tenth day. There was never any rise in temperature after the operation. The fistula discharged freely for three days, and then the discharge steadily decreased. It had almost closed on December 19th.

The shape of the stone accounts for the absence of such intense jaundice as usually occurs in these cases. The attacks of biliary colic were doubtless due to axial rotation of the stone in the duct. The attacks of shivering, which are a very constant feature of cases of stone in the common bile duct, are, I think, instances of nervous traumatic fever, analogous to urethral fever following the passage of a catheter, and occurring in cases in which the element of sepsis is entirely excluded.

In conclusion, I think this case illustrates the importance of operative interference in cases of biliary calculus, and even when the age is suggestive of the presence of irremediable conditions associated with the gall stones.

TWO OPERATIONS PERFORMED UPON THE LIVER THROUGH THE CHEST WALL.

By G. P. NEWBOLT, F.R.C.S.,
Honorary Surgeon, Royal Southern Hospital, Liverpool.

GENERALLY speaking, in cases of abscess or hydatid of the liver one is able to evacuate the contents of the cavity by an incision made through the abdominal wall, but there are instances in which it is only possible to do this by resecting a rib and thus reaching the collection. The two following cases are good examples of this condition.

CASE I.—Hydatid of Liver opened through the Chest Wall.

History.—A. F., aged 12, was transferred from the medical wards of the Royal Southern Hospital to my care on August 7th, 1900. She was admitted on July 23rd, 1900, and had been ill for some months suffering from shortness of breath and discomfort in the right side. She had lived on a sheep farm in North Wales.

State on Examination.—The abdomen was swollen in the upper two-thirds,