This statement by Sir Richard Thorne-Thorne of 1898 is still true to-day. The death-rate from phthisis pulmonalis is steadily decreasing, but that from tabes mesenterica is still steadily increasing. On the other hand, we have it stated on high authority that tuberculosis in English cattle is rapidly increasing.

The majority of deaths in infants occur between the ages of 3 and 6 months, just at a time when raw milk is consumed in greatest quantity; from that time the mortality is less and sinks rapidly as age increases.

I have for some time been inclined to the view that tabes mesenterica is not true tuberculosis—that is, that it is not caused by the same tubercle bacillus as that which produces phthisis pulmonalis.

When visiting the splendid laboratories of Professor Koch a few months ago I was much impressed with the distinctive cultural peculiarities of human and bovine tuberculosis, and was bound to admit that the cultures appeared to be distinctive. Professor Koch was kind enough to give me pure, fresh cultures of both human and bovine tuberculosis, and I made several subcultures of both. The growths on blood serum were always distinctive in each case.

From a recent case of tabes mesenterica I made cultivations of the mesenteric gland on blood serum with glycerine, and to my surprise the growth exactly corresponded to bovine tuberculosis or *Perlsucht*. Again, from a case of strumous knee-joint I made a cultivation of the lymph from the interior of the joint, with an exactly similar result. The cultivations which I have previously obtained from phthisical sputum have generally shown the usual characteristics of human tuberculosis and never bovine. Professor Ravenel affirms—

That the tubercle bacillus from bovine sources has in culture fairly constant and persistent peculiarities of growth and morphology, by which it may tentatively be differentiated from that ordinarily found in man.

From these two cases, together with a general survey of clinical facts, I feel there is some reason to suppose that whilst bovine and human tuberculosis are distinct and separate diseases, yet the human subject is susceptible to bovine tuberculosis in the early period of life especially, and that bovine tuberculosis may set up serious lesions in the human body.

I hope in a future paper to be able to substantiate this view after more precise work on tabes mesenterica, strumous glands, and tuberculous joints.

CONCLUSIONS.

In view of the foregoing, I venture to think there is some evidence to assume:

1. That there are two distinct varieties of tuberculosis affecting the human body, one produced by human tubercle, the other produced by bovine tubercle.
2. That human and bovine tuberculosis are separate and distinct.
3. That bovine tuberculosis entering the alimentary canal in milk may set up tabes mesenterica, more especially in children.
4. That bovine tuberculosis is probably the cause of enlarged lymph glands, tuberculous joints, and lupus.
5. That true human tuberculosis or phthisis pulmonalis is always conveyed from one person to another by infection, and generally from advanced cases of phthisis.
6. That every effort should be made to stamp out tuberculosis in cattle, and that milk should be boiled before use by children.

SOIL POLLUTION AND DISEASE IN CAMPS.

By MAJOR R. CALDWELL, F.R.C.S., D.P.H.

THE influence of soil upon health is a matter which has long occupied the attention of the profession, and the interest attaching to this subject has of late been materially heightened by certain speculations relative to the part played by camping grounds in the incidence of enteric fever on armies in the field. These speculations naturally tend to be narrowed to the question as to the influence exerted in this connexion by camping grounds in prolonged occupation by troops.

The soil of a camp which has been occupied for any length of time may be compared to a filter or contact bed for the purification of sewage, in which, owing to lack of rest, or to what would have the same effect, excess of material to be disposed of, the nitrifying organisms are unable to meet necessary requirements, with the production of harmful and offensive

bodies as a result. In the case of the artificial filter or contact bed, not only are periods of rest most carefully regulated, but every effort is made to ensure, as far as possible, that the sewage to be treated should be of a more or less uniform character, as any sudden alteration in its composition is likely to overcome the powers of the living agencies which are utilized for its purification.

In the case of a camp it may fairly be admitted that neither of the above facts receives or, from the nature of the case, can possibly receive, serious practical consideration. Masses of human faeces, horse dung, kitchen refuse, and filth of all sorts are often of necessity thrust *en masse* into the soil, without regard to their ultimate fate once they are covered by a layer of earth of varying thickness.

I have no present intention of discussing questions relative to the variability of bacteria, except in so far as to state that when we consider how external conditions such as food and climate can so affect living protoplasm as unmistakably to alter, within a few generations, the physical and mental characteristics of the human race, we should feel no surprise if lowly organisms such as bacteria, which germinate at such a rate that the "number of individuals which might arise from a single bacterium in three or four days is almost inconceivable" (Hewlett), should develop new and sufficiently remarkable characteristics under the influence of altered surroundings, including possibly the power of producing specific disease. Whether the above theory is correct or not, the practical fact remains that as a general rule in every camp enteric fever and dysentery sooner or later make their appearance. The cause of the trouble is in most cases sought for in the water supply, which, from a sanitary point of view, is usually regarded as the root of all evil when defective, and, on the other hand, when above suspicion, as a panacea for a host of those ills incidental to service in the field.

While it would be difficult to exaggerate the importance of such a vital question, experience shows that water is by no means the sole channel by which diseases such as those under consideration are spread amongst troops. Several cases have come under my notice in which men, who for weeks had been employed in blockhouse duty on the veld, and who had only partaken of water in the form of tea or coffee, and therefore presumably safe, have been attacked by a virulent form of enteric; and officers also during the recent campaign have been victims who have assured me that for many weeks they have never partaken of unboiled water, including in the above term all forms of mineral drinks. For reasons which are too lengthy to be entered on here, the conveyance of enteric fever by milk or food amongst the troops actually in the field, as far as my own experience goes, need not be considered.

A farm, utilized as an outpost in the Harrismith district of the Orange River Colony, was to my personal knowledge a veritable hotbed of enteric fever of the worst stamp, although all drinking water was religiously boiled. The soil, however, had been much fouled by passing columns. In a camp, of which I recently had charge, also in the Harrismith district, enteric fever and dysentery occurred in spite of the fact that the water was of unimpeachable quality.

Many other instances from my personal experience could be given; let it suffice to say that, conversing with other medical officers who have recently been engaged in active service in the field, I have been deeply interested to find that their observations, like my own, point to the constant occurrence of enteric fever under circumstances which justify the absolute elimination of water as a possible cause of the disease in question, but where soil pollution was excessive. I was also much struck, during the earlier stages of the war in Natal, by the fact that men, while trekking, were as a rule singularly free from sickness, but that a return to standing camp was invariably the precursor of a lower standard of health.

The condition of a camp, under which heading blockhouses may for sanitary purposes be classed, must in any event unavoidably tend to the wholesale spread of micro-organisms of all kinds. The constant movement of men, the passage of heavy waggons with horses, cattle, or mules, the necessary trenching and digging, and apart from these, in South Africa, the heavy rainfall, are all factors in disintegration, bringing to the surface organisms which might otherwise have remained harmlessly in the soil. The influence, also, of the movements of ground air and water should not be forgotten. In particular, the effect of movements of ground air in this direction have been most ably pointed out by Sir C. A. Cameron. In his opinion, specific organisms carried into the

atmosphere by this means are one cause of enteric fever in Dublin.

In every camp, also, there is, in spite of all possible efforts at removal, a certain amount of organic matter, largely the result of pollution by animals, which decomposes on or close to the surface of the soil, and, having dried in the sun, is distributed in the form of dust.

The remedy for these evils must vary with the circumstances of the case. Where military considerations permit of such a measure, frequent change of camping ground is naturally to be recommended, but the occasions on which this step is feasible must be most infrequent. The position of individual tents may, however, be readily altered, and this should be done as frequently as possible; the surface covered by a tent rapidly becomes foul, and, the temperature in the interior of the tent—in bell tents, at least—being higher, as a rule, than that of the surrounding atmosphere, must naturally tend to cause undesirable emanations from the soil. The aspirating influence of a blockhouse on the ground air would be greater than that of a tent, and danger to health would be proportionately increased. This defect, as far as blockhouses are concerned, could be remedied by some form of impermeable flooring. Such an arrangement would constitute something more than a theoretical advantage, as the soil pollution tending to occur inside these structures must exercise a marked influence on the health of the occupants.

When the necessary implements and labour are available, buckets may be substituted for trench latrines, and all refuse removed and buried at such a distance from camp as to render pollution of the soil a matter of no practical importance. When the above method cannot be carried out—and this will usually be the case—and sufficient space is available, latrines and refuse pits should be shallow in order to subject their contents rapidly to the influence of nitrification, and frequently filled in, in order not to overtax the agencies on which the process depends. The contents of latrine trenches and rubbish pits dug several feet deep are removed from the influence of those nitrifying agencies, which, it must be remembered, being aerobic, are to be found in the upper layers of the soil only. The conditions in such a case probably resemble those found in a septic tank, liquefaction of organic matter taking place under the influence of anaerobic organisms. Percolation under these circumstances would seriously endanger neighbouring water supplies. These disadvantages must, however, at times be overlooked, and under some circumstances need never be considered. In actual practice every method is found to have its disadvantages, and even with the greatest care and with unremitting attention all organic matter will, as already stated, be neither removed nor buried, as a certain amount—the result largely of faecal pollution by cattle—must under ordinary circumstances remain adherent to the surface of the soil.

Much may be done to minimize risk; but, having regard to the ordinary circumstances of warfare and to the present state of our knowledge, a certain degree of danger remains, which must be regarded as an unavoidable incident of active service in the field.

THE SUPPLY OF PURE WATER TO TROOPS IN THE FIELD.

By JAMES FERGUSON LEES, M.D.,

Late Medical Officer 1st Battalion the Connaught Rangers, South African Field Force.

OWING to the liability of filters to go wrong or break, as well as the immense labour necessary to produce a sufficient supply of pure water in this way, purification by filtration is unfitted for general use in the field. Chemical purification is also, *per se*, of little use, since the addition of any chemical agent in sufficient quantity to destroy infection would be inadmissible.

The most efficacious method of sterilizing water is undoubtedly by boiling, the main objection to this method being the length of time required for cooling before the water is fit to drink; but a constant supply of water, sterilized in this way and at any time available, might be attained by preparing each night in advance the water supply for the next day.

With this object in view, I have designed the water-cart shown in the accompanying diagrams.

On arrival at the final halting place for the night, the water-carts would be driven up to the source from which it was proposed to draw the following day's supply of water, and the settling tanks removed by a man on either side of

each cart catching hold by the lugs, G, and swinging the tank with its attached furnace over to the ground, using the hinge, K, as a fulcrum. When the tank is in position on the ground—as shown in dotted lines in Fig. 1—the hinge at K is open and the tank then loose from the cart.

Fig. 1.—Water-cart consisting of an iron tank B, the carrying tank, fixed on the body of the cart, provided with a tap, T, and covered by a lid, L, which should be bolted down, but at the same time easily removable, for occasionally cleansing the tank. Attached to the carrying tank is a semi-rotary hand-pump, C, with a suction-pipe S, and a discharge-pipe D. This pump is worked by the handle K, which fits over the square-head E. Outside and inverted over the carrying tank is another tank, A, called the settling tank, provided with a furnace, F, and hinged to the cart at K by an open hinge. Lugs, G, are attached to the settling tank on either side for lifting it.

The water-carts, with their carrying tanks full of pure water previously prepared at the last night's halting place, would then drive to their allotted positions in the camp, and from them the whole water supply for cooking and drinking would be drawn. In the meantime, the settling tanks would have been filled with water from the proposed source, fires lit in the furnaces, and the water in each boiled for half-an-hour.

Should the water be thick with suspended organic or mineral matter, alum (6 gr. to the gallon of water) should now be added to precipitate this. A previous addition of iron perchloride ($2\frac{1}{2}$ gr. to the gallon) may be necessary for this precipitation should there be no lime present in the water. The water should be left standing in the tanks all night to allow of it cooling and settling.

FIG. 2

FIG. 3

Fig. 2.—The water cart with the settling tank removed.

Fig. 3.—The settling tank on the ground. P, a pipe for drawing off the water fitted with a perforated rose, R, covered with an asbestos filtering cloth.

In the morning, by connecting the pipe, P, with the hand-pump at S by means of a flexible hose, the supernatant clear water in the settling tanks could be drawn off into the water-carts in a few minutes. The settling tanks would then be cleaned out and placed again in position over the carrying tanks.

By thus preparing each night a quantity sufficient for the next twenty-four hours, a supply of pure water which would be available at any time during the march or on arrival at