there was insufficient evidence to show what was the cause of the inflammation.

In the most recent textbooks I have failed to find much guidance on the matter to enable me to conclude what is the pathology of the condition. In the *Clinical Journal*, however, for August 6th, 13th, and 20th, 1902, are three lectures delivered at the London Polyclinic by Mr. F. C. Wallis, entitled the *Diagnosis and Treatment of Diseases of the Rectum*. In speaking of ulceration of the rectum, he describes a condition which he says is "the commonest form of ulceration which has come under his notice" since his connexion with St. Mark's Hospital, and which is due, in his opinion, to sepsis. In his own words: "By this I mean ulceration due to some septic discharge from the outside attacking the greater portion of the mucous membrane, and so progressing; or, what was a common cause in the past, post-operative sepsis." In many respects there was a resemblance between this case and those described by Mr. Wallis as being of septic origin. There was, however, an absence of some other signs which, as described in the lectures, should have been present if the ulceration was of this nature. Thus there was no oedema of the muco-cutaneous margin around the external sphincter, with fissuring and ulceration between the oedematous portions.

The virulence of the inflammation as evidenced by the large slough which came away strongly suggested an acute septic process, and in the absence of any other discoverable cause, the question arose whether the gonorrhoeal urethritis previously mentioned had any bearing on the rectal ulceration.

I felt that the best chance of quickly improving his condition was to produce complete and immediate physiological rest in the rectum, and this could only be secured by colotomy. I therefore advised that this should be done, and through the courtesy of Mr. Joseph Thompson he was admitted into the General Hospital under my care. On March 3rd I did the first stage in the operation for inguinal colotomy by bringing a portion of the descending colon out through a small wound in the abdominal parietes, and fixing it there by passing a glass rod through its mesentery, the ends of the rod resting on each side of the wound, on the belly wall. There being no stitches used and no distension existing, as so often is the case when a colotomy has to be performed, this operation is perhaps the simplest in surgery, and occupies three or four minutes only. Three days later I opened the bowel in the wound, and the faeces were subsequently discharged through the opening thus made above the ulcerated rectum, which latter was by this proceeding placed at complete physiological rest.

For about a week after the colon had been opened I had the rectum washed through once daily, and on three or four occasions 1 oz. of iodoform emulsion was thrown into it after the irrigation.

There is but little to add to this part of the case, and at the end of less than a month all the ulceration was healed, and he was discharged from the hospital wearing a colotomy belt.

Before he went home I informed him that should he wish it I would close the colotomy wound later on, but a few weeks' additional rest would do the rectum no harm in view of the severity of the inflammation to which it had been subjected.

As he was 72 years of age I thought it doubtful if he would desire any further operation, but he came to me in August saying that the opening was a nuisance and prevented him working, and he was readmitted to have it closed. This was done on August 26th by the method described by the late Mr. Greig Smith, without opening the general peritoneal cavity. The upper opening in the colon I first shut off by a plug of cotton wool, so as to prevent soiling during the manipulations, and an incision having been made through the skin just outside the margin where the mucous membrane joined the skin, this was deepened until subperitoneal fat was reached. The finger was then able to strip the peritoneum off the inside of the abdominal parietes in the neighbourhood of the wound to the extent desired, so that it, with the attached colon with its opening in it, could be delivered well outside the belly. A continuous suture was now applied, after trimming the edges of the opening in the gut, which took up mucous membrane only and closed the bowel in a transverse direction. A series of Lambert sutures was then placed over this, thereby inverting the first union, and the gut was closed. The muscles were brought together with buried silk sutures, a tiny drainage tube being placed just beneath them in case any leakage

from the bowel occurred, and the skin united with fishing gut.

There was never any leakage from the bowel, and the tube was removed next day. The whole thing was healed eight days after the operation and the stitches removed. There was some difficulty in getting the bowels to act at first, but this was eventually overcome by the use of enemata and some castor oil on the eighth day after the operation, and since then they have acted normally each day.

At the present time (November 19th) he is perfectly well and at work regularly, and there is moreover not the slightest indication of any narrowing of the lumen of the gut at the place where the ulceration existed.

He was cured of his disease solely by the application of "physiological rest" to the diseased part; and I am of opinion that no other agency which we could have employed would have so quickly, so effectively, and so satisfactorily produced this effect.

CASE OF PUERPERAL HYPERPYREXIA OF MALARIAL ORIGIN.

By THOMAS WILLIAM CLAY, L.R.C.P., L.R.C.S.,
Holyhead.

THE following case seems to be of sufficient interest to call for publication.

Mrs. A., aged 28, a primipara, was delivered by forceps of a male child at 6 a.m. on September 20th, 1900. The delivery was effected with but little difficulty; the perineum was slightly torn and was stitched. Progress was in every respect satisfactory, the last visit being made on the fifth day when the temperature was, as it had been all along, normal. On the morning of September 26th (seventh day) I received an urgent message to attend immediately. On my arrival, two hours after the onset of the attack, I was informed that the patient had been suddenly taken with a very severe rigor, felt icy cold, and soon afterwards became very hot, but that now she was feeling better. I found her bathed in perspiration, with a temperature of 102.4° F. After careful examination I utterly failed to satisfy myself as to the cause of the attack. The stitches were removed, the perineum being quite healed. The vagina was washed out with an antiseptic solution. Although the nurse told me that the bowels had acted the day previously, an enema of soap and water was given, with the result that a huge motion was passed. Three hours from the onset of the first rigor, and in my presence, she again had a rigor of such severity that her teeth chattered and the bed shook. Her temperature immediately began to rise, and rapidly reached 107.8° F.; she was now delirious, the pulse so weak and rapid that it could not be counted; in fact, she appeared to be *in extremis*. Stimulants were freely administered; a bath was not available, but ice was immediately procured from a neighbouring hotel; it was applied to the head, rubbed and applied directly to the skin of the abdomen and thorax. The effect was most gratifying; the temperature at once began to fall, and when 103.4° F. the ice was discontinued. The patient was dried and wrapped in warm blankets, perspired freely, and felt much better. Quinine in doses of 4 gr. was given every three hours. She had two or three slight shivers after this. The temperature, intermittent in character, never again rose higher than 103° F., and was normal in four days. The bowels acted naturally the day following the enema, the quantity being again quite exceptionally large. After this she made a good recovery.

The case caused me a considerable amount of anxiety, and I pondered a great deal over the possible nature and causation of the attack. The entire absence of all local signs of septic infection, as well as the non-elevation of the temperature until the seventh day, to my mind practically excluded a diagnosis of puerperal septicaemia. The quite unusual and extraordinary loaded state of the intestine appealed to me as a possible cause, absorption from the bowel, being, I believe, recognized as a cause of severe febrile disturbance during the puerperium.

The same patient, in April, 1902, after an easy, non-instrumental labour, gave birth to a second child. Things went on well until the evening of the twelfth day, when I was hurriedly summoned. The husband, who was much alarmed, informed me that she had had an attack resembling that after the first confinement; she complained of feeling very cold, shivered, and on taking the temperature he found it to be over 103° F. Arriving an hour later I found her perspiring freely, with a temperature of 101° F.

Remembering how puzzled I had been about the cause of the disturbance after the first confinement, and finding the patient again in a condition bearing such a close resemblance to the previous attack (also the absence this time of constipation), whilst at the bedside the cold, hot, and sweating

characters of the two attacks suddenly occurred to me. Malaria suggested itself, and I found, on questioning, that the patient had not only resided abroad, but that four years previously she had suffered from a severe attack of malarial fever.

Acting on this discovery, a dose of quinine (10 gr.) was given, followed by 4 gr. every three hours; she had no more rigors; the temperature was normal in twenty-four hours. Here, then, probably was the explanation which in the first attack was unsuspected. Malarial puerperal fever presents, I believe, some difficulty in diagnosis from puerperal septicaemia: but the main points of distinction were well brought out in this case. First, in both attacks, the onset of fever not occurring until after the fifth day, probably always before this in septicaemia. Secondly, the particularly well-marked rigors and intermission of temperature, much more pronounced than in septicaemia. Thirdly, the short duration of the symptoms. The loaded state of the bowel possibly was a factor which contributed to the intensity of the first attack.

The severity of the first attack—the rarity, at all events, in this country of this condition—and the possibility that others like myself may overlook malaria as a cause of fever in the puerperium, are my reasons for wishing to record this case.

RESEARCHES ON HIBBERTIA VOLUBILIS.

By JOHN REID, M.A., M.D., C.M.,
Southfields, S.W.

THE above plant was identified by my old and esteemed friend the late Baron Sir F. von Müller, who kindly assisted me in the bibliography, and told me that, as the order had been little investigated, my work would serve to indicate the general properties of its members. Tannin and silica are the products found.

The specimens used were obtained at Coraki, on the Richmond River and Manly Beach in N.S.W., in the months of January and February; the pistil and fruit with envelope removed were extracted after being dried.

The alkaloid dillennine may be extracted by acetic acid or spirit by the usual methods. The plant residue is then boiled in methylated spirit to yield the oil, and the residue with alkali to yield dillennic acid, from whose solution a colouring matter—stephenine, named after the late Rev. Geo. Stephen, Fordyce, Banffshire—may be skimmed off, an acid from an alkaline fluid. There is also a brown-coloured neutral resinoid obtained along with the alkaloid, when the latter is precipitated by ammonia from solution.

Dillennine is a white powder depositing acicular crystals from solution in chloroform. Its acetate forms acicular crystals, the dillennate broken plates. Solution in spirit diminishes the colouring power of tincture of guaiacum on potato surface. It is slightly volatile, decomposing partly with the odour of urea or burning feathers—charring. The usual alkaloid reactions are given, with phosphomolybdic acid, $KIHgI_2$, and iodurated potassium iodide. Given internally, the tongue becomes dry and then moist; the circulation is enfeebled; no head symptoms appear. The pupils are dilated, remaining so for a few hours. There seems to be no stimulation of the liver.

Dillennic acid precipitates with lead and lime water, from which precipitates it may be obtained. It is very insoluble in water; its solution does not precipitate iron perchloride, gum, or gelatine. It decomposes sodium bicarbonate, intensifying the sherry-red colour of its solution. Ammonia makes the colour still deeper. Alkaline solutions are very soluble. An alkaline solution gives a dark ink-like solution with iron salts to transmitted light, with a lighter colour to reflected light. Silver nitrate causes a dark grey precipitate, insoluble in dilute nitric acid but readily soluble in ammonia.

Stains of silver on the fingers are readily removed by dillennic acid and ammonia. Lead in a neutral or vegetable acid solution partially precipitates, in alkaline solution entirely. Alum partially precipitates, lime water entirely or lime salts partially, the lime precipitate being redissolved by acetic acid, etc. Potash solution is partly precipitated by acetic acid; the residue gives a precipitate with HNO_3 ; its residue a precipitate with alum. The acid burns without odour, and seems to smoulder away when heated in a glass tube.

Solution of the acid very speedily caused tetanic spasms in

a common earthworm and death, the worm *post mortem* presenting a flattened appearance. In man internally it seems to act violently on the liver, causing black motions of a watery character, with some griping and nausea. There is no irritant or astringent effect on the mouth, etc. The solution of alkaline salt or acid keeps readily, hence it has antiseptic properties worthy of investigation.

Stephenine, like carmine, dissolves readily in ammonia, and is a nuclear stain; it is soluble in glycerine, insoluble in spirit. It can be readily mounted in dammar, and will stain silk and paper of a rose colour. The oil, soluble in boiling, insoluble in cold methylated spirit, gives the sickly odour of the plant. It is solid and, like essential oils, is non-saponifiable.

BIBLIOGRAPHY.

Phyto. Aust., xl, 94: Müller also, viii, 125; xl, 94. Ersch and Gruber. *Encyc. Bentham, Aust. Flor.*, i, 19. Rosenthal's *Synopsis Plantarum Diaphoricarum*.

HUMAN AND BOVINE TUBERCULOSIS.

A PRELIMINARY NOTE.

By NATHAN RAW, M.D., M.R.C.P. LOND.,

British Member of the International Committee for the Prevention of Consumption; Physician, Mill Road Infirmary, Liverpool; Physician, Sanatorium for Consumption, Heswall.

THE controversy amongst scientific workers as to the exact relationship between human and bovine tuberculosis is still active, and the startling statement of Koch at the International Congress in London in 1901 has, if it has as yet led to no other result, at least stimulated men to active research throughout the world. Professor Koch only this year in Berlin has reiterated his theory, and is as firmly convinced as ever he was that the two diseases—namely, human tuberculosis and bovine tuberculosis or *Perlsucht*, are quite distinct and different, and that bovine tuberculosis cannot be conveyed to the human.

The question is too large to be discussed here, and it is, of course, quite beyond my powers, in the absence of careful and prolonged experimentation, to offer a solution, but although I was taught to believe that the two diseases were really identical, previous to the paper by Koch in 1901, I have after careful observation entirely changed my view, an opinion which is based chiefly on clinical observation.

During the last five years, namely, from 1898 to 1902, I have had under my care in Mill Road Infirmary 1,750 patients suffering from various forms of phthisis pulmonalis, and 273 children suffering from *tabes mesenterica*. In addition to these cases, there have been a great number of patients affected with varying forms of tuberculosis of joints, bones, glands, and skin, as would be expected in a large Poor-law infirmary containing 800 beds.

Careful *post-mortem* examination with bacteriological and microscopical observations have been carried out in the laboratory, together with inoculation experiments conducted at the Thompson-Yates Laboratories, University College.

Up to the present time it has been customary to classify all the lesions called tuberculous in one general category of disease, and to say they were all produced by one bacillus, namely, the tubercle bacillus of Koch. The chief of these lesions are:

Phthisis pulmonalis.
Tuberculosis of pleura.
" " peritoneum.
" " lymph glands.
" " joints.
" " bone.
" " skin.
" " other organs.

If we consider for a moment the disease called *tabes mesenterica*, which is classified as tuberculosis, we must ask ourselves the questions, "How does the bacillus gain entrance to the mesenteric glands?" and "Why is the disease not decreasing with other forms of tuberculosis?"

The late Sir Richard Thorne-Thorne in his Harben lecture of 1893, says:

The matter, too, assumes a still more serious aspect if we limit ourselves to the first year of life, when raw milk is most largely used as food, for then we find that the reduction in the rate of death from the various forms of tuberculosis, which reduction has been going on at all ages for about half a century, not only disappears, but is actually transformed into a large increase, reaching no less than 27.7 per cent.

This in itself is grave enough, but its significance is still further emphasized when we remember what are the circumstances under which this increase in the rate of death from *tabes mesenterica* has gone on synchronously with a decrease in that from other forms of tuberculosis.