forward case in a rather desperate condition. An incision was at once made over the caecum, and a quantity of greenish gangrenous fluid at once escaped with flakes of lymph. I then drew gently on a mass presenting in the wound, and at once a knuckle of small intestine came forward from behind the caecum in a state of advanced gangrene and quite rotten. This was secured in the incision and opened there, or rather it could be wiped away. All this was done in a very short time; but the relief came too late to arrest the general abdominal sepsis, and the patient died. Cases are, of course, on record where this condition has been suspected or diagnosed and relieved successfully by operation.

Broken-down caseating glands in the right iliac fossa (No. 6), simulating a subacute appendicitis, are not so uncommon. Here the patient has probably had antecedent attacks of pain and more or less trouble with the bowels from the localized tuberculous peritonitis. One case from among a number will illustrate the difficulty of diagnosis. Some years ago I was asked to see a little girl, 10 years of age, in the country, who had been ill for some time with all the symptoms of subacute epityphlitis. The temperature was up and down: there was a diffused swelling in the right iliac fossa, not very painful. On opening over the caecum, I at first thought that I had really found a thickened appendix containing concretions lying among adherent intestines and omentum. But on closer examination this proved to be a chain of caseating glands running from the caecum through the mesentery. Some had burst, some were just breaking down. I removed as many as I thought could be safely taken away, and the patient recovered after a tedious convalescence.

No. 7 (ileocaecal carcinoma with abscess) must be a fairly uncommon condition, and yet I remember such a case several years ago here. It seemed typical, and there was a large abscess over the anterior iliac spine. This I opened and drained. It was not until the woman's death a long time afterwards that the necropsy revealed the fact that there was a ring of cancer in the valve and a perforation which had led to the abscess.

But I remember a much more puzzling case which I had to operate on two years ago down in Kent. It was that of a lady of 55 who had a round and perfectly movable tumour in the right iliac fossa with some symptoms of obstruction not acute and little or no fever. Here I was uncertain until I had opened the abdomen as to whether we had not to deal with a malignant growth of the ileocaecal valve. And even while I was clearing the puckered and hard mass I was preparing for an enterectomy. It turned out, however, that the mass was a chronically inflamed appendix included in tough adhesions. Finally I cleared the appendix, which was very long, thick, and hard removed it, and closed the abdomen. Recovery was perfect, and I have heard within the last few days—while seeing another case in her town—that she is quite well. Other cases I have seen suggested the same difficulty.

Remember that there are cases of appendical inflammation which are quite latent. A patient may have a lump in the iliac fossa with few or no symptoms of appendicitis, with little or no tenderness and no raised temperature, and it may last for months in this condition and yet after all turn out to be a suppurating appendix. Some of you must remember a strong, healthy, country labourer, aged 26, from Surrey, who has just left the hospital. He looked perfectly healthy but had for some time had an occasional pain in the right iliac fossa. When he arrived here he had no trouble except the lump and no fever at all, and yet he had a considerable abscess containing three stercoliths behind the caecum. And at the present time, and since this lecture was delivered, I have a still more striking case under my care in private. A particularly healthy, ruddy, and well-nourished gentleman of 45 came to me a couple of weeks ago at his doctor's request to consult me about a lump in his right groin the size of a large egg. This he had had for many months. It was smooth and painless, and very hard. It could be moved easily from side to side, and only when pressed far inwards was there a sense of dragging. His temperature was steadily a little under normal. Beyond dyspepsia and an occasional attack of flatulence and diarrhoea he had no symptoms of intestinal derangement, and he went about his ordinary work without trouble. I was quite prepared at his age to find a new growth, but when I operated five days ago I found an appendical abscess containing about half an ounce of pus.

In reference to No. 8 you might ask how could a haematoma of the broad ligament simulate appendicitis? And yet it can closely. Some years ago I was called up here to operate on a case of acute appendicitis in a little girl, aged 10 years. She had been seized some hours before with violent pain in the

right side of the pelvis, had vomited, become distended and tender, and her temperature was high. There was a considerable swelling on the right side above Poupart's ligament, which was very tender. Every one of those who saw the case thought it was a typical appendicitis. When I opened the abdomen on the right side there ran out a quantity of blood-stained fluid, of which the pelvis was full. The appendix was perfectly normal, and there was no peritonitis. But the right broad ligament was distended with a large black haematoma, from which the serum had exuded. I only wiped the pelvis dry, and returned the broad ligament to its place, and all went well. I suppose this was an abortive attempt at menstruation. The case is fully recorded in the *Transactions of the Clinical Society*.

As to No. 9, I only recall one case in which reduction of a hernia *en masse* had to be considered in this connexion. It has already been alluded to in discussing the case of perforated gastric ulcer. But in that case it was certainly not out of the question.

No. 10 (intussusception) ought never to be mistaken for appendicitis. And yet I was called into Surrey some years ago by a very shrewd practitioner, whose telegram ran, "Intussusception case; please come at once prepared to operate." I went, but found the case to be one of acute appendicitis in a boy of 6. On this I operated with an excellent result, the appendix was removed, and a quantity of inflammatory fluid drained off, recovery following without much trouble.

I cannot follow up this subject further just now. The mere enumeration of the conditions which may simulate appendicitis and the relation of one or two cases illustrating each has led us far enough to-day. On some future occasion I may try to sift the cases and offer you clues which may help in differential diagnosis. But just now my aim has been to remind you that the diagnosis of appendicitis must be approached with great care. In these days when we are operating on very large numbers of typhilitic cases in a routine way I think we need to be reminded of this from time to time. The ultimate diagnosis will in many cases only be arrived at when the abdomen is opened. But if we do not recognize the possibility of finding perhaps any of the conditions mentioned above and have not prepared our minds beforehand to deal with any of them, valuable time may be lost and mistakes in manipulation may occur. Moreover, we may not have made proper preparations for the operation beforehand, and so be landed in difficulties.

The Erasmus Wilson Lectures

ON THE
PATHOLOGY AND SYMPTOMS OF ADENOCARCINOMA OF THE RECTUM.

Delivered before the Royal College of Surgeons of England.

By SIR CHARLES B. BALL, M.CH., F.R.C.S.I.,
HON. F.R.C.S. ENG.,

Regius Professor of Surgery in the University of Dublin.

LECTURE II.—ADENOCARCINOMA. PATHOLOGY.

As in a simple adenoma, the essential element in adenocarcinoma is the fact that it springs from and involves the mucous membrane, it has already been pointed out that the only positive difference is that the malignant form penetrates the basement membrane and becomes widely diffused, it is therefore possible to assert positively that a tumour of the rectum which does not involve the mucous membrane is not an ordinary cancer; and as quite 95 per cent. of the cases of rectal carcinoma are of the adenomatous type, the importance of this clinical point is obvious.

The degree of malignancy of a cancer as indicated by the tendency to recur after operation we know well varies greatly with its situation; for instance, an epithelioma of the lip is less malignant than the same disease occurring in the tongue; and epithelioma on the scrotum less malignant than when it occurs upon the penis. What then is the relative malignancy of cancer of the rectum to similar disease in other parts of the body? If the case comes under notice before the disease has completely penetrated the intestinal tunic, the prospect of permanent cure by removal is very good indeed, unfortunately, however,

it is usual to find when the case is first diagnosed that the disease has already extended beyond the limits of the rectal tunic and then it must be admitted the malignancy is as great as in any other region of the body.

There are occasionally to be found within the rectum malignant growths which are not adeno-carcinoma, such as the various forms of sarcoma and in the lower portion of the anal canal squamous epithelioma is sometimes seen, all are rare, and put together probably do not make up 5 per cent. of the entire number of cases of malignant disease met with in this region.

Adeno-carcinoma, like the benign variety, is so easily diagnosed in the rectum as compared with other portions of the intestinal tract that its relative frequency in this region appears to be greater than it really is; it is, however, probable that it is actually much more common in the lower bowel than elsewhere.

Like other forms of carcinoma, adeno-carcinoma of the intestine is met with more frequently after middle age, but quite a number of cases are recorded in which it commenced in comparatively early life. I have myself seen cases at the age of 16 and 18 years respectively. When met with before middle age it is usual for the case to be highly malignant and run a rapid course.

Both sexes appear to be equally affected; recent large statistics tend to show that the incidence of malignant disease of the intestines in males and females is practically identical.

It is hard to estimate usefully the probable duration of these cases, as owing to the trivial early symptoms it is usually quite impossible to fix the date of commencement, while in the latter stages death frequently ensues from the onset of what may be called accidental complications. Roughly speaking, however, it may be said that from the first occurrence of symptoms sufficiently marked to justify a diagnosis until death, the average duration in unoperated cases is two years.

CLINICAL TYPES.

Clinically adeno-carcinoma, as met with in the rectum, presents one of two tolerably distinct types: (1) Commences

Fig. 1.—Tuberous adeno-carcinoma of the rectum, which has in the centre broken down into a craterform ulcer, and spread almost round the entire circumference of the bowel, but has not as yet extended to the perirectal tissues.

as a flat tubercle in the mucous and submucous tissue; it increases somewhat rapidly by exuberant growth at the margin, while the centre breaks down and forms a craterform ulcer; it tends to spread round the circumference of the bowel somewhat more rapidly than it extends in the vertical direction, thus making an oval ulcer. Perforation of the muscular coat occurs early, with involvement of the surrounding structures and pelvic lymphatics, and the disease runs a somewhat rapid course. This is the form more commonly met with in comparatively young subjects. (2) Commences as a more superficial ulceration, and has not the exuberant edges met with in the other variety; the adenomatous tissue of the new growth is less in quantity and more atrophic, but there is a considerable formation of dense connective tissue surrounding the ulcerated surface, which tends to contract and narrow the lumen of the bowel, producing intestinal obstruction, which indeed may be the first symptom to draw attention to the disease; this apparently is the schirro-contracted rectum of the older writers, it runs a much more chronic course, and is usually met with at a more advanced age than the first variety.

Fig. 2.—Atrophic adeno-carcinoma of the rectum producing tight stricture; on the right side enlarged lymphatic glands are seen beneath the peritoneum.

MORBID HISTOLOGY.

If a section be made of the growing edge of an adeno-carcinoma it will be found that there is a considerable pro-

Fig. 3.—Photomicrograph of the edge of a tuberous adeno-carcinoma ($\times 20$); to the left normal mucous membrane is seen and in the centre the growth of adenomatous tissue in the submucosa, the basement membrane having been perforated.

liferation of the mucous glands, that the adenomatous tissue breaks through the muscularis mucosae and spreads in the submucous tissue separating the bundles of muscular fibres, the essential characteristic by which adeno-carcinoma is distinguished from the non-malignant form, it eventually perforates the entire thickness of the intestinal tunic and spreads in the tissues of the pelvis, any of which may become implicated in the growth.

The lymphatics soon become involved, first the glands at the back of the rectum, then those along the great iliac vessels, the glands in the groin escaping involvement except where the skin of the anal canal is encroached upon, or at a late period where there is a widely diffused contamination of the lymphatic system. Next in

order of frequency to the lymphatic system, metastatic growths are met with in the liver, and they occasionally have been observed in the lung and elsewhere, but wherever found

Fig. 4.—Photomicrograph of adeno-carcinoma of rectum ($\times 30$) the adenomatous tissue is seen invading the circular muscular coat of the intestinal tissue.

they reproduce accurately the original type of disease—namely, an arrangement of adenomatous tissue similar to that met with normally in the Lieberkühn follicles of the intestine (see Fig. 5).

Fig. 5.—Photomicrograph ($\times 50$) of secondary adeno-carcinoma in lung, the primary disease having been situated in the rectum. To the left above, the characteristic adenomatous tissue is seen.

Associated with adeno-carcinoma, it is not uncommon to meet with polypoid growths (Figs. 6 and 7) which, from their pedunculated growth and freedom from involvement of the deeper structures are essentially benign tumours, these may either be of the tuberculated surface commonly found, or the tumour may be split up so as to be formed of a mass of filiform processes—the so-called villous growth—it appears probable that these neoplasms are the result of the direct irritation of the discharge from the cancerous ulcer.

Fig. 6.—Simple adenoma found in connexion with a typical adeno-carcinoma. The benign character of the growth is seen by the way the muscular coat has separated from the mucous coat in preparing the specimen.

Fig. 7.—Simple villous adenoma in connexion with a typical adeno-carcinoma of the rectum. Photomicrograph ($\times 5$).

COLLOID DEGENERATION.

In a small percentage of cases it will be found that adeno-

Fig. 8.—Colloid adeno-carcinoma of rectum producing tight stricture, the dilatation of the intestine above the stricture is well marked.

carcinoma of the rectum has undergone colloid degeneration, which is characterized by a somewhat wide infiltration of the tissues by a smooth gelatinous material; clinically the importance of this condition is that it is liable to be mistaken for tertiary syphilis, but the latter is usually characterized by much greater density and firmness to the touch, and it more frequently implicates the anal canal.

Fig. 9.—Photomicrograph (x 15) of colloid adeno-carcinoma of rectum, showing penetration of the muscularis mucosae and infiltration of the submucosa with colloid material.

SYMPTOMS.

Diarrhoea.—The early symptoms of cancer of the rectum may be of the most trivial character, and as the hope of successful surgical treatment depends altogether upon early diagnosis, it is important to pay great attention to the slightest indications of this terrible disease; unfortunately a large proportion of cases when they first consult a surgeon are so advanced that complete removal is impracticable. Usually the first symptom to attract attention is a slight chronic diarrhoea, more marked in the morning. If the stools are inspected the motion is found mixed with mucus, which later on is generally blood-stained, and not infrequently the small motion will be found to consist of mucus and blood only. This large increase of mucus is due to the great increase of gland tissue which is such a characteristic of the disease, and as soon as the surface becomes ulcerated the mucus is mixed with blood. It is essential that all cases of diarrhoea lasting more than a few days, especially if mucus or blood is present in the dejecta, should be submitted to a careful digital examination of the rectum. In this way only can early diagnosis be arrived at.

Obstruction.—As the lumen of the bowel becomes encroached upon, obstruction becomes marked, and if the disease is situated low down the motions are never properly formed, being tape-like or in little pieces; and although there may be slight discharge of bloody mucus with thin faeces several times daily, increasing doses of purgatives become necessary to effect a satisfactory faecal evacuation, the abdomen becomes distended, and the colon loaded with masses of scybala. As the disease progresses more positive symptoms become prominent, but at the same time they usually indicate such extension of disease that complete removal has ceased to be possible.

Pain.—In the early stages pain is not a prominent symptom, unless the lower section of the anal canal is involved, when it may be acute; but this is extremely rare. In the latter stages, when the disease has spread into the other pelvic structures, pain may be severe, especially if the sacral plexus of nerves be encroached upon, or the bladder opened. Involvement of the bladder is first indicated by frequent micturition, which becomes extremely painful.

once fistula has formed, the escape of faeces into the bladder setting up a violent cystitis, while the masses may be sufficient to block the urethra and so cause retention of urine, or the first symptom may be passage of flatus per urethram: a truly miserable condition urgently demanding the operation of colotomy. In like manner in the female the vagina or uterus may be opened into, forming faecal fistula, while an opening up of the loose areolar tissue of the pelvis may set up stercoral abscess, demanding external opening in either buttock. Among the rarer complications due to the spread of the disease in the pelvis may be mentioned oedema of one leg from involvement of the iliac vein, or hydronephrosis from blocking of one of the ureters.

DIGITAL EXAMINATION.

In all cases where the remotest suspicion of cancer of the rectum exists digital examination should be insisted upon, and for this purpose it is desirable to have the bowel as completely emptied as possible. The well-lubricated finger is to be introduced as far as possible, and each portion of the surface of mucous membrane carefully searched, in the large majority of cases when the characteristic symptoms are present the disease will be readily detected, as the most common situation is easily within reach of the finger in the lower portion of the rectal ampulla, a rough irregular mass can usually be felt obviously originating in the mucous membrane, sometimes confined to one aspect of the bowel but more often surrounding the entire tube, and when this is the case the sensation conveyed to the finger is like that of an irregular and craggy cervix uteri, with an opening in the centre like the os, if in an early stage of obstruction the tip of the finger can be passed through this, and possibly feel healthy mucous membrane above. If the disease is situated higher up more difficulty may be experienced in the diagnosis, sometimes the surgeon may feel that there is a mass a little higher up that can be actually touched with the finger, folds of mucous membrane intervening and preventing the characteristic sensation being obtained. In these cases additional information may be obtained by getting the patient to stand up and strain down, or, still better, by the bimanual method used by gynaecologists, conducted, if necessary, under an anaesthetic. An adeno-carcinoma which

Fig. 10.—Adenocarcinoma of ileocaecal valve producing an intussusception which extended to the rectum, and was removed by excision after protrusion through the anus. The vermiform appendix and divided ileum are seen above.

originates high up the bowel may sometimes be found in the rectum having produced an intussusception, as in the case from which Fig. 10 is taken. A woman, aged 75, came under notice for complete intestinal obstruction, and a digital examination of the rectum revealed the presence of a malignant mass, as the finger could be readily passed round the circumference of the tumour without feeling any attachment to the wall of the rectum. The diagnosis of intussusception was readily made, and with little difficulty the growth was drawn down through the anus and removed by circular excision. The first intimation of the site of origin was the

appearance of the vermiform appendix when the peritoneum was opened. The intussusception had evidently been a very chronic one as there were no inflammatory adhesions. The divided ileum was sutured to one angle of the divided colon and the rest of the colon wound closed, the line of junction being returned into the rectum. The patient, who appeared to be little disturbed by the operation, suddenly sat up in bed some ten hours afterwards, and fell back dead.

Besides the mere recognition of the disease, much more must be made out at the preliminary examination. First, as to mobility: We can recognize three degrees of mobility of the tumour. First it is freely mobile, and can be pushed up and down in the pelvis; this is highly important, as indicating that the disease is still confined to the rectal tunic, and is eminently satisfactory for surgical treatment. Secondly, it can be moved to a more limited extent, apparently carrying with it some of the other pelvic structures; this indicates that the disease has spread through the coats of the bowel, and if this has occurred to only a limited extent there is a possibility of removing the tumour, but the prognosis is not at all as good in reference to recurrence as in the first case. In the third case the disease is absolutely fixed, indicating that the disease has spread so far that the bony pelvis has become involved; when this has occurred, it usually means that attempts at complete removal will be quite futile.

Next to the character of the growth itself, the condition of the surrounding structures in the pelvis have to be examined—notably the hollow of the sacrum searched for enlarged lymphatic glands, the condition of the prostate in the male, and the uterus and adnexa in the female. Lastly, as the finger is being withdrawn, the length of healthy tissue below the disease and the condition of the anal canal should be estimated. Having completed the digital examination of the rectum, the abdomen should be carefully palpated, secondary tumours in the liver sought for, and also enlarged glands along the course of the iliac vessels; finally, the patient's general condition, especially the state of the kidneys, is to be inquired into, and having completed this the surgeon is in a position to express an opinion as to whether the case is one suitable for excision. If freely movable, and no enlarged glands or metastatic growths can be felt, no matter how high up the disease is situated, provided the general condition of the patient is good, the case is one eminently suitable for removal. If there is limited adhesion to the vagina, or portion of the pelvic diaphragm, or if lymphatic involvement is confined to a few small glands in the hollow of the sacrum, operation may be undertaken although the prognosis as to recurrence must be estimated as unfavourable, while if the disease has involved the prostate, bladder, or uterus; has widely involved the lateral or posterior aspects of the pelvis; is attended with extensive involvement of the lymphatic system; or still more if metastatic growths can be detected in the abdomen, operation must be absolutely declined.

SCHOOL HYGIENE IN FRANCE.—The French League of Medical Practitioners and Families for the Improvement of the Physical and Intellectual Hygiene of Schools, the formation of which was announced in the BRITISH MEDICAL JOURNAL some time ago, recently held its first annual meeting, under the presidency of Dr. P. Le Gendre, who was assisted by Dr. A. Mathieu, the General Secretary. The Minister of Public Instruction was represented by M. Rabier, Director of Secondary Education. The meeting passed the following resolutions: (1) That an official organ of the League, to be entitled *L'Hygiène Scolaire*, should be founded; (2) that a National Congress of Physiological Paedagogy should be organized; and (3) that the Committee of the League should be instructed to urge on the Departmental Councils the need for the effective organization of medical inspection of schools in all departments in which, notwithstanding the obligation imposed by the law of 1886 and a decree of 1887, no such inspection exists. M. Rabier congratulated the organizers of the League on the success of their undertaking, and assured them that their representations would always receive favourable attention.

An Irish patriotic banquet will be held at the Trocadero Restaurant, W., on Monday, March 16th, the eve of St. Patrick. The chair will be taken by Dr. C. J. Barry, who has practised in London for the last twenty-five years, and is a member of the Fulham Board of Guardians. Tickets for the banquet, at which the usual loyal toasts will be given, may be obtained at Keith, Prowse, and Co.'s box office; or from Mr. David Barry, 449, Fulham Road, S.W. The price of letters to admit lady or gentleman is half a guinea.

CASE OF DILATATION OF THE PYLORUS FOR SIMPLE STRICTURE.

By HORATIO P. SYMONDS, F.R.C.S. EDIN.,
Surgeon to the Radcliffe Infirmary, Oxford.

THE case to be described was that of a man, aged 40, who was admitted to the Radcliffe Infirmary, Oxford, on March 15th, 1902, suffering from dilatation of the stomach. He had been ill, off and on, for eleven years, suffering from great flatulent distension of the stomach, with sickness. The latter symptom did not occur frequently, but when it did, the vomit was great in amount, and had a very putrid smell. When it happened it usually came on a little time after taking food. The patient also suffered from chronic constipation; the bowels never being properly relieved except after the use of an enema. He had constantly to leave off work. He was a fairly well nourished man. No tumour could be felt in the abdomen, but by percussion the stomach was found to be very much distended. There was no history of vomiting of blood. This last fact, together with the long continuance of the symptoms, appeared to exclude malignant disease and to point to a simple stricture of the pylorus.

The man's condition of discomfort was so great that he asked if no operation could be undertaken for his relief, and I therefore determined to try the effect of dilatation of the pylorus. For a week previous to the operation the patient was kept on a bland diet, and received 8 gr. of benzoc-naphthol every four hours.

On March 25th the abdomen was opened through the rectus muscle below the xiphoid cartilage. The stomach was found markedly distended. The pylorus was steadied with the left hand, and with the index and middle fingers of the right a portion of the anterior wall of the stomach was invaginated through the pyloric orifice, and dilatation of the latter was carefully accomplished by separating the fingers. The anterior wall of the stomach was then drawn out through the abdominal wound, and a longitudinal tuck was run in it by a continuous suture of the applied peritoneal coats. This tuck included that part of the stomach which had been used in the invagination manipulation. The abdominal wound was then closed. The following day the patient vomited a little blood, but otherwise made an uneventful recovery. He remained in hospital for a month on a careful diet, and then went home. Since the middle of August he has been at work, has had no stomach symptoms, and expresses himself as feeling a different man.

The operation for dilatation of the pylorus for simple stricture is usually associated with the name of Loreta, who seems to have used two procedures. In the operation as usually attributed to him the viscus is opened and the pylorus dilated by the direct introduction of the finger into it. Wharton and Curtis in their *Surgery*, however describe Loreta's operation as having been performed by him in the manner in which I operated in the above case. Loreta is said to have performed the operation thirty times. It has been twice performed with a fatal result in New York, and Mr. Hagyard has recorded a successful case in which the operation was performed by himself. In this case the stomach was opened and there was some troublesome bleeding from the dilated pylorus. The patient made a good recovery, but there appears to have been little relief of symptoms. Cheyne and Burghard, in mentioning the operation, condemn the invagination method on account of the serious results which may follow in that part of the stomach which has been pushed through the pylorus. As I performed the operation, however, the invaginated part is included in the tuck which is run in the stomach wall, and as it no doubt becomes digested and thus destroyed, no evil effects can follow so long as the suturing of the peritoneum in the tuck has been properly performed.

A SIMPLE METHOD OF OPERATING ON PILES.

By A. B. MITCHELL, M.B., F.R.C.S.I.,

Surgeon to the Royal Victoria Hospital, Belfast, and to the Ulster Hospital for Children and Women.

THE following method of operating on haemorrhoids, which I have employed for the past four years, has in my hands proved both simple and efficient. It is quite possible that other operators have adopted the same principle, but I have never seen it described.

The sphincter having been dilated and the piles brought fully into view, the mucous membrane is sponged with a solution of corrosive sublimate 1 in 1,000.