quite well up to two days before admission. This patient was a Nubian, who took part in the mutiny of 1898, since when he had been in the Entebbe Gaol. He, like the others, had never been outside the Uganda Protectorate, and had no recollection of being bitten by flies of any sort. His temperature rising to over 103° the same afternoon, gradually sank, almost reaching normal on the fourth morning after admission, on which day it rose again above 100° in the afternoon, and then became subnormal on the fifth day. The headache and oedema also disappeared on the fifth day. The pulse in this, and also in the first two cases, rapid at first, gradually resumed a normal rate. The parasites, more numerous at first than in the first two cases, diminished in number on the fourth day, and on the fifth day disappeared entirely.

REMARKS.

The parasite, in both fresh and stained preparations, is apparently exactly similar in all respects to the trypanosoma described by Dr. Dutton, and also to that which I saw in Dr. Manson's case at the London School of Tropical Medicine last November. Unfortunately undue pressure of other work has prevented my making blood counts in sufficient quantity to be worth recording, though, in the few attempts I made at differential count of leucocytes, there was increase in the percentage of lymphocytes and large mononuclear cells and decrease in the polymorphonuclear cells. I have all these cases still under observation, and shall therefore be able to note any fresh manifestation of symptoms which may occur. They are interesting as showing that the trypanosoma, hitherto found in man only in West Africa, has a wider geographical distribution. They are also remarkable in comparison with Dutton's and Manson's papers on account of the short duration of the symptoms, the absence of hurried respiration, and of wasting, and in the first two cases, of splenic enlargement and oedema of the skin. One is led to think that a certain proportion of cases of short duration, such as I have met with here, in which microscopical examination of the blood reveals no malarial parasites, might reasonably be due to a trypanosoma infection where the number of parasites is small (as in Case II), and therefore easily overlooked. Another point worthy of note is that these cases have occurred in a country reputed to be free from tsetse fly disease among cattle, though the natives have brought me in a fly which Colonel Bruce has assured me is undoubtedly of the tsetse fly group, and he has kindly forwarded specimens to the British Museum for further identification. I have been fortunate in being able to show these cases to Dr. Castellani, Colonel Bruce, R.A.M.C., and Dr. Nabarro, of the Royal Society's Commission for the Sleeping Sickness; and I am indebted to Dr. Moffat for permission to forward these notes.

THE USE OF ACID PHOSPHATE OF SODIUM IN ALKALINITY OF THE URINE.

By ROBERT HUTCHISON, M.D. EDIN., M.R.C.P.,

Assistant Physician to the London Hospital, and to the Hospital for Sick Children, Great Ormond Street.

A FEW years ago, whilst investigating the behaviour of acids and alkalies in the body, I had occasion to make some experiments on the influence of different drugs on the reaction of the urine. In particular I tried to determine what agents are best adapted to increase the acidity of the renal secretion. The method I adopted was to estimate the acidity of the urine secreted during the night by an individual on a fairly fixed diet on several successive occasions, so as to obtain a standard of normal acidity. The substance the effect of which was to be tested was then administered at bedtime, again on several successive occasions, and the average urinary acidity obtained compared with the standard already observed under normal conditions. The acidity of the urine in these experiments was estimated in two ways:

1. By the method of Freund, which consists in throwing down the alkaline phosphates by barium chloride and subsequently estimating the acid phosphates by titration with uranium nitrate solution.

2. By direct titration with standard alkali solution, phenolphthalein being taken as the indicator.

I am aware that the latter method is open to objection as a means of determining the absolute acidity, but it gives quite trustworthy results for comparative purposes, and, as a matter

of fact, I found that it always afforded the same indications as the more accurate method of Freund above described.

My first observations were made with mineral acids—hydrochloric, sulphuric, and phosphoric; I found that the influence of these, even in the largest doses that could be conveniently administered, was extremely slight. The reason for this, no doubt, is that mineral acids are neutralized in the liver or circulation partly by fixed alkalies, partly by ammonium carbonate, and excreted in the form of the corresponding salt. In accordance with this view, I found that the ammonia excretion was always greatly increased after the administration of those substances.

2. Organic acids, of which I tested tartaric, acetic, citric, and lactic, had a slight but distinct tendency to increase the acidity of the urine. On the other hand, the effect was never very great, some of the acid being neutralized as above, as evidenced by an increased excretion of ammonia, and some probably being oxidized in the body and excreted in the form of carbonate.

3. Benzoic and boric acids had no influence on the reaction of the urine.

4. Acid salts were next tried, especially acid phosphate of sodium (NaH_2PO_4), for, that salt being the chief cause of the acid reaction of normal urine, one might reasonably expect its administration to have some effect. This expectation proved to be correct, for sodium acid phosphate had a more direct and marked influence on the reaction of the urine than any other substance tried.

Some of the results are shown in the following table, which is compiled from the averages of many observations:

Influence of Medicinal Doses of Various Substances on the Acidity of the Urine (Average of Several Observations).

	Amount of Urine Secreted.	No. of c.cm. of N_{10}NaOH required to Neutralize 10 c.cm. of Urine.
Normal average	240 c.cm.	5.7
$\frac{1}{2}$ dr. ac. phosphor dil.	230 "	6.6
2 oz., $\frac{1}{2}$ per cent. lactic acid	345 "	5.5
2 oz., $\frac{1}{2}$ per cent. acetic acid	390 "	4.3
10 gr. benzoic acid	368 "	5.3
45 gr. tartaric acid	270 "	6.7
45 gr. sodium acid phosphate	250 "	10.0

One may state the result from sodium acid phosphate in terms of Freund's method as follows:

	C.cm. of Uranium Solution required to Precipitate Acid Phosphates in 100 c.cm. Urine.	C.cm. of Uranium Solution required to Precipitate Alkaline Phosphates in 100 c.cm. Urine.
Normal average	46	23
Average after administration of 60 gr. of sodium acid phosphate	92	41

It will be observed that by no means all the acid phosphate given is excreted in that form, but that a considerable proportion is neutralized in the body. This is also shown by the fact that the ammonia output is decidedly raised after the administration of the acid phosphate.

Thus the total ammonia excretion, stated in terms of $\text{N}_{10}\text{H}_2\text{SO}_4$ is: Normal, 148; after 60 gr. sodium acid phosphate, 170. The ammonia increase, however, is always more marked when either a mineral or an organic acid is given.

Having observed these results, one was encouraged to try the effect of acid sodium phosphate in cases of alkaline urine. This I have done in a great many cases, and the results show that the power of the drug to increase the acidity of the urine is quite manifest. It would be tedious to recite cases, but it will be sufficient to say that it has now been used pretty largely in the London Hospital during the last two years in cases of alkalinity of the urine from cystitis and other causes with very satisfactory results. I do not wish to claim too much for it. I do not say that it will immediately render an ammoniacal urine acid, or that it can be depended upon to bring about that result in the severest cases, but I think it will often be found of great assistance. My colleague, Mr. Barnard, has already expressed the high opinion he has formed of the acid phosphate as an aid in keeping the urine

acid after operations on the bladder²), and my chief object in writing this paper has been to secure for it, if possible, a more extended trial.

Sodium acid phosphate has the advantage of being very soluble in water and easily administered. It may be given in doses of from 30 to 60 gr. every three hours, but it is better to give smaller quantities oftener so as to secure continuous elimination. A convenient plan is to dissolve 2 drachms in a pint of water and allow the patient to drink small quantities of this from time to time. The only unpleasant effect which I have ever seen it produce has been diarrhoea, but that can be at once checked by suspending the administration and giving a dose or two of chalk mixture.

I should add that sodium acid phosphate will be found a useful adjuvant to urotropin, for the latter, as is well known, acts best in an acid urine.

REFERENCES.

¹ See also *Virchow's Archiv*, 125, p. 566, 1891. ² *Lancet*, November 11th, 1901

CLINICAL OBSERVATIONS ON THE TREATMENT OF INOPERABLE CANCER BY FORMALIN.*

By A. F. MEREDITH POWELL, M.B. & C.M. EDIN.,
Minster, Thanet.

[ABSTRACT.]

FORMALIN, as commercially known, is a 40 per cent. solution in water of formic aldehyde made by passing a mixture of methyl alcohol vapour and air over heated platinum and conveying it into water. This preparation is a powerful, highly-diffusible, non-toxic antiseptic and deodorant, possessed of escharotic qualities which must be guarded against in using it. It also possesses the property of hardening tissues, and I may here state that I have taken advantage of all its properties except its escharotic quality in attaining the results I propose to describe. I was led to try my present method from my observation, twelve months ago, that formalin 2 per cent. solution had the property of coagulating egg albumen when whole eggs were immersed in it for a few weeks. Thinking over this action of formalin, and knowing its hardening properties on tissues generally, I argued that if I hardened the morbid tissue, if possible without seriously interfering with the healthy tissues, I should convert the growth into a foreign body, and Nature would proceed to dislodge it. My inference has, I venture to say, proved correct.

In from three to seven days after application the cancer mass begins to detach itself in the same way that necrosed tissue separates in gangrene. A line of demarcation is formed at the edge of the malignant mass, and separation takes place in a few days, the time varying with the size and nature of the disease.

Some twenty-four to forty-eight hours after the first application the objectionable foul-smelling discharges cease, thus showing the powerful deodorizing quality of formalin even in 2 per cent. solution. In proportion as the fetid discharges lessen the patient becomes more bright and cheerful. In from twelve to sixteen days I was able to remove the growths in three cases—a sarcoma of the breast, a scirrhus of the breast, and an epithelioma of the lip (recurrent), leaving in its stead healthy granulating tissue, the skin gradually growing and closing the wound up. I have repeatedly reapplied formalin during the healing process without any effect on the granulation tissue left, thus showing that no signs of the original growth remained, for skin will not grow over unhealthy tissue. The process of separating is completed by the aid of a pair of forceps and scissors to snip the fibrous bands that pass into the underlying granulations. It would appear that a fight for supremacy had been going on between the healthy tissue and morbid growth, the former being unable to dislodge its saprophytic enemy until the formalin reduced the greater vitality of the malignant mass, then Nature proved itself superior in the hitherto unequal fight.

The strength of formalin solution should be between $\frac{1}{2}$ per cent. and 2 per cent. If weaker than this the hardening process is not satisfactory, and the application must be continued for a longer period, thus losing time, and the process is extremely painful to the patient. Likewise if 2 per cent. is exceeded the application is painful, the diseased mass becomes surface hardened, separation is difficult, and the risk

of destroying the skin around becomes greater the stronger the solution used, until the escharotic qualities of the formalin manifest themselves.

The method of application is as follows: Absorbent lint soaked in 2 per cent. formalin solution (made by adding 19 parts distilled water to 1 of commercial formalin) and laid on the tumour; this is covered with jaconet and cotton wool and bandaged on. The dressing should be changed six-hourly. After the third or fourth dressing the discharges and fetor ceases; the further process is an aseptic one. In three to seven days the tumour loses its elasticity and becomes darkened, friable, and insensitive—the further use of formalin is painless, and separation takes place as above described. Formalin 2 per cent. solution appears to exert a selective power on morbid tissue, or perhaps it is that the tissue, being more tender by virtue of its younger age and rapid growth, is less resistive to its action than the more mature tissue. It also acts as a powerful stimulant, thus aiding Nature to get rid of its unwelcome parasite. The treatment of the healing surface resolved itself into the ordinary method with alternate boric fomentations or sal alembroth gauze as occasion suggested itself.

Dr. J. D. McFeeley, of Dublin, read a paper at the Surgical Section of the British Medical Association last year on the treatment of rodent ulcers with formalin (published in the *BRITISH MEDICAL JOURNAL* of November 8th). His method and mine are widely different; and I may here state I devised my plan of treatment quite independently, not having seen his published papers till my attention was drawn to it lately. McFeeley avails himself of the escharotic qualities of formalin, which he applies, full strength, under an anaesthetic, either local or general, and keeping his patient under the influence of morphine until the escharotic action is produced; the hardened tissue formed is subsequently allowed to slough off. The disadvantage of his plan of treatment appears to me to be that the whole of the disease may not be attacked (in case of a large sarcoma); the leathery eschar formed takes five or six weeks to slough off, it is therefore not an aseptic process; it is extremely painful to the patient; there is the risk of destroying healthy surrounding skin, in short, the process is similar in all respects to what obtains when chloride of zinc is used.

My method is practically painless to the patient: with it no narcotics are required, and no anaesthetics local or general being necessary. The formalin has time to select other foci should such exist. The whole of the malignant mass is permeated and hardened throughout, and after separation only healthy tissue is left. No sepsis occurs, and the fetid smell is entirely destroyed after a few hours' treatment. The malignant mass is completely removed in sixteen to twenty-four days.

[Several cases were shown illustrating the results obtained by this formalin method.]

GUAIACOL IN THE TREATMENT OF SMALL-POX.

By J. J. RIDGE, M.D. AND M.D. (STATE MED.), B.S., B.A.,
B.Sc. LOND.,
Enfield.

THE continued prevalence of small-pox in some parts of England prompts me to report the results and treatment of a small number of cases which came under my care last year as medical superintendent of the temporary small-pox hospital erected by the Enfield Urban District Council.

The results may be presented briefly in a tabular form, giving those treated at Enfield and a nearly similar number sent to South Mimms Small-pox Hospital.

Enfield.				South Mimms.			
	Recovered.	Died.	Total.		Recovered.	Died.	Total.
Adults ...	34, 1 unvac.	1, vac.	35	Adults ...	47, 4 unvac.	3, vac.	50
Children	9, 4 unvac.	4, unvac.	13	Children	3, 2 unvac.	1, unvac.	4

The ages of the children treated at Enfield were five under 1 year; one between 1 and 2; three, 3 years; one, 4; one, 7; one, 8; and one, 9. Those that died were 10 months, 4 months, 4 months, and 2 weeks.

* Read before a meeting of the South-Eastern Branch of the British Medical Association.