length of the inspiratory and expiratory sounds is not modified in health; (c) the vocal resonance, especially in men, is more marked on the right side; (d) vocal fremitus is more marked on the right side.

3. Of these differences the increase in the vocal fremitus is by far the most constant. Differences in the percussion note are only noted in half the cases examined.

4. The differences, except in percussion, are not so much alterations in the character of the signs as modifications in transmission of the sounds.

5. The modifications in the signs are apparently due to the direction and form of the main bronchus, which are not the same in the two lungs. The percussion note modification may possibly be influenced by the thicker covering of the right side of the chest, but is most probably also dependent upon the position and size of the bronchus.

CONSUMPTION IN THE NAVY.

By W. E. HOME, M.D., M.R.C.P.E.,
Staff Surgeon.

MR. SICHEL'S important paper on Consumption in the Navy, read at the annual meeting of the Association at Manchester, published in the BRITISH MEDICAL JOURNAL of October 4th is full of interest, and his demonstration of the waste of money and loss of efficiency produced by the disease is most valuable.

Personally, I think that he has underestimated the damage, as he has only considered cases returned as tuberculous. The Blue Book for 1900 gives in addition, at page 9, materials from which we may calculate the total deaths ascribed to phthisis, and these must be added to the "tuberculous," that we may have complete statistics of deaths from tuberculous infection. Diseases of the respiratory system are reported on page 9, and at the end of the very valuable annual discussion of the prevalence of these diseases in the training service, it is said that in that service about half (really two-thirds) the invalids from respiratory diseases and about a quarter of the deaths are due to phthisis, probably tuberculous. If this proportion obtains in the service generally, we must add to the 49 deaths which occurred in 1900 from tuberculous disease one-fourth of the 106 deaths reported to have resulted from respiratory disease, that is, 26, and thus we get 75 as the total number of deaths in the service from tubercle in 1900, a rate of 0.79 per 1,000. This increased proportion gives even more weight to Mr. Sichel's contention that there is still room for improvement in the navy.

The loss of the training service from tuberculous diseases is nowhere shown in the Blue Book, but in the last ten years there has been on an average a yearly death-rate of 0.4 per 1,000 from phthisis and pleurisy, which may be pretty safely grouped together. The rate has been falling of recent years, such cases being increasingly returned as tuberculous. There is some reason for believing that in 1900 deaths from phthisis were proportionately more numerous in the training ships than in the general service, but there were only 3 in the training ships. In the three years 1896-7-8 the ratio per 1,000 of invalids and deaths for phthisis pure and simple was higher in the training service (5 per 1,000 invalids and 3 per 1,000 deaths) than in the general service (3 per 1,000 invalids and 1 per 1,000 deaths). This is a serious consideration when it is remembered how lately those boys were passed into the service as free from physical defect.

Owing to alterations made in the form of the health statistics of the navy it is no longer possible to compare the death-rate of this disease in the army and navy, but from their respective reports for 1896, 1897, 1898 the deaths in each service on the large home station (in each case involving about half the men of the service) may be extracted. To get a complete account of the deaths from tubercle I add to those specified as "tuberculous" those also ascribed to phthisis, hæmoptysis, hæmorrhage of the lung, pleurisy, empyema, and broncho-pneumonia. We then find an average:—Army, force at home, 97,666; deaths, 67; rate per 1,000, 0.68; 1896-8; navy, force at home, 34,481; deaths, 31; rate per 1,000, 0.9; and as our men are better clothed and fed, I suppose the excess must be due to their closer aggregation in small cubic space, as Mr. Sichel believes; in fact, to the greater chance our life offers for the spread of infection. To reduce this risk we must diminish overcrowding by improving ventilation, and as in the management of other infectious diseases we must strive for early recognition and isolation or removal of

the patient. Mr. Sichel has done a most useful work in bringing this matter before the Association, and has greatly lessened the difficulties of securing increased prophylaxis by his skilful investigation and telling announcement of the loss to the navy caused by tubercle, furnishing us with a financial argument for improved sanitation which cannot but arrest the attention of those who concern themselves with the defence of the empire.

OBSERVATIONS ON EMPYEMATA IN CHILDREN.

By P. STANLEY BLAKER, M.R.C.S.ENG., L.R.C.P.LOND.,
Late Resident Medical Officer, East London Hospital for Children,
Shadwell, E.

In this paper I have put together the facts and impressions gathered from a personal experience of 81 cases of empyemata in children.

I think it is on the whole admitted that the disease during childhood is a less serious complaint than during adult life, but this statement does not hold good for all ages of childhood. It will not be denied, for instance, that the disease is a very serious complaint during infant life—that is, when it attacks a child under 2 years of age; under a year old, as the figures below will show, the recoveries are few and far between. Hence a record of a personal experience of the above number of cases may be deemed of value, and more especially the points relating to treatment.

I range myself with those who believe that an empyema is an empyema from the beginning—that it starts as such, and very seldom as a simple effusion, which later on becomes purulent. Only in one case did a secondary puncture (after an interval of a fortnight) of the chest reveal pus when the first exploration withdrew a syringe of clear pleural fluid. In this case, moreover, I am rather inclined to the belief that pyogenic organisms were introduced into the chest at the first operation. In every other case pus was found from the beginning.

Classification.

If I were to arrange all the 81 cases in some order of classification, I should, from a clinical standpoint, be disposed to divide them up into two main groups—those that I would associate with the name of "acute" or "primary," and the "late" or "secondary," which latter would include about 80 or 85 per cent. of all cases. To explain this classification, I would begin by saying that nearly all empyemata seen at a children's hospital are associated with an attack of pneumonia, either of the croupous or of the catarrhal form. An empyema due to initial disease of the pleura alone must be a very rare condition indeed, and then it would probably be associated with a similar condition of the other serous membranes or with some form of pyæmia. In the 81 cases to which reference is made, it was not found possible to attribute any one of them to an absolute primary morbid condition of the pleura. Further, many cases that were examined *post mortem* showed some trace of associated pneumonia. Of course, it is difficult to imagine how the lung tissue could escape implication when the contiguous pleura was diseased and vice versa, and hence it would be practically impossible to say which was the seat of the primary mischief. Now, the affection of the pleura is manifested either concomitantly with the attack of pneumonia or else subsequently to it, this period varying from ten days to three or four weeks, and even much longer; and hence those in the former group would be classed as acute or primary, and those in the latter as chronic or secondary.

In the acute cases the signs of fluid in the chest develop simultaneously with those of pneumonia, and the patient is very ill indeed from the first onset of the disease. The illness starts quite suddenly with vomiting and marked dyspnoea, the respirations are hurried and grunting, the *alae nasi* work vigorously, there is a good deal of recession of the lower part of the chest, the face is dusky and the lips markedly cyanotic; the temperature is found to be raised to 104° or 105°. On making a physical examination of the chest, there is marked dullness with increased resistance on one or both sides, and loud tubular breathing over a very extensive area. A puncture with an exploring needle withdraws a thin watery purulent fluid, very often blood-stained. Such cases practically always die when the patients are under 2 years old. They may live for a few days or as long as a fortnight after the onset, but they eventually die, not from the empyema, but mostly from extensive consolidation of the lungs, if they escape such complications as pericarditis, meningitis, and peritonitis.