Remarks

ON THE

OPERATIVE TREATMENT OF CHRONIC FACIAL PALSY OF PERIPHERAL ORIGIN.

(With Special Plate)

BY

CHARLES A. BALLANCE, M.S., F.R.C.S.,

Senior Assistant Surgeon and Lecturer on Surgery, St. Thomas's Hospital;
Surgeon to the National Hospital for the Paralysed
and Epileptic, Queen Square;

HAMILTON A. BALLANCE, M.S., F.R.C.S.,

Assistant Surgeon to the Norfolk and Norwich Hospital, Norwich;

AND

PURVES STEWART, M.A., M.D., M.R.C.P.,

Assistant Physician to the Westminster Hospital.

THE prognosis in facial palsy of extracerebral origin is a very variable one. Whilst in the majority of so-called "rheumatic" cases the prospects of recovery are good, yet in other cases, and especially those due to traumatism or to involvement of the Fallopian aqueduct by a suppurative otitis media, recovery may not occur and the patient is doomed to lifelong disfigurement.

The following series of cases is an attempt to deal with the uncured residue by performing an anastomosis between another healthy nerve and the distal segment of the paralysed facial. The rationale of this procedure is based to a large extent upon the conclusions arrived at in a work published by two of us over a year ago,¹ according to which regeneration occurs in the distal segment of a divided nerve, even when separated from the central. Such regeneration, however, does not reach full maturity unless the distal segment is joined to the proximal, so as to permit of transmission of impulses between the nerve centres and the periphery.

According to the other view, hitherto the more popular one, regeneration occurs by a process of downgrowth from the central into the distal segment. The latter plays an entirely passive part, and unless reunited becomes reduced after a time to a mere fibrous cord along which the downgrowth of nerve fibres from the central segment towards the periphery with any hope of reaching the muscles is practically impossible.

If, on the other hand, as we have demonstrated, regeneration occurs (immature, it is true) in the distal segment itself, no interval of time ought to be too long for attempted reunion so long as any muscle fibres survive which can be innervated by the regenerated and reunited nerve. Thus, for example, in one of our cases the paralysis had existed for almost three years and yet recovery subsequently occurred. And we see no reason on *a priori* grounds why this limit should not be exceeded, providing always, as already stated, muscle fibres are still present to be innervated by the recovering nerve.

The method of procedure in the present series of cases was as follows:—Having first assured ourselves by galvanic stimulation that muscle fibres still survived on the paralysed side of the face, the facial nerve was exposed by operation at its point of exit from the stylo-mastoid foramen. The nerve trunk was cut across as high up as possible, and a minute portion of the distal segment was reserved for microscopic examination. The spinal accessory nerve was then exposed, its sheath incised at a level convenient for union with the divided facial, and into it the distal segment of the facial nerve was fixed by means of fine silk sutures. After healing of the wound, the muscles on the paralysed side were assiduously stimulated by daily galvanism for months, until faradic excitability reappeared, when faradism was substituted.

In every case where microscopic examination was carried out, new nerve fibres were demonstrated in the distal segment of the paralysed facial. (Compare Figs. 8, 9, and 10.)

HISTORICAL NOTE.

In 1895 the facial nerve was united to the spinal accessory by one of us at St. Thomas's Hospital for the relief of facial palsy due to ear disease. (Case 1.) In 1898 Faure² united the trunk of the facial to the trapezius branch of the spinal accessory, the branch being divided and end-to-end anastomosis being carried out on the posterior belly of the digastric

muscle. The result was not satisfactory. In 1899 Kennedy³ divided the facial nerve for facial spasm, and united it to the spinal accessory (end-to-side junction). The spasm was cured, and subsequently remarkably free movement of the face occurred on movement of the shoulder. In January, 1903 one of us united the facial to the hypoglossal (end-to-side junction), but the result of this operation cannot be determined for some months yet.

CLINICAL CASES.

CASE I.—G. G., a boy 11 years of age, was operated on in 1895 for left-sided otitis media. The left face became paralysed after operation, and facio-accessory anastomosis was performed six months later. The boy disappeared from observation for seven years, but reappeared again in 1902. His condition now is interesting. There is apparent left facial palsy, flaccid in type, with drooping of the angle of the mouth and widening of the palpebral aperture. (See Fig. 1.) There is also an old internuclear strabismus of the left eye, dating from infancy and quite unconnected with his facial palsy. All the muscles of the face react briskly and normally, both to faradism and to galvanism. Taste in the anterior part of the tongue is still absent on the affected side. No voluntary dissociated movement can be carried out in the left face, but on raising the left shoulder against resistance, marked contraction occurs on the left side of the face, and is maintained as long as the trapezius is innervated. (See Fig. 2.) Occasional fibrillary tremors occur in all the muscles of the left face.

CASE II.—Miss Q., 23 years of age, suffered from left-sided otorrhoea for twenty years. In addition, within the last seven years she had also had numerous attacks of left facial palsy. A fortnight after her seventh and last attack of palsy in April, 1901, the ear condition was operated upon, a cholesteatomatous mass being removed from the mastoid. The bony wall of the Fallopian aqueduct was found to be eroded, and the facial nerve lay exposed on an open shelf instead of within a closed canal. The facial muscles on examination were completely paralysed, had lost their excitability to faradic shocks, and the galvanic reactions though present were sluggish and with reversed polar phenomena. Taste was lost in the anterior two-thirds of the tongue on the affected side. In December, 1901, eight months after the onset of the last attack of facial palsy, electrical treatment having meanwhile proved unavailing, facio-accessory anastomosis was performed. Some transient weakness of the left sterno-mastoid and trapezius muscles followed, but these two muscles did not lose their faradic excitability, and rapidly recovered. It was not, however, until October, 1902, nine months after the operation, that any return of motor power was noticeable. At that date, when the left shoulder was elevated against resistance, there was a twitch of the orbicularis palpebrarum and an outward movement of the angle of the mouth. The previous flaccidity of the face was also much diminished, and the deformity of the face very slight when at rest. Taste was still absent in the anterior part of the tongue on the affected side.

CASE III.—The case was one of a married woman, aged 33, who had been under the care of Dr. Wright, of Eye. She was admitted into the Norfolk and Norwich Hospital on August 12th, 1901, suffering from right otitis media, for which, on August 16th, the mastoid operation was performed. Complete right facial paralysis followed the operation with wasting of muscles and loss of faradic contractility. Six months after operation although the constant current had been regularly applied, there was no sign of return of voluntary power. The difference in the two sides of the face when at rest, was, however, not so marked as immediately after the operation, a certain amount of contracture having ensued. At this period it was noted that all the muscles supplied by the facial nerve reacted well to the galvanic current—a very slight response to the faradic current was also noticed in the orbicularis palpebrarum, levator labii superioris, levator labii superioris naeque nasi, orbicularis oris, and depressor anguli oris. On February 23rd, 1902, that is, a little more than six months from the date of the mastoid operation, the patient was placed under chloroform and a faradic current of a strength greater than she could bear on the skin of the face without an anaesthetic was applied and all the muscles responded well. The right spinal accessory and facial nerves were exposed. A faradic current was then applied directly to the trunk of the facial nerve, and all the facial muscles immediately contracted vigorously. The nerve was divided in the canal $\frac{1}{2}$ in. above the stylo-mastoid foramen by chiselling away the overlying bone, and the distal portion, after a length of $\frac{3}{4}$ in. had been removed for microscopic examination (see Fig. 10), was brought down and attached end-to-side by three sutures of 0000 French silk to the spinal accessory trunk. The sheath of the spinal accessory was vertically incised for $\frac{1}{2}$ in. at the spot selected for the union, and it was thought that two of the sutures included a few fibres of the nerve in addition to the cut edge of the sheath. The wound was closed and the patient's head inclined to the right shoulder by means of bandages to relieve the nerve junction of all tension. Three weeks after the operation there was no response to faradism in any of the facial muscles, and the orbicularis oris and levator labii superioris were alone excitable to galvanism, and that to only a very slight degree. A week later, however, all the muscles responded to the constant current. There was considerable wasting of the sterno-mastoid as a result of the second operation, and it was not till a month had elapsed from that date that any voluntary power could be detected in it. The facial paralysis certainly appeared to be more pronounced after the operation than when examined immediately before it, and this was quite noticeable a month later when she left the hospital. Three months after operation a little voluntary power was first detected in the muscles of the right side of the face, more marked in those of the lower than in those of the upper half. This movement was

associated with contraction of the right sterno-mastoid and trapezius. The facial muscles showed no reaction to the faradic current, but responded well to galvanism. Eight months after operation voluntary power had returned more completely, especially in the lower half of the face. She was able by a strong effort to close the eye entirely. The sterno-mastoid had regained a considerable amount of power (Figs. 3, 4, and 5 were taken at this period). Every voluntary movement in the right side of the face, however, was associated with contraction of the sterno-mastoid and trapezius, and conversely, contraction of these muscles was associated with movement of the face. When the patient was working her sewing machine with the right hand, each revolution of the wheel was accompanied by a contraction of the face, there being no power of dissociated movement. She had developed the habit, when wishing to move the right side of the face, of gripping the right hand with the left, thus partly fixing the shoulder and thereby making its movement less noticeable. During emotional excitement, for example, laughing, the left side of the face alone moved, the right being motionless. All the facial muscles reacted to the faradic current, though less briskly than on the healthy side; they also reacted well to the constant current. Ten months and a-half after operation the patient stated that she felt more power in the face, that food no longer lodged in the cheek, and that the movement of the shoulder associated with that of the face and that of the face with the shoulder was much less. The associated movement of the face when turning the handle of the sewing machine had almost disappeared. Upon examination, this diminution in the associated movements was quite evident and yet there was no absolute dissociation of the two. In laughing the left side of the face still moved alone. The facial muscles reacted well to galvanism, the frontalis better than the rest. To faradism the reaction had improved, but the frontalis was not so easily excitable as the other muscles, and voluntary power had not recovered in it to the extent that it had in the rest of the face.

CASE IV.—Miss Z., 36 years of age, suffered from left-sided chronic otitis media, for which an operation was performed in October, 1901. On the second day after operation total left-sided facial palsy appeared, with loss of taste in the anterior two-thirds of the tongue on that side. In spite of electrical treatment the paralysis remained permanent. After four months slight contracture of the paralysed muscles developed, and the electrical reactions were those of profound degeneration. In February, 1902, four and a-half months after the onset of the palsy, facio-accessory anastomosis was performed. Under the anaesthetic no reaction could be obtained in the paralysed muscles to strong faradic shocks applied through the skin, but direct stimulation of the exposed nerve trunk caused brisk contraction of all the paralysed muscles. After the operation the contracture disappeared, the paralysed side of the face again becoming flaccid, and there was superadded a paralysis of the left sterno-mastoid and upper part of the trapezius. These two latter muscles were found, three months after the operation, to have recovered. Four and a-half months after operation, on elevating the left shoulder against resistance, slight movement was observed in the zygomatic muscles of the paralysed side, and this movement gradually increased in strength, until after the lapse of seven months from the operation there was marked elevation of the angle of the mouth (Fig. 7) and of the ala nasi, with some contraction in the orbicularis palpebrarum. The flaccidity of the face also disappeared, so that at rest it was practically symmetrical (Fig. 6). In November, 1902, nine months after the operation, faradic excitability was found to have reappeared in the muscles of the left face, and the galvanic reactions, though sluggish, now showed normal polar differences. Fibrillary tremors were also visible at the angle of the mouth. No voluntary movement could yet be performed without simultaneous elevation of the shoulder—that is, the patient could not dissociate the movements of the face from those of the shoulder. Moreover, the facial movements were so easily elicited by slight shoulder movements that the patient had to carry her parasol or umbrella in the right hand instead of the left, otherwise involuntary facial movements so readily occurred that awkward misunderstandings with strangers resulted. The loss of taste was as before. In January, 1903, eleven months after operation, elevation of the inner end of the eyebrow reappeared.

CASE V.—At the age of 15 years Miss X. fell through the skylight of a conservatory, a distance of over 15 ft., sustaining a fracture of the base of the skull, with right facial palsy and deafness of the right ear. The deafness cleared up in about a year and a-half, but the facial palsy remained complete when she came under our observation nearly three years after the original injury. The right facial muscles were then totally paralysed, flaccid, and exhibited the electrical reactions of degeneration, loss of faradic excitability with sluggish response to galvanism, the anodal contraction at closure being greater than the cathodal. Taste was lost in the anterior two-thirds of the tongue on the affected side. The most assiduous treatment by massage and electrical stimulation ever since the original accident had been carried out but had failed to produce the slightest improvement. Accordingly an operation was recommended and in April, 1902, two years and eleven months after the accident, facio-accessory anastomosis was performed. Whilst the patient was under the anaesthetic, the strongest faradic shocks applied through the skin failed to produce any reaction in the paralysed muscles. But when applied direct to the trunk of the facial nerve after its exposure in the wound, it produced faint contraction in the orbicularis oris and in the lower-lid fibres of the orbicularis palpebrarum. For two months after operation, the sterno-mastoid and upper portion of the trapezius were paralysed, with loss of faradic excitability; they then recovered completely. Careful galvanic treatment was meanwhile applied to the paralysed facial muscles, and in September, 1902, five months after operation, slight movement first appeared in the face when

the right shoulder was elevated against resistance. The earliest facial movement to return was a horizontal outward movement of the angle of the mouth. A few weeks later, on innervation of the trapezius as before, the eye could also be partially closed and the angle of the mouth was drawn outwards and upwards. The loss of taste remained as before. Microscopic examination demonstrated the presence of young nerve fibres in the distal segment. (See Figs. 8 and 9.)

CASE VI.—M. K., a girl, aged 18 years, had chronic middle-ear disease, for which she was operated on in February, 1902. Ever since, the right face had been totally paralysed, and in August, 1902, six months after the onset of the palsy, no improvement having occurred meanwhile, one of us was requested to admit her to St. Thomas's Hospital in order that facio-accessory anastomosis should be carried out. The electrical reactions were those of profound degeneration. When last seen in January, 1903, four months and a-half after the operation, no recovery had yet taken place, the face was still flaccid and paralysed, and the electrical reactions as before. Galvanic treatment is still being carried out.

CASE VII.—Facio-hypoglossal anastomosis:—Mrs. B., aged 44, had suffered from right facial palsy for twelve months. On admission to the National Hospital for the Paralysed, Queen Square, on December 10th, 1902, there was complete right peripheral palsy. No contraction to faradism was obtained in the face muscles. There was also anaesthesia in the distribution of the inferior dental nerve, and loss of taste in the anterior two-thirds of tongue. The condition was due to a tumour of the right ramus of the lower jaw. On January 9th, 1903, the right ramus of the lower jaw was removed, together with a cystic tumour. The hypoglossal nerve was defined, the occipital artery and descendens nerve being divided so as to free the hypoglossal. The facial nerve was divided at the stylo-mastoid foramen and sutured (end-to-side junction) to the hypoglossal. On February 2nd there was considerable atrophy of the right side of the tongue, which was protruded towards the paralysed side. The facial muscles are being treated by galvanism. As yet, there is, of course, no recovery in them.

MODES OF OPERATION.

In Case 1 (1895) the spinal accessory was half cut through, and in the niche thus formed the trunk of the divided facial was fixed by fine silk sutures.

Faure's (1898) first method was the end-to-end suture of the divided trapezius branch of the spinal accessory with the end of the divided facial. The union was effected on the surface of the posterior belly of the digastric. Faure's second method consisted of the division of the trapezius branch of the spinal accessory. This was then brought upwards outside the belly of the digastric and fixed end-to-side to the uncut trunk of the facial.

Kennedy's method (1899) consisted in division of the spinal accessory, with the exception on one side of its perineurium, and the suturing, end-to-side, of the divided trunk of the facial into this gap.

Fig. 11.—Facio-spinal anastomosis in the dog. Thirteen months after operation. F, facial nerve. SA¹, spinal accessory (central end). SA², spinal accessory (peripheral end). L, silk ligature. An, point of anastomosis. After Manasse. (*Langenbeck's Archiv*, 1900. Bd. lxi.)

Manasse⁴ (1900) recommended the division of the facial nerve at the stylo-mastoid foramen and the suturing of the cut trunk, end-to-side, to the spinal accessory (Fig. 11). This operation required, of course, the division of the posterior belly of the digastric, and it is practically the one which we now recommend with very slight modification (see Fig. 12).

Barrago-Ciarella⁵ (1901) performed some interesting experiments on dogs. He found that recovery of movement in the face after facio-accessory anastomosis commenced in four and a-half months and was complete in six months, and that

Fig. 3.—Case III at rest.

Fig. 4.—Case III; during elevation of shoulder.

Fig. 5.—Case III; during elevation of shoulder.

Fig. 1.—Case I; at rest

Fig. 2.—Case I; during elevation of left shoulder.

Fig. 6.—Case IV; at rest.

Fig. 7.—Case IV; during elevation of shoulder.

Fig. 8.—Facial palsy of three years' duration. Case v; longitudinal section of distal segment: Van Gieson stain. ($\times 200$.) Shows proliferation of neurilemma cells and their arrangement in imbricating rows.

Fig. 9.—Facial palsy of three years' duration. Case v; longitudinal section of distal segment. Weigert stain. ($\times 300$.) Shows young nerve fibres in the distal segment.

Fig. 10.—Facial palsy of six months' duration. Case III; longitudinal section of distal segment. Kulschitzky-Wolters method. ($\times 300$.) Shows young nerve fibres in the distal segment.

recovery after facio-vagal anastomosis commenced in six months and was complete in eight. Our cases show that in man recovery after facio-accessory anastomosis takes about the same time as in the dog. (In Case III recovery commenced three months after operation.)

Fig. 12.—To illustrate the method of end-to-side anastomosis, recommended for facio-accessory or facio-hypoglossal anastomosis. The spinal accessory or hypoglossal nerve (SA) is shown with a longitudinal incision in its sheath which gapes, and thus exposes the nerve fibres. A short longitudinal incision in the sheath of the facial (F) is also made so as to expose freely the nerve fibres, which retract somewhat into the sheath when the trunk of this nerve is severed. The figure to the left shows the fine silk threads passing through the sheath only of each nerve, that to the right shows the sutures drawn tight and tied.

From the above series of facio-accessory cases it will be observed that in five patients out of six voluntary movement has already reappeared in the paralysed facial muscles, the nerve prior to operation having been paralysed for periods varying from five months to three years. In the sixth case, only four and a half months have elapsed since operation, and we therefore anticipate that recovery will yet occur. (In Case II the first sign of returning movement did not appear until nine months after operation.)

One of the immediate results of operation is invariably to produce a temporary palsy of the sterno-mastoid and upper part of the trapezius, due to affection of the spinal accessory. This clears up completely in all cases. Another point of interest is that if the paralysed muscles have already developed contracture, to however slight a degree, they at once become flaccid again; in fact, in several cases where clinically the muscles before operation appeared to be flaccid, this flaccidity seemed to become actually exaggerated, a condition for which we do not offer any explanation, except that possibly in such cases the nerves, though completely disconnected from the brain, do in some way, by virtue of the regeneration that has taken place in them, exert an influence on the muscles.

An observation to which it is desirable to draw attention is that the facial muscles during operation (see Cases v and vi), in which the patient was completely anaesthetized, though unresponsive to the strongest faradic shocks applied through the skin, nevertheless reacted when the peripheral end of the divided facial nerve was stimulated directly. This is possibly a clinical evidence of peripheral regeneration.

After operation, motor power returns in the affected muscles before faradic excitability; thus in Case IV, where motor power returned at four and a-half months, faradic excitability did not reappear until the ninth month.

Contemporaneously with the recovery of motor power, the muscle tonus in a certain number of cases also returns, so that at rest the face becomes again symmetrical, instead of distorted as before. This is well seen in Figs. 3 and 6. Such recovery from disfigurement is in itself no small benefit, especially in the case of lady patients. We must admit, however, that the flaccidity does not always disappear, even where motor power returns, as in Case I. (See Fig. 1.)

So far, in our series of facio-accessory cases we have not observed any independent movement of the face, unassociated with that of the trapezius and sterno-mastoid, although in several cases a minimal innervation of the trapezius is sufficient to cause facial contraction. The accomplishment of dissociated facial movement is, of course, a matter of education of the cortex, or of that part of it from which shoulder movements are initiated. Such a patient will require to educate afresh part of the shoulder centre at the junction of the upper and middle thirds of the ascending frontal gyrus,

to do the work of the old facial centre some distance away, in the lower third of the same gyrus. (See Figs. 13 and 14.)

The left hemisphere of brain of Chimpanzee

Fig. 13.—Sherrington and Grünbaum's motor areas. (Note relative positions of face, tongue, and shoulder centres.)

Bearing this in mind, we considered whether it might not be practicable to select for anastomosis with the distal segment of the paralysed facial some other healthy nerve supplying muscles whose cortical centre is nearer to the face centre than in the case of the shoulder. The hypoglossal nerve and the cortical tongue centre fulfil these requirements.

As will be seen from the accompanying diagrams (Figs. 13 and 14) the face centre and the tongue-centre overlap one another, and we would therefore anticipate that the prospects of dissociated facial movement through the tongue-centre would be greater than when working through the shoulder-centre. Moreover it may be recalled that the movements of the tongue are closely associated with those of the lips, so much so that in health the orbicularis oris and the transverse fibres of the tongue always contract together. It is more than probable that the facial nerve derives its lip fibres from the hypoglossal nucleus. There is, therefore, a close relation, not only between the cortical face-centres and tongue-centres, but also between the facial and hypoglossal nuclei in the bulb.

Accordingly, in a case of facial palsy due to tumour in the parotid region, one of us, after removal of the tumour, united the distal segment of the facial to the healthy hypoglossal (Case VII). And this latter operation of facio-hypoglossal anastomosis is the one which we would recommend in future in preference to that of facio-accessory union.

In a letter to one of us in criticism of Case III, Professor E. A. Schäfer states that, in his opinion, although the cortical face centre had been cut off from its peripheral connexions for a period of six months, yet it would not have lost its power to start impulses, and since the movements of the two sides of the face are commonly associated, he would expect one facial centre to retain sufficient functional activity for both.

Professor Schäfer also suggests the glosso-pharyngeal as a suitable nerve to which to attach the divided facial, on the ground that its motor nucleus is much nearer to and is apparently serially homologous with the facial nucleus. With reference to this we think that the small size of the glosso-pharyngeal and its relative inaccessibility would make the operation one of extreme difficulty.

If the cortical facial centre still retained the power of generating impulses six months after division of the facial trunk, then the absence of the recovery of voluntary dissociated movement might be due to the inability of the "association fibres" running from the centre of the face to that of the shoulder to transmit such impulses.

Note.

CASES I, II, IV, V, VI, and VII were operated on by C. A. Ballance, who has also operated on another case of facial palsy, but the operation could not be completed, as the trunk of the facial nerve had been carried away by a bullet.

CASE VII is of special interest, as in this the trunk of the facial nerve was united to the hypoglossal. If in this case dissociated movement of the face occurs it will open the question as to whether the previous cases of facio-accessory anastomosis might not with advantage be re-operated on.

Fig. 14.—Schema to illustrate the cortical connexions between the movements of the face, tongue, and shoulder respectively, and the relations of the corresponding nerves in the medulla. It will be seen that the centres of the movements of the tongue and the face overlap in the cortex, whilst that for the shoulder is some distance away from the face. It will also be observed that the facial nerve in the medulla derives some of its fibres from the hypoglossal nucleus.

CASE III was operated on at Norwich by H. A. Ballance. Purves Stewart independently examined Cases I, II, IV, V, and VI, and directed their electrical treatment. He also prepared the microscopical preparations (see Figs. 8, 9, 10), the histological portion of the work being carried out at the neuro-pathological laboratories of King's College, by kind permission of Dr. Ferrier.

CONCLUSIONS.

1. Peripheral facial palsy is remediable by facio-accessory anastomosis, but the extent of recovery appears to be limited

to associated movements in conjunction with the shoulder. In most cases the previous deformity disappears when the face is at rest.

2. For reasons above stated we would in future recommend facio-hypoglossal anastomosis rather than facio-accessory.

3. The cases suitable for operation are those in which the paralysis has lasted so long that no recovery is to be expected, say, facial palsy lasting six months without any sign of

recovery. In our opinion the sooner the operation is done after this date the better.

4. A suppurative causal condition producing an infective neuritis renders the prognosis after operative treatment less favourable than in cases due to trauma.

REFERENCES.

¹ *The Healing of Nerves*. By C. A. Ballance and Purves Stewart. 1901. Macmillan and Co. ² *Traitement Chirurgial de la Paralyse Faciale par Anastomose Spino-faciale*. Congrès de Chirurgie. 1898. ³ *Phil. Trans. Roy. Soc. Lond.* Series B, vol. cxciv. 1901. ⁴ *Langenbeck's Archiv* 1900. Bd. lxxii, fasc. 4. ⁵ *Il Policlinico*, Rome, February, 1901, fasc. 3, p. 21 (ref. by Bréavoiné in *Chippault's Travaux de Neurol.-Chirurg.*, 1901, p. 92).

CASE OF BRAIN TUMOUR.

By A. C. DUTT, B.A., M.B.CANTAB.,

With Remarks by D. Lowson, M.D.,

Surgeon, Hull Royal Infirmary.

ON Sunday, October 19th, 1902, I was sent for in the evening to see S. P., aged 39, who, I was told, was in a "fit." On my arrival I found him lying on the sofa, unconscious, cyanosed, and breathing stertorously. His friends thought it was a "bilious attack," to which he was subject at intervals.

History.—Eleven years ago he had had a similar attack, with violent headache and delirium, when he kept to his bed for a fortnight. As a child he had had "rheumatism" very badly. Six years ago he had pleurisy on the right side. Every few days he had headaches; when an attack came on out-of-doors his wife said "he crawled along the road as if he could not see." He had been getting thinner. Last winter in Belgium he had a severe headache, when he walked about in his bedroom for three hours. His mother died when he was 10 years old; cause of death unknown. Father died of pneumonia. Of his two brothers, one died of phthisis. His two sisters are alive and well. The patient was a traveller for his brother, a fruit merchant in this city, and had been in America several years, where he had had a rough and arduous life. He had great business capabilities, and was the trusted adviser of his brother, who valued his services highly. He has several young children, quite healthy, and of normal development. His wife said he was very irritable at times, but this was attributed to a "sluggish liver."

State on Examination.—The patient was of medium height, spare build, and very dark for an Englishman. The attack was undoubtedly epileptiform. He was ordered to be put to bed, and a mixture of bromides with belladonna was prescribed—with 3 gr. calomel at bedtime. In the morning he was considerably better, and conversed rationally and intelligently. Some years ago, he used to take alcohol freely, and at the age of 19 had contracted syphilis in America for which he had been treated for two months or so. There was no history of any other serious illness, except a slight attack of ague abroad, and several influenzal seizures at home. As he complained of indigestion and sickness a mixture of bismuth and salol with pepsin was ordered. The improvement being maintained, on October 22nd I gave nitromuriatic acid with strychnine before, and pepsin pills after meals. On the 23rd there was a good deal of flatulence necessitating the use of sodium bicarbonate and Bragg's charcoal which gave temporary relief. On the 24th violent headache was complained of, and the dyspeptic symptoms were worse. Phenacetin and caffeine relieved the headache; $\frac{1}{2}$ m creosote pills relieved the flatulence better than the charcoal powder. From this time the patient could not do without the phenacetin powders. Mercurial ointment was ordered to be rubbed into the axilla for three or four nights and a 5 per cent. ointment of oleate of mercury was applied to the scalp. This quickly produced stomatitis and had to be abandoned. On October 31st 3 ss. of pot. brom., 10 gr. chloral, and 10 m tr. hyoscyamus was given for insomnia, but it did not produce the desired effect. The draught was repeated and 12 gr. of sulphonal given—producing two hours' restless sleep. On November 3rd quinine and phosphoric acid was tried, but this had no effect on the "neuralgia" (as the patient called it) and made him worse. All this time there was no tenderness of the scalp anywhere; mercurial inunction was begun again, but the next day the mouth was as bad as ever. On the 4th blisters were put behind the ears, which seemed to do a little good. The patient now remarked he could only see one of my eyes and half my face, and that the ornaments on the mantelshelf seemed double. Every remedy having so far

failed to ameliorate the symptoms, and mercurial inunction being impracticable, I thought of the possibility of a syphilitic growth in the brain, and on November 5th prescribed potassium iodide beginning with 5 gr. t.i.d. The temperature so far had been subnormal, and the pulse 60 to 68; respirations 16 to 20. The iodide was increased by 2 gr. t.i.d. every second day till he took 20 gr. t.i.d. without apparent discomfort. There was now homonymous hemianopsia, defect of colour vision, and alexia or word blindness.

On November 8th Dr. Lowson saw the patient with me, took great interest in the case, and agreed with me in my diagnosis of an intracranial growth. On November 18th the teeth were tender and the mouth inflamed. Washing the mouth and brushing the teeth with "listerine" greatly improved this condition. Dr. Daly saw the case with me on November 27th and during my absence on the two following days; his opinion coincided with ours. In addition to alexia, the patient was apraxic—that is, forgot the use and significance of common objects, for example, knife and fork. Both alexia and apraxia pointing to a lesion in the angular gyrus extending back into the occipital convolution, we came to the conclusion it was probably a gumma. The patient could not call to mind objects named and point them out at request. He could recognize letters, but could not form them into words—for example, he could name the letters in *Yorkshire Post*, but could not say the name of the paper; he would call red and green blue. He could understand spoken words, but not words written or printed. He could not read, and could not recall objects when the names were shown to him. He could write voluntarily and at dictation fairly well, but could not copy, there being some paraphasia, "ryadd" being written for "Sunday"; there was no power to write the names of objects seen, heard, felt, tasted, or smelt; knife, spoon, fork, not being recognized. There was no power to read aloud or inaudibly, but he could write his name well and read it. He could also read my signature. He could not recognize a letter when traced with a pencil on the palm of his hand. The power to call up mentally the appearance of an object was absent, but he could read figures, calculate, and count both money and in numbers. There was papillitis more marked in the left optic disc. Occasional twitching of the left hand, and frequent retching were complained of, but never any sickness, the patient eating two or three good meals a day. Sometimes he was troubled with irritation in the left nostril and deafness in the left ear. There was some paraphasia, persistent insomnia, and occasional delirium. The headache was worse; he would knock his head on the wall to get relief.

On the December 20th the temperature was 100°, and stiffness of the neck was superadded to the other symptoms. Localized sweating of the scalp was noticed on December 26th. On December 27th the temperature was 102°, and Drs. Lowson and Daly saw the patient with me again. On the 28th he complained of difficulty and pain in passing water, a few drops being passed frequently.

A prolonged course of potassium iodide having failed to do any good, and the prognosis being desperate, I pointed out to the patient's friends that the case was hopeless, the only prospect—and that a very slender one—of alleviation being an operation. Their consent having been obtained, Dr. Lowson operated on December 31st, chloroform being the anaesthetic chosen. A cap of lint, soaked in 1 in 20 carbolic had been worn over the shaved scalp twelve hours before, and a quarter of a grain of morphine was injected hypodermically an hour before the operation. According to Treves the morphine has the effect of contracting the arterioles of the brain, and hence of lessening haemorrhage; it also allows less ChCl, to be given.¹

Operation.—A single flap was raised at the region of the left angular gyrus perpendicular to Reid's base line. The outline at the flap formed a shallow curve, being free below. The pericranium being separated by the rugin over the area exposed, and all bleeding arrested by pressure forceps (there being some difficulty owing to the thickness of the scalp), the trephine was applied and a disc of bone removed. Immediately the dura bulged out, due to great intracranial pressure. But on incising the dura no growth was seen. Dr. Lowson put his little finger into the wound, and felt some adhesions about $1\frac{1}{2}$ in. to 2 in. posterior to the opening made. The hole was enlarged by chipping off the bone all round, and the dura opened at this spot. On passing a grooved director, much pus-like material oozed out. This was subsequently kindly examined by Dr. Daly, who pronounced it to be most probably the debris from a breaking-down gummatous growth.