staining reaction of the bodies, which I can confirm from my own experience. In using the methylene blue and eosin stain of Romanowsky the nuclear point of the cell-inclusion easily stains red, whereas with the ordinary nuclear dyes it is stained only with great difficulty. The zone immediately around this point remains unstained, and the protoplasm stains a faint blue and the capsule a darker blue. This reaction of the bodies to this particular stain is precisely the same as that of the unicellular animal organisms—for example, amoebae, the malarial parasite, the trypanosomata, etc.—and in all these instances there is also the unstained zone immediately around the nuclear point.

There has not been sufficient time as yet to confirm or disprove the part of his book referring to the development, multiplication, etc., of the bodies.

So far, then, has this particular part of the cancer question reached. If we consider the points I have brought forward above, it will be, I think, impossible for us to shut out the parasitic theory, for not only does the origin of cancer, but also its growth and the differentiation of its cells, find complete explanation in this theory.

Moreover, it seems to me, if we think of the clinical course of the disease, its beginning in one spot, its extension to distant parts by lymphatic or blood ways, the cachexia out of all proportion to the extent of the disease, the spread by contagion, the occurrence in certain parts of the body, and its return after years of quiescence, we are driven, from this side, too, on to the parasitic theory, in which (as in no other) all these events find their explanation.

Lastly, the only hopeful outlook in cancer would seem to rest, too, on the ground of the parasitic theory, for was it not Pasteur who said that the mind of man shall become lord over all infectious and parasitic diseases?

BIBLIOGRAPHY.

I have thought it better not to encumber the above with references, but the following works have been especially mentioned or consulted: Behla, *Die geographisch-statistische Methode als Hilfsmittel der Krebsforschung*, 1899; *Die Krebserkrankungen der Stadt Luckau von 1878 bis 1899: Ueber "Cancer à Deux" und Infection des Krebses*. Borrel, *Annales de l'Institut Pasteur*, 1901 and 1903. Czerny, *Beiträge zur klinischen Chirurgie*, Bd. xxv, Heft 1. Feinberg, *Das Gewebe und die Ursache der Krebsgeschwülste*, 1903. Gaylord, *Reports of the New York State Pathological Laboratory*, Buffalo; *Orth-Festschrift*. Guelliot, *Gazette des Hôpitaux*, Paris, 1892. Hanau, *Fortschritte der Medizin*, 1890. Jensen, *Centralblatt für Bakteriologie*, I, Bd. xxxiv, 1903. Lubarsch, *Pathologische Anatomie und Krebsforschung*, 1902. Morau, *Archives de Médecine Expérimentale*, 1894. Nawaschin, *Flora*, 1895. Nösske, *Deutsche Zeitschrift für Chirurgie*, Bd. lxi, Pianese, *Zeitschrift für Hygiene*, Bd. xxxvi; *Histologie und Ätiologie des Carcinoms*, 1886. Podwyszozi, *Centralblatt für Bakteriologie*, Bd. xxvii. Rüffer, *Sur les Parasites des Tumeurs Epithéliales Malignes*, 1896. Schueller, *Die Parasiten im Krebs und Sarkom*, 1901. Von Hanseemann, *Die mikroskopische Diagnose der bösartigen Geschwülste*, 1897; *Studien über die Specificität, den Altruismus, und die Anaplasia der Zellen, mit besonderer Berücksichtigung der Geschwülste*, 1893. Von Leyden, *Ueber die Parasiten des Krebses*, 1902. *Verhandlungen des Comité für Krebsforschung*, Hefte 1 und 2. *Veröffentlichungen des Komitees für Krebsforschung*, 1902.

THE BEHAVIOUR OF CANCER WITHIN NERVE AND TROPHIC AREAS.*

[WITH SPECIAL PLATE.]

By G. LENTHAL CHEATLE, C.B., F.R.C.S.,

Surgeon, King's College Hospital; Italian Hospital; and Hospital for Epilepsy and Paralysis, Maida Vale.

It would be advisable for those who read this paper to separate the consideration of cancer into three parts. I regard these three headings as essential to the study of carcinoma.

1. The genesis, which includes those matters which appertain to the actual agent which induces epithelial proliferation.

2. The incidence, those matters which relate to the site or soil in which cancer primarily begins; and

3. The spread, that which concerns the area of occupation of the cancer when considered apart from its secondary deposits.

There can be no doubt that the genesis and incidence must be closely related, but the fundamental influence at work in each may be quite different. I do not intend to enter into the genesis of carcinoma under a separate heading in this paper, but will refer to it during the observations on the incidence and spread of cancer.

The Incidence of Cancer.

In my last paper¹ I pointed out that there were grounds for considering the incidence of cancer had possibly a direct or

indirect connexion with the nerve area on which it fell. My reason for so doing was, among others, that it is impossible to omit the subject of irritation from among the etiological factors of cancer. Cancer, and especially squamous epithelioma, more often than not begins upon the actual site of irritation.² As constant irritation applied over a long period produces marked intracellular changes in the ganglia of the posterior sensory roots, it will also probably induce profound physiological changes in the areas of their distributions as well as at the actual sites of irritation.

Again, the incidence of squamous epithelioma and rodent ulcer are very closely associated with the distribution of the fifth cranial nerve. When rodent ulcer is multiple, the points of incidence are nearly always on the area or areas of one or both fifth cranial nerves respectively, and chiefly on those parts which are not overlapped by the second cervical spinal nerves (see Fig. 1). I also pointed out that squamous epithelioma and rodent ulcer often appeared on those points at which nerves become cutaneous. Dr. Head has described points in the peripheral distribution of the posterior spinal ganglia, which he terms maximum points.³ The maximum

Fig. 1c.—Dr. Colcott Fox's case of multiple rodent ulcers on one fifth cranial nerve. The two upper ones have united to form a single ulcer.

points are those foci within the areas of referred pain in visceral disease at which the pain is most acute. Herpes zoster, when its distribution is complete, maps out the whole area of referred pain, but when incomplete generally appears on the maximum points. These maximum points somewhat closely correspond to the situations at which nerves become cutaneous. Whether maximum points are necessarily connected with such sites I do not know. I understand that Dr. Head's maximum points are the maximum points of central influence, but I do not know whether even a peripheral nerve can or cannot exhibit or claim its maximal points of central influence. To-day I attempt to show that the incidence of cancer frequently falls on these maximum points, described by Dr. Head. I have Dr. Head's kind permission to reproduce three plates (Figs. 2, 3, and 4) from the articles to which I have already directed attention. By comparing the rodent ulcers in Figs. 1, 5, 6, 7, 8, 9, 10 with the maximum points in Dr. Head's plate (Fig. 4), my meaning will be clear. Mr. Harold Barnard has collected ten cases of tar cancer in which only one lesion occurred in each case. Fig. 11, Diagrams A, B, C, show the distribution of Mr. Barnard's cases. Diagram D is one of my own. These cases show points of incidence.

The rodent ulcers are only examples of my meaning; but

* The substance of two papers recently read before the Pathological Society of London.

Fig. 2.—The maximum points are shown on the left side.

the tar cancers represent the distribution of all the tar cancers on the face which I can now produce.

The number of rodent ulcers which appear on the face are so numerous that one requires a great many more cases than I have been able to collect to show fairly the proportion of cases arising on maximum points to those which do not. However, those cases in which a rodent ulcer appears on the trunk are uncommon, therefore it may be reasonable to compare the one with the other. The maximum points, too, do not correspond so closely with what the limit of our knowledge at present recognizes as embryonic clefts. And the points on the trunk have an extra advantage of appearing on a bigger field, which makes the proportion more valuable as evidence. I have collected 10 cases of rodent ulcer on the trunk, not with the special object of collecting rodent ulcers on the body, but simply taken out of my collection. Out of 10 separate cases 7 began exactly on maximum points. In 1 of the remaining 3 the ulcer was extensive, and its history does not record where it began. Another of the cases which is corresponding to part of the distribution of the twelfth dorsal could not have begun on the maximum point; and in the drawing of the remain-

Fig. 3.—The maximum points are on the left of the figure.

Fig. 4.

ing case so many lesions are depicted it is difficult to say which are rodent ulcers and which are not. This proportion may be accidental, but the fact justifies attention being drawn to it.

With regard to the tar cancers (Fig. 11) there are 11 cases; 4 began on the centre of the maximum point; 2 began at maximum points, but towards their margins. Of the 11 cases 8 are in relation to clefts; I do not see how tar

cancers can be etiologically related to clefts. With regard to clefts it is interesting to point out that the incidence of cancer is generally at the same point in the same cleft region, as if some other influence than the mere presence of a cleft was an agent involved in the process.

Rodent ulcer more frequently begins on the lowest part of the fronto-nasal maximum point where the infra-trochlear

nerve becomes cutaneous than on any other part of the body. The temporal, fronto-temporal, fronto-nasal, maxillary, and mandibular are maximum points on which cancer of the face frequently falls.

Fig. 5.—Dr. Sequeira's case of rodent ulcer on chest (5th dorsal).

It must be recognized in considering this question of possible relationship between maximum points and the incidence of cancer that the maximum point of an area does not claim to possess qualities and properties which do not exist elsewhere on that area, but only to possess them in a more accentuated degree. It is possible to imagine that constant peripheral irritation over a long period would produce profound physiological changes in a nerve area; and in cancer, as well, one sees marked intracellular changes in posterior spinal root ganglia.

With these ideas in mind, one can understand that providing the irritation includes a maximum point and part of the same nerve area, the more likely place for the incidence of cancer would be on the maximum point, that is to say if cancer has any connexion with peripheral irritation and direct or indirect nerve influence. (See Fig. 30, special plate, which is the reproduction of a drawing made from the right forearm of a man suffering from tar cancer and under the care of Mr. F. S. Eve, who kindly gives me leave to publish the case here.) The

Fig. 6.—Mr. Reid's case of rodent ulcer on back of shoulder (2nd dorsal).

Fig. 7.—Mr. J. Hutchinson's case of rodent ulcer of front of axilla (2nd dorsal).

patient was a tar worker, his whole hand and forearm were exposed to and suffered from the effects of the tar irritation, but the place at which the squamous epithelium became cancer was on the maximum point of the first dorsal root. See Dr. Head's diagram in Fig. 2. If, however, the maximum point be not included in the actual focus of peripheral irritation, then the irritation ought to be greater or of longer duration to induce the same effect. At present I cannot see any evidence which leads me to suppose that

Fig. 8.—Mr. J. Hutchinson's case of rodent ulcer (maxillary).

the incidence of cancer in the referred areas can be related to visceral lesions. If one is careful to recognize the central cause of herpes zoster it is a very interesting disease with which to compare cancer from a genetical point of view. Because in herpes zoster, for some reason, either from the nerve stimulus, vasomotor, toxic, or other causes, the epithelium dies. Whatever is the cause of death, it is

Fig. 9.—Dr. Sequeira's case of rodent ulcer (fronto-temporal).

inseparably connected with changes in the posterior spinal root ganglia. One could understand that part of, or modifications of, the same process which induces death of the epithelium may also lead to proliferation of epithelium, and hence the possibility of a direct or indirect nerve agency being

Fig. 10.—Dr. Sequeira's case of rodent ulcer (temporal).

concerned in the genesis of cancer ought to be borne in mind. Again, if I can show that, when spreading, the growth of cancer stops at lines of normal and abnormal nerve limitations, whether natural or artificial nerve boundaries (vide infra), and often erode denser tissues as bone and fasciae after they have reached those lines, then the argument is justified that it may possibly be a direct or indirect nerve influence, and probably not a mechanical or purely uncontrolled vascular influence, which is at work limiting spread.

Fig. 11 A.

Fig. 11 B.

Fig. 11 C.

Fig. 11 D.

Now if such a nerve influence may limit spread, and as it were shut the gates, even if in some cases it appeared to be temporarily, then one is justified in arguing, when considering the question of incidence, that it may allow the area to be opened, or

Fig. 12.—Dr. Sequeira's case of rodent ulcer.

when considering the question of genesis may open the area. The fact that the glans penis, cervix of uterus, and alimentary mucous membranes are common sites for carcinoma indicates that merely tactile sensation apart from pain sensation may not have so much to do with the matter as vasomotor or other impulses of which we are at present unaware. But, whatever the explanation, the fact appears to me to be that maximum points described by Dr. Head are points too frequently associated with the incidence of cancer to be regarded only

as belonging to the dictates of accident.

Spread of Cancer.

The other point I ventured to suggest was that a possible direct or indirect nervous influence was the chief agent in directing the areas occupied by cancer. I now more firmly believe in the possibility of a direct or indirect nervous agent, above all others, influencing the spread, a belief which is constantly receiving support by the addition of fresh cases and continued investigation. But one cannot look at Figs. 12, 13, 14, 15, 16, 17, 18, 19, and compare them with Fig. 20, without being at once struck with the similarity between the arrangement of cutaneous muscles and the distribution of these ulcers. For example, Fig. 13 (special plate) seems to exactly correspond to the area of the levator labii superioris alaeque nasi muscle, and Fig. 16 shows a distribution which can be explained from a possible nerve point of view by saying

the ulcer occupies an area formed by the first and second division of the fifth cranial nerve, whilst the third division is quite free; even that part escapes which is supplied by the buccal branch at the angle of the mouth. But it can also be explained more or less from a cutaneous muscular limitation point of view by saying that the ulcer covers the anterior belly of the occipito-frontalis, the orbicularis palpebrarum, the zygomaticus minor, and part of the orbicularis oris muscles. Where the growth seems to have transgressed these muscular borders in Fig. 16, it may be explained by assuming that the shapes and directions of the contours of the muscles are maintained by a regular excentric growth from the margins at which arrest had previously occurred. It is interesting to observe here that if one closely watches the facial muscle of a clean shaved thin man in the act of public speaking, one can often detect a demarcation between that part of the orbicularis oris at the angle of the mouth and the point which may be the outer border of the levator labii superioris or zygomaticus minor muscle. (See Fig. 15, special plate.) I think these two cases alone will save me from being accused of paying too much attention to this point. Before publishing my last paper I fully considered it among other pos-

Fig. 16.—Mr. Malcolm Morris's case of rodent ulcer.

sibilities as chief limiting factors of spread. I can see that it will be a point raised against the possibility of a nerve influence, and I can see that muscles by their fasciae, intrinsic lymphatics, and vascular arrangements, and also that other vascular arrangements, hair follicles, and cutaneous glands outside the sphere of muscles, do exert a subsidiary action, but only within certain specified lines which I have all along ventured to believe marked the normal or abnormal limits of nerve areas. I will try to point out that this is so.

First of all I will call attention to the intimate anatomical relationship which exists between the cutaneous muscles of the head and neck and the skin. These cutaneous muscles lie ensheathed in comparatively great masses under the cutis vera. They break up into small bundles which can be readily detected, occupying an area within the cutis vera at a level of the sebaceous and sweat glands and hair follicles. With more careful investigation still smaller and more isolated fibres and bundles of fibres can be seen here and there almost reaching the ridges of the squamous epithelium. This arrangement of the facial muscles is most intimate, so far as I can judge, in the orbicularis palpebrarum and the orbicularis oris muscles. Even at their actual beginning it is almost impossible for

rodent ulcer and squamous epithelioma to be too superficial to be in direct anatomical relations to the cutaneous muscles of the region.

Now take the microscopical appearances of the rodent ulcers, at first they spread superficially on the top of the fasciae over the deepest and largest of the three masses of muscles and among the two uppermost sets which I have just described, and completely destroy these and all structures lying above that fascia; after a time they penetrate it and the subjacent muscles, periosteum, and bone. The edges of rodent ulcer for the purpose of this paper may be divided into two—the spreading margin and the margin of arrest. Most edges have much in common; they are all seen situated between the fascial covering of muscle (where there is muscle below) and the atrophying squamous epithelium above. The tumour is closely in contact with both. Hair follicles, sebaceous and sweat glands can be seen being, as it were, pushed before the growth where it impinges upon them, and, indeed, from their assumed shapes, they appear to be offering obstruction to the oncoming tumour.

The most marked factor of mechanical limitation in the cutis vera is probably fibrous tissue. Can the sheaths of the scattered muscular bundles, either collectively or individually, be the means by which growth is arrested? Clinical

Fig. 17.—Mr. Henry Morris's case of rodent ulcer.

evidence is here much more valuable than microscopical evidence, for the appearances at the edge of a tumour not rapidly spreading are relatively about the same as those at the edge of a growth arrested by encountering strands of fibrous tissue. This is so because around the edges of a tumour or that part of a tumour which is not rapidly spreading, the fibrous tissue may proliferate to such an extent as to resemble the edge of a growth mechanically arrested by fibrous tissue.

Take Fig. 13 and realize the clinical fact that a rodent ulcer, whether covering a muscle or fascia, can be penetrating deeply into bone and yet remain limited superficially. This is not common, but I have seen a case. A barrier formed by a cutaneous structure would surely give way before bone or even temporal fascia ulcerated, were it the predominating cause of limitation.

I can quite understand and appreciate the argument that in a large ulcer as represented in Figs. 14 and 15 (special plate), in which the lateral views of the same case are reproduced, the ulceration of bone is most likely to occur in those parts on which the ulcer has existed longest. The rodent ulcer in this case began on the forehead. From this point in ten years it extended all over the face, ulcerated through fasciae, muscles, and bones, too far away (double the distance) from the point

of incidence to permit the argument that mechanical limitation stopped the further ulceration on the scalp. It may be said that it was attracted by the greater vascularity in the direction of spread. But I shall presently show that when a rodent ulcer destroys the nerve supply of the skin and renders it completely anaesthetic and analgeic the growth does not spread into that senseless area. By looking at the devastation wrought by this rodent ulcer in Figs. 14 and 15, no doubt can exist as to the certainty of destruction of all sensory nerves going to the scalp from the orbit. I believe this is the reason of the want of spread further on the scalp in this case.

Fig. 17 is the drawing of a case under the care of Mr. Henry Morris, which he kindly allowed me to photograph and reproduce. It is a rodent ulcer, which from a cutaneous-muscle point of view may be said to cover part of the orbicularis palpebrarum muscle, and be guided towards the ear by the attrahens aurem muscle (the exact point of posterior limitation is not accurately depicted). Fig. 18 represents the same case six or seven years after, and shows that the ulcer has in the meanwhile destroyed the larger amount of one-fifth cranial nerve area, bones included, and yet the posterior

Fig. 18.—Mr. Henry Morris's case of rodent ulcer.

limit cannot justify the assumption that muscle, fascia, and greater advantage of blood supply form adequate reasons for the want of spread posteriorly on the ear and scalp after six or seven years' ulceration of bone anteriorly. It is a unilateral fifth cranial nerve lesion to which further reference will be made when discussing the effect of destruction of the nerves on the spread of the disease.

Fig. 19 shows a case in which the rodent ulcer began on the side of the head just above the ear, in the region of the attollens aurem and attrahens aurem muscles and temporal artery. Muscles and fascial bands connected with them cannot be held to have arrested growth at the posterior margin when the zygomatic process, the temporal fascia, have been eroded and the facial nerve destroyed by the same ulcer further away from the starting point anteriorly. The spread here was not merely circumferential, for when it reached the posterior margin it stopped, while spread continued anteriorly. Eight months after this photograph was taken the ulcer had crept up towards the scalp without transgressing the posterior limit seen in this photograph. Had it been an uncontrolled

arterial limit one would have imagined that during all this time the free collateral anastomosis would have taken its share in inducing the spread in all directions, but the posterior limit of the fifth cranial nerve was not transgressed.

In Fig. 12 the rodent ulcer began at the position of the cross mark, and at first occupied the position of the dotted lines. The lowest level (which it almost at once reached and has maintained so far throughout) corresponds with the posterior limit of the second cervical posterior spinal root ganglion, as demarcated by Professor Sherrington on monkeys (see Figs. 22 and 23).

It is in the monkey 1.5 cm. below the external occipital protuberance, and therefore below the attachment of the posterior belly of the occipito-frontalis muscle, hence it is at once impossible to give this muscle any share in the limitation of the growth at this line. Also the direction in which the ulcer first spread was towards the maximum point of this area, and transversely across the direction of the fibres of the muscle to which it is in topographical relation. The diagram shows the distribution of the ulcer six months after the case was first seen. It is still seen to be in the second cervical nerve area, and has not transgressed the posterior limit at which it has remained from the start. Again, it is exceedingly difficult to see how an uncontrolled arterial distribution could cause limitation posteriorly and allow its progress forward. The direction of the blood flow and the comparatively smaller

Fig. 19.—Dr. Sequeira's case of rodent ulcer.

vascular supply behind this line would not account for its sharp definition at the back and side of the neck.

It may be of interest to remember here that a muscle does rarely exist which joins the posterior belly of the occipito-frontalis with the platysma on the cheek, the occipito-facialis muscle, even supposing the existence of the muscle or its undeveloped representative, neither could account for the limitation, as the line is too low. There is an important line which in the macacus rhesus

extends from the mid-dorsal line of the scalp at about the junction of the posterior with the middle third of the distance from the glabella to the external occipital protuberance. Thence it takes an almost rectilinear course to the base of the pinna about 4 millimetres behind the anterior surface of that structure; the boundary then ascends the hinder (mesial) surface of the pinna, finally attaining to and winding over its upper edge. On the anterior (lateral) surface of the pinna the boundary descends a little in front of the fossa of the helix, sweeps forward below the concha to the tragus, and then dips over it slightly into the concha, and winds out again in the notch between the tragus and the antitragus. Having thus left the ear, the boundary line proceeds along the outer surface of the vertical ramus of the lower jaw, running slightly in front of the posterior edge of it. It crosses in front of the angle of the jaw, and turns horizontally inwards toward the mid-ventral line of the neck, which it reaches a few millimetres above the thyroid cartilage (Sherrington). See V in Fig. 22 or 23.

FIG. 13.—Rodent Ulcer, Guy's Hospital Museum.

FIG. 14.—Rodent Ulcer, Royal College of Surgeons' Museum.

FIG. 15.—Rodent Ulcer, Royal College of Surgeons' Museum.

FIG. 30.—Mr. Eve's case of Tar Cancer at the bend of the elbow.

FIG. 24.—Rodent Ulcer, the Cancer Hospital Museum.

FIG. 29.—Rodent Ulcer of left orbit, scalp and ear, Westminster Hospital Museum.

FIG. 36.—Dr. Lunn's Case of Rodent Ulcer.

FIG. 37.—Rodent Ulcer, St. George's Hospital Museum.

Fig. 20.—From Gray's *Anatomy*.

Compare Fig. 22 with Frohse's posterior lines of the human fifth cranial nerve (Fig. 21), and compare these lines to the rodent ulcers in Figs. 18, 19, 27, 28, and 29 (special plate). This line transgresses all possible muscular boundaries, and even vascular relations must be secondary to it.

The rodent ulcers of Figs. 24 (special plate) and 25 represent a type of distribution which I have seen partly mapped out in six cases, one of which is a squamous ephthelioma, kindly sent from the Hague by Dr. Schoemacher. Compare these two rodent ulcers with Heiberg's conception of the fifth cranial nerve (Fig. 26) and Dr. Head's mandibular area. There are exceptions to my statements, and they occur when cancer begins in mixed

Fig. 21.

Fig. 22.—A, nipple; B, the mid-ventral line of the limb. Fig. 23.—A, the mid-dorsal line of the limb.

Fig. 25.—Mr. Lynn Thomas's case of rodent ulcer which began on the mandibular maximum point.

The man shown in Fig. 36 (special plate) has completely lost those sensations of pain and touch over the left superciliary ridge, and incompletely lost those sensations on either side of it (at the shaded areas), and it will be seen that the spread of the rodent has respected the horizontal line of skin at which anaesthesia and analgesia begin. It must be said that the seventh nerve supply to the cutaneous muscles of the forehead has been destroyed. Although it began at the left inner canthus, the ulcer is not spreading over that line, at which it has been situated for at least six months, although its spread has been in all other directions.

Fig. 26.

Mr. T. H. Cheattle, Burford, Oxon, has now a case of rodent ulcer under his care which began in the same place as Fig. 36, and occupies the same area and has the same absence of pain

and touch sensations in the same region. The disease has been under observation for seven months, and the ulcer has spread in all directions except in the anaesthetic and analgesic area on the forehead, where it has remained stationary.

Compare these cases with Figs. 14, 15, 18, and 37. A very limited ulceration of the scalp over the frontal eminences in these cases has occurred, and, from the amount of ulceration of the superciliary ridges and sides of the heads the cutaneous nerves had gone. Now from the histories of these two living cases limitations at these lines probably have been of com-

Fig. 27.—Rodent ulcer (St. Bartholomew's Museum).

paratively long duration, because so much ulceration has extended so far and so deep from the point of incidence as to lend great support to my suggestion that the destruction of the nerves is in causal relation to the limitation. In fact the distribution of Fig. 36 and Mr. Cheattle's case in marked degrees resemble that of Fig. 37 (special plate).

In Fig. 37 the escape of the skin supplied by the mental branch of the third division of the fifth cranial nerve is very suggestive that the destruction of this nerve (which has certainly occurred as it leaves the mental foramen) has been the cause of the tissue's immunity of spread; the same may be said of the mental region in Fig. 14. In Fig. 18 the immunity of the scalp is explicable by the same reasons.

Fig. 28.—Dr. Harrison's (Bristol) case of rodent ulcer.

It may well be asked how, if this is so, Figs. 18, 19, 27, 28, and 29 are so extensively ulcerated before limitation by ulcerated nerves. The answer is that the rodent generally spreads superficially over large areas, and only penetrates deeply in the nerve strata later, and then only in small parts of them. It will also be seen that the rodents have attacked the supra-orbital nerve area from the side, and would likely destroy the nerve branches longitudinally instead of its trunk at the point of emergence, and thus the parts invaded have not been rendered analgesic and anaesthetic.

With this relationship in mind, compare this state of things to the ulcer in Fig. 19, in which the auriculo-temporal nerve is probably destroyed. The spread of the disease and its occupation of the auriculo-temporal nerve area ~~must have occurred before the nerve was destroyed, and its subsequent spread after destruction of that nerve was probably due to the presence of the supraorbital and other branches of the fifth cranial nerve.~~ If the spread or normal spread of the rodent ulcer demands the presence of the tactile or pain sensory nerves, how is it that lines like that on the scalp of Fig. 29 and the posterior line in Fig. 19 are obtainable? If there is truth in the last observation just noted, it is complicated by these cases definitely proving that limitation can also occur in the presence of active nerve impulses, which may be explained that the one set of active nerves are in a different condition of irritability to the opposing set; the difference of this irritability being due to the profound physiological changes, the possibility of whose presence I have already mentioned. But the matter is further complicated by the presence of the

Fig. 33.

second cervical overlap in the auriculo-temporal region; and the question may well be asked if involvement (whether anabolic or catabolic) of nerves is concerned, why does not it (the second cervical nerve root overlap) assume the function and aiding the diffusion of the growth? The full explanation is wrapped up in mystery probably for many reasons, and at present one can only say that in the spheres of influence belonging to two or three overlapping roots nobody can tell with certainty which root is predominant or functional. Besides, the question is still further complicated by the fact that in the lowest form of reflex act an afferent impulse does not reach the centre, but, as it were, "short circuits," and passes down the same nerve as an efferent impulse to reach the periphery; hence the spread of a cancer may not be due to only central nerve impulses. For example, an efferent impulse in a vaso-dilator effect is capable of demonstrating peripheral expression by either the higher or the lower form of reflexes to which I have just referred.

If my observations are confirmed and it is proved that cancer does not spread in areas deprived of certain nerve impulses, then the importance of the thing is not merely academic; practical questions from the point of view of treatment arise. In considering this possibility it must be

realized that the abolition of the nerve impulses by the destruction of a peripheral nerve by a cancer is permanent so long as the cancer exists, and that the destruction by excision of a peripheral nerve is not so permanent, and that pain and tactile sensations can recur in a spot from which cancer has been excised. Possibly the curative action of x rays are related to trophic changes which of necessity include nerve changes.

The knowledge and the extent of these areas of occupation and hence the likely directions of spread would be a fundamental guide in making incisions so long as the removal of cancer by the knife remains a recognized practice.

I have shown that the spread and limitation of cancer in the skin and muscles bear a marked relationship to the presence of nerves, and the absence of nerves where normally they previously existed bears a marked relation to the limitation of cancer. But I cannot say whether cancer can genetically arise in a completely denervated area. Dr. Trevor kindly allowed me to cut a section from the forehead skin in Fig. 37, and Dr. Lunn has allowed me to do the same in Fig. 36. One can detect some rodent ulcer cells among the tissues, hence I know this disease can exist for a considerable time in a completely denervated area, although it does not spread in that area.

For fear of falling into the error that rodent ulcer alone follows definite areas, I would remind the reader that I pointed out that remarkable limitation of cancer, and curiously enough benign but diffuse papillomata of the tongue show remarkable middle line limitations.

The squamous epithelioma beginning on the lip or elsewhere on the naso-labial region, described by Dr. Head, with remarkable regularity covers the whole area, and apparently rather than leave that area it fungates.

I must offer my best thanks to Dr. Sequeira for sending me many cases; to Dr. Lunn, Dr. Harrison of Bristol, and Mr. Lynn Thomas I owe similar debts of gratitude. The ideas contained in this article are rendered more tenable to those who possess a complete knowledge of the published works of Dr. Head and Dr. Sherrington.

REFERENCES.

- 1 BRITISH MEDICAL JOURNAL, April 18th, 1903. 2 F. S. Eve, On the Relation of Epithelioma to Irritation and Chronic Inflammation, BRITISH MEDICAL JOURNAL, April 2nd, 1881. 3 *Brain*, vol. xvi, p. 1 and vol. xvii, p. 339. 4 *Philosophical Transactions of the Royal Society of London*, Series B, vol. (1898) pp. 45-126. Experiments in Examination of the Peripheral Distribution of the Nerves of the Posterior Roots of Some Spinal Nerves, by Charles S. Sherrington, M.A., M.D., F.R.S.; and *Die oberflächlichen Nerven des Kopfes*, von Dr. med. Fritz Frohse.

An Address

ON THE

NATURE AND PHYSIOLOGICAL ACTION OF RADIUM EMANATIONS AND RAYS,

WITH OBSERVATIONS ON OTHER RAYS.

Delivered before the Royal Scottish Society of Arts.

By DAWSON TURNER, M.D.,

President of the Society: Physician in Charge of the Electrical Department, Edinburgh Royal Infirmary.

OF all the scientific discoveries of the last year one stands out pre-eminent, not only from its scientific importance, but also from the great public interest it has excited. I refer to the discovery by Monsieur and Madame Curie, of radium. This discovery was not, like that of the Roentgen rays, made suddenly, but had been led up to by the researches of Becquerel and of Roentgen, and was ultimately due to the painstaking and laborious analytical work of the Curies. Roentgen's great discovery of a radiation that would render the human body transparent was made in 1895, and Becquerel followed this in 1896 by finding that uranium and some of its compounds emitted a radiation akin to that of the Roentgen rays, but in very feeble amount; lastly the Curies, after examining almost every known element, found that thorium and certain ores of thorium and uranium—namely, the pitchblendes—were also radio-active, and that the latter were even more energetic than the element uranium itself. The next step they took was to separate pitchblende into its constituents, each of which was carefully tested for radio-activity; proceeding slowly and laboriously by this method they gradually separated the chief source of the radio-activity of pitchblende—the elements polonium and radium. As my interest

in this subject is chiefly medical, I will commence by referring to the forms of radiation which have been more or less recently made use of in practical medicine and surgery. They can be divided into five groups, as follows:

1. Radiant heat rays.
2. Ordinary light.
3. Ultra-violet light.
4. Roentgen rays.
5. Radium emanations and rays.

The *effluve* or brush discharge produced by high-frequency currents and by static electricity has some of the properties of the last two.

Light.

About three of these forms of radiation a good deal is known. Thus, below the ordinary spectrum of white light we have the long radiant heat waves, while above are the short ultra-violet rays; the latter, together with the blue, indigo, and violet, are called after the great Danish doctor who first used them in the treatment of disease, the Finsen light. The chief properties of ultra-violet light are that it is:

- a. Powerfully actinic.
- b. Can excite fluorescence.
- c. Can discharge an electrified body, particularly if it be negatively electrified.
- d. Has clinical and bactericidal effects.
- e. Can produce nuclei for cloud condensation in moist air.

Roentgen Rays.

Coming next to the Roentgen rays we find that their nature is still unknown, though evidence has been accumulating in favour of Sir George Stokes's theory that they are due to irregular impulses in the ether comparable acoustically to a noise. Thus it has been shown by Blondlot that they travel with the velocity of light and that the rays are polarized, but they differ from ordinary light in not being capable of regular reflection or refraction: they can, however, excite fluorescence, affect a photographic plate, and discharge an electrified body. The latter properties are shared by ultra-violet light, so that the Roentgen rays more closely resemble ultra-violet light than ordinary visible light. They differ chiefly in their powers of penetration; ultra-violet rays have but little penetrative power as regards our tissues and are absorbed by the thinnest film of blood, while the Roentgen rays pass through our skin and soft structures with great facility, but throw shadows of the bones. Transparency to Roentgen rays is a question of density, but transparency to ultra-violet light is selective; thus rock-salt and ice are transparent, while glass, mica, and gelatine are opaque. Further, the Roentgen rays appear to have no bactericidal effect, while ultra-violet light has marked bactericidal effects; a very important clinical difference is this, that while the Roentgen rays will produce, if pushed, a severe inflammation and destruction of the skin and nails, the Finsen light never produces such untoward effects.

Whether the clinical effects of the Roentgen rays are due entirely and purely to the rays, or whether the electro-static field and ionization round the tube also play a part, is an unsolved and difficult problem; there are for instance certain diseases, principally of the skin, in which the Roentgen rays usually produce a curative effect. I have, to test this question, treated three such cases with the cathode breeze and have produced the same beneficial results, and in fact in one case where the Roentgen rays utterly failed the cathode breeze was successful. These results favour the view that the curative action produced by exposing a patient to an x -ray tube is not due wholly to the Roentgen rays.

Radium Emanations and Rays.

The distinguishing radio-active properties of radium, according to Professor Rutherford, are that it gives off a radio-active emanation and three kinds of rays— α rays, β rays, γ rays.

It also maintains itself constantly at a higher temperature than its surroundings, and it continually emits heat at an estimated rate of about 100 calories per gram per hour. It is thus in its behaviour in outstanding antagonism to some of our fundamental physical laws.

The emanation is of the nature of a luminous gas which can be condensed by intense cold and which imparts radio-activity to objects in its path; more than two-thirds of the heating effect of radium is due to this self produced emanation, and 1 c.cm. of it would probably instantly melt down the glass tube which contained it.

The α rays consist of a stream of positively charged particles of mass about twice that of the hydrogen atom and projected