which it performs jointly with other organs is apt to become defective in old people, especially the gouty. The skin of old people is mostly drier than in earlier life, partly, no doubt, from the obliteration of some of the capillaries. All active exercises help in keeping the skin more active, but one of the most powerful means is the bath. Many persons with an active circulation can use cold baths from early life to very old age, others with poor reaction do better with a hot bath. Again, others begin best with a warm bath, and then let in cold water and sluice themselves, head included, rapidly with cold water at the end of the bath. The active rubbing and drying with a rough towel after the bath does not only dry and massage the skin, but obliges the bather to make many movements with the arms, which he would not do otherwise. Thus the bath acts as a kind of gymnastics to the skin itself but induces also gymnastics of the arms. One of the great advantages of the bath, especially the cold bath, is that it hardens us against chills by increasing our resisting power, in a similar way as the constant stay in the open air does. It is most useful to combine with the bath the sponging or sluicing the whole head with cold water and thoroughly rubbing and drying it afterwards. Many people can keep themselves free from "colds" by this process. This hardening is further helped by keeping the head uncovered whenever possible not only indoors, but also in walking and driving. If a full bath cannot be managed it is advisable to take a hip bath, and to sluice the head with cold water; and if this, too, is impossible, the whole body may be rubbed with a wet towel.

Before I leave the attention to the different organs of the body I may mention that I have repeatedly seen in elderly persons some advantage from massage of the eyes, ears, and nose in retarding the decay of the corresponding senses. Massage of the thyroid gland is possibly useful in maintaining general energy. Massage of the scalp not only retards the loss of hair, but also prevent some forms of headache.

Amongst the important matters into which I am unable to enter are the clothing, the house to live in, and the climate. A regular change for six to eight weeks every year from one part of the country to another, or to the seaside, or to the Continent, to different elevations above the sea, is a great help to mind and body, and, through this, a powerful agent in the prolongation of life. The exact choice depends on individual points. To some people travel from one place of interest to another is preferable to a long stay at the same place. Travelling is, in fact, a great promoter of longevity. Old people whose mental activity begins to be stagnant ought to go to localities where their attention is stimulated by art, by history, by scenery, by the manners of the people, such as Egypt, Rome, Naples, Palermo, Florence, Venice, Athens, Constantinople, Vienna, Munich, Berlin, Paris, etc.

With increasing years the resisting power of the organism decreases, in some more and in others less; in some earlier, in others later. While in youth and in the prime of life the cold, the dampness, the high winds, the fogs, and mists are fairly well borne, they cause in many old people, especially those who neglect regular open-air exercise, catarrh, rheumatism, inflammation of the lungs, mental depression, and other ailments, and through these lead to disease, premature senile decay, and death. Warmer clothing is useful, but often not sufficient to counteract the inclemencies; it is therefore prudent for many of those who have lost much of their resisting power to spend the winter and early spring in milder climates, where their mind is cheered by sun and flowers, where their accommodating powers are less tried, such as the South and South-west of England, the Riviera, Corsica, the Canaries, Sicily, Algiers, or Egypt. In addition, the larger amount of sun in these localities and the flowers cheer the mind and render it more hopeful, and hope and cheerfulness exercise a most beneficial influence on the whole body. The first Lord Brougham showed his wisdom in this matter by spending the winters of his later life at Cannes, where he enjoyed much better health and prolonged his life considerably.

CONCLUSIONS.

The main points of my advice may be comprised in a few sentences:

Moderation in eating, drinking and physical indulgence.

Pure air out of the house and within.

The keeping of every organ of the body as far as possible in constant working order.

Regular exercise every day in all weathers; supplemented in many cases by breathing movements, and by walking and climbing tours.

- Going to bed early and rising early, and restricting the hours of sleep to six or seven hours.

Daily baths or ablutions according to individual conditions, cold or warm, or warm followed by cold.

Regular work and mental occupation.

Cultivation of placidity, cheerfulness and hopefulness of mind.

Employment of the great power of the mind in controlling passions and nervous fear.

Strengthening the will in carrying out whatever is useful, and in checking the craving for stimulants, anodynes, and other injurious agencies.

In laying down these precepts I wish not to create a life of privation, but to promote a long life and a useful and happy one to the end without suffering, and I can assure you that it has been my good fortune to succeed in a great many cases.

NOTES AND REFERENCES.

- ¹ Old Age in Health and Disease, *Med. Soc. Trans.*, vol. xx, 1897: Arterial Hypertrophy, *Med. Chir. Trans.*, 1897-8; I may also refer to Charcot's lectures on senile diseases. ² On Senile Plethora, Hunterian Society's Transactions, session 1895-6, p. 38; Rise of Blood Pressure in Later Life, *Med. Chir. Trans.*, vol. lxxxvi, 1903. ³ *Die Kunst der Verlaengerung des menschlichen Lebens*, Bonn, 1890. ⁴ *Die Kunst das menschliche Leben zu Verlaengern*, Wiesbaden, 1891. ⁵ I may refer amongst many other works to Dr. G. W. Balfour's *The Senile Heart*, London, 1894. ⁶ *On Centenarians*, 1890. ⁷ See the statistics in Dr. Savill's paper already quoted. ⁸ The loss of weight varies according to circumstances, such as the size and weight of the walker, the amount of fluid taken before starting, the duration of the walk and whether combined with much climbing, the temperature and humidity of the air, the stillness or agitation of the air. ⁹ Macmillan, 1901, pp. 71 to 148. ¹⁰ BRITISH MEDICAL JOURNAL, June 23rd, 1888, p. 1312. ¹¹ *Lectures on Dietetics and Dyspepsia*, by William Roberts, M.D. Smith Elder and Co., 1885, p. 29. ¹² *Journal der praktischen Arzneykunde und Wundarzneeykunst*, vol. v. Stück, 4, Jena, 1798. ¹³ Humphry, *Old Age*, p. 53. ¹⁴ *The Code of Health and Longevity*, 5th edition, London, 1833, p. 249.

The Harveian Lectures

ON

THE TREATMENT OF SOME ACUTE VISCERAL INFLAMMATIONS.

Delivered before the Harveian Society of London, November 5th, 12th, and 19th, 1903.

By D. B. LEES, M.A., M.D., F.R.C.P.,

Senior Physician, Hospital for Sick Children, Great Ormond Street; Physician, St. Mary's Hospital.

LECTURE III. PNEUMONIA.

It may be well to combine the suggestions of the preceding lecture into an outline plan of treatment for pneumonia, though it may involve some repetition.

Every case in which a rigor occurs and the temperature rises should be sent to bed at once in a well-ventilated room without draughts, the warmth of the room being maintained at 60° F. If pneumonia is apparently developing, a trained nurse should be obtained from the first. The temperature, pulse-rate, and respiration-rate should be observed and recorded on a chart, and this should be repeated every four hours. If the patient, when first seen, is cold and at all collapsed, it is desirable to give him a "hot pack," by swathing him in a sheet wrung out of hot water of temperature of 110° F. (the head being kept cool), and covering him with blankets. Some hot brandy and water may be given to him to drink. He should be kept in the pack for about twenty minutes, then the sheet should be removed, the patient dried quickly, and placed in a warmed bed.

When he has thus been rendered warm, let the practitioner make a careful examination of (1) the tongue, mouth, throat, glands, (2) the left heart, (3) the right heart, (4) anterior pulmonary regions, (5) posterior and lateral pulmonary regions, (6) liver, spleen, abdomen. If pneumonia is developing, it is usually possible, by a very careful percussion, to detect some slight indication of the coming trouble, and it is extremely important to make sure of the diagnosis as soon as possible, for before the expiration of twenty-four hours from the onset there is a chance of arresting the disease by vigorous treatment. There will probably be pain on one side of the chest, with somewhat limited expansion of that side in inspiration, and some slight local impairment of resonance at base or apex. Over this area there may be a very little subcrepitant *râle*, but the chief auscultatory indication will be local feebleness of breath sounds. This comparative absence of breathing

in the earliest stage of pneumonia is mentioned by Professor Osler, but is not generally recognized: it is certainly a fact.

Put two hot-water bottles to the patient's feet, and, as soon as possible (every hour is of importance), fill two icebags with small fragments of ice, and apply them as already directed over the suspected part of the lung, one in front and one behind. If the mouth and fauces are foul, a sanitas mouth wash should be employed, and the throat sprayed with 1 in 2,000 perchloride of mercury lotion. This should be repeated every three hours for the first two days. It is probably desirable in every case, for the infection of the air passages doubtless often starts from the mouth, and the spraying can be easily effected during the early days of the attack, when there is little dyspnoea. The diet should consist of milk, or milk and barley water, given every two hours, and water if desired.

The patient should be seen again the same evening and again carefully examined. Any other area of dullness that can be detected should be covered by a third icebag. If pain in the side has not been already relieved by the ice, a subcutaneous injection of $\frac{1}{4}$ gr. to $\frac{1}{2}$ gr. of morphine should be administered, and a night draught of bromide and chloral-amide should be ready if the patient does not sleep. This must on no account be overlooked.

If the attempt at arrest is successful, on the second day the dullness will be found not to have increased—possibly it may already have diminished; the air will enter the suspected area more freely, the temperature will be lower, and the pulse-rate less frequent. It will in this case be necessary simply to persevere steadily with the treatment, but the greatest care must be employed to detect any fresh inflammatory foci, and to attack them immediately. Carelessness in percussion will lose the possible chance of saving the patient from a dangerous illness.

In proof of the assertion that if a case of pneumonia comes under observation within twenty-four hours after the initial rigor it is sometimes possible to arrest it by vigorous treatment, I give the two following cases:

CASE I.—W. B., 18, carman, was seized on the evening of October 31st, 1895, sixteen hours before his admission into St. Mary's Hospital, with a rigor which lasted an hour. Next morning he had fever and pain in the right side. On admission his skin was hot and dry, and there was some labial herpes. Temperature 103.6° , pulse 120, respirations 40. When I first saw him on the evening of November 1st, twenty-four hours after the rigor, I found dullness at the base of the right lung in front below the fourth rib, extending into the lower axilla, with some tenderness. The breath sounds were feeble over the dull area. No bronchial breathing, but a little crepitation at the end of inspiration. Behind, at the right base, breathing weak, and some impairment of resonance. Three icebags were at once applied.

November 2nd. Temperature, 101.8° ; pulse, 100; respirations 34. Feels better. Dullness decidedly less extensive.

November 3rd. Temperature 100° , rising to 101.8° , falling to 99° ; pulse, 100; respirations, 30. Dullness still diminishing. Says he is "a lot better."

November 4th. Temperature, 98° ; pulse, 72; respirations, 26. Now only a small dull area in lower axilla. Ice removed (after sixty hours).

November 5th. Temperature rose to 99.8° , but fell to 98° .

November 6th. Temperature rose to 100.2° , but fell to 98° .

November 7th. Temperature, normal; pulse, 64; respirations, 20. Very slight impairment of resonance could now be detected.

CASE II.—E. N., 14, admitted May 22nd, 1896, twenty-four hours after immersion in a canal and twelve hours after a rigor. He had a headache and dyspnoea. On admission dullness was found in the right axillary region, and an icebag at once applied. Temperature, 103.5° ; pulse, 120; respirations, 40.

May 23rd (10 a.m.). Both cheeks very flushed. Obvious dyspnoea. Temperature, 103° ; pulse, 120; respirations, 40. Dull in the right lower axilla, not behind scapular line nor to inner side of nipple. Just below angle of right scapula there was distinct fine crepitation, with inspiration only; this was so typical that I made all my clinical clerks listen to it. Breath sounds diminished over the dull area; no bronchial breathing. Dullness and diminished breathing in right suprascapular fossa also. Some pain on left side of abdomen on taking a deep breath, but no rub could be heard and there was no dullness. Heart normal. Three more icebags were ordered, making four in all; two to the right base, a third over the right apex behind, and the fourth over the left axilla. After one hour the temperature fell to 100° , and the ice was removed. It then rose to 102° , but at once fell again.

May 24th (forty-eight hours after the rigor). Temperature, normal; pulse, 74; respirations, 34. He had slept well, was now not flushed, and the right axilla was less dull. The temperature remained subnormal for thirty-six hours. There was a short rise to 100° on the 25th, and to 99.5° on the 26th. After this it was normal, and the boy was quite well, and the right axillary region was of normal resonance.

It will be observed that in each of these cases there was no crisis, but an immediate and rapid subsidence of temperature, physical signs, and symptoms. In such cases as these it is

reasonable to claim that the disease has been arrested. But it is not always possible to arrest a pneumonia even when it is treated very early, and after twenty-four hours there is little hope of success. This is not surprising when we remember how rapidly micro-organisms increase in number under favourable circumstances. Washbourn and Eyre found, on cultivating the pneumococcus in nutrient broth, making plate cultivations from the broth culture and counting the living cocci present at different periods, that 140 colonies increased in three hours to 6,149, and in six hours more to 13,680; twelve hours later they were "innumerable."

As the normal temperature of the human body is only 1° F. below the optimum temperature for the growth of the pneumococcus, it is clear that if an attempt to arrest the development of a pneumonia is to have any chance of success, it must be made very early and very vigorously.

But it is always possible to influence the course of a pneumonia, to diminish its intensity, and often to shorten its duration. This of course is difficult to prove, because of the uncertainty of the time of occurrence of the crisis in the disease when untreated. But there is nothing really improbable in the assertion that there is reason to believe that the ice treatment sometimes brings about an earlier crisis. For however the crisis may be caused, whether by the manufacture of an antitoxin or by a failure of further growth of the pneumococcus, it seems clear that any treatment which can to any extent inhibit the growth of the microbe and thus check the amount of toxin which it produces, will to that extent facilitate the earlier termination of the struggle between the attacking and the defending forces, in other words, it will hasten the crisis.

If the attempt to arrest the disease is unsuccessful, on the second day the area of dullness will be larger, and over it may be heard inspiratory crepitation, or sharp *râles* of double rhythm in children, or some prolonged expiration, or distinctly bronchial breathing. A third or a fourth icebag should now be applied, the sites for their application being outlined in blue.

It is desirable at this period to administer two or three grains of calomel, followed after three hours by a seidlitz powder. When a sufficient evacuation has been obtained the purgative should not be repeated, for in the later days of a pneumonia there is a tendency to diarrhoea.

On the second evening the hypnotic must be again given if necessary, and morphine if pain is present, for the patient must have sleep. It may, perhaps, be desirable to remove one or two of the icebags during the night, leaving two only in position. It might be thought that the necessity for disturbance would be fatal to sleep, but the relief of pain and dyspnoea is so great that the patient easily falls asleep again, provided that his right heart is not over-full. In the case of young children the temperature should now be taken every two hours (hourly for babies), and it can be done without disturbing them. If any icebags have been removed at night, they should be replaced early next morning.

On the third morning the physical signs in the lungs must again be most carefully determined, and directions given for the alterations of position of the icebags necessitated by the changes found. But now special attention must be given to the right heart. If the dullness of the right auricle is found to extend two finger-breadths in the fourth right space, and there is distinct dyspnoea and some slight lividity of lips, or cheeks, or finger tips, leeches should be applied over the lower ribs on the right side below the nipple level. One should be used for a baby under six months, two for a child under two years, four for a child of ten years, six for an adult, eight for a robust man. If not used at once the leeches should be held in readiness, for they may possibly be required in the evening if the patient is to sleep. Some tins of malted milk, and one or two cylinders of compressed oxygen should be procured.

On the third evening it will in most cases, unless the ice has already caused a marked improvement in the physical signs, be advisable to apply leeches—if they have not been already used—an hour or two before the time for sleep. The relief thus given to the right heart will often induce sleep without any hypnotic, but one must be given if needful. Even morphine may be used safely under these circumstances.

On the fourth morning, if the leeches have been applied, the patient will feel more comfortable, though the physical signs may have increased in extent. The same minute care in determining the physical signs in both lungs must be

practised. Watch carefully for fresh areas of dullness, especially if there has been any fresh rise of temperature, and attack them at once.

The right auricle having now been relieved it will be desirable to give considerable quantities of water, both to satisfy thirst, and to promote diuresis and the elimination of toxin. During the twenty-four hours following the use of the leeches, 3 or 4 pints of water may be given, in quantities of 8 to 10 oz. every three hours; for a child 4 to 6 oz.

If the patient has not come under treatment until the fourth day of a severe attack, he will probably be in considerable distress. Dyspnoea and discomfort will be marked, cyanosis distinct, and the dullness of the right auricle may measure from two to two and a-half fingerbreadths in the fourth space, one or one and a-half in the third, and half a fingerbreadth or more in the second. This should be ascertained at once, before any attempt is made to discover the amount of disease in the lungs. The call for bleeding is urgent and imperative. A larger number of leeches must now be used than would have sufficed on the previous day. Two must be employed for a baby, three for a young child, four to six for an older child, eight to twelve for an adult. A venesection is often preferable: 4 oz. for a young adult, 8 oz. for a strong man.

An hour after the bleeding both lungs should be carefully examined and the outlines of the dull areas marked on the chest. Two icebags must be applied at once to the worst inflammatory foci, an hour or two later a third, and before long a fourth. We are now in the thick of the fight, and it is necessary to call up the reserves and have all our forces in readiness for the struggle of the next three or four days. The subcutaneous injection of strychnine should be begun and maintained systematically in increasing amount or frequency. Now is the time also to begin the administration of oxygen; this, too, should be regularly continued throughout. Milk and also water may be given in considerable quantities after the venesection. At night sleep will probably come naturally, the right heart having been relieved and the pulmonary congestion diminished by the ice, but if not a hypnotic must be given, and even morphine if necessary; the patient must have sleep.

On the fifth day, if the patient has been vigorously treated with leeches and ice, there is often a marked improvement in the physical signs, and much less tendency to extension. But a severe case may be not yet controlled. It will now be desirable to limit the amount of fluid given to the patient, so as to lessen the strain on the right heart. The diet for the next two or three days should be simply malted milk powder dissolved in milk, a tablespoonful in two ounces every two hours for an adult, two teaspoonfuls in one ounce for a child. The icebags must be continuously applied, and their position altered as may be necessary, special care being taken to discover and attack fresh or spreading areas of inflammation. If leeches have been used on the third day, it is desirable to examine the right heart again, very carefully on the evening of the fifth day. The relief will almost always last for forty-eight hours, but by the fifth evening some more leeches may be required. In determining this point, especially when the left lung is mainly involved, it is very necessary to guard against being misled by overdistension of the right lung into an underestimate of the size of the right auricle. In case of doubt, let the decision be for the leeches. The amount of sleep which the patient has hitherto obtained is also of importance in deciding this question. If he has slept well, and the right auricle does not measure more than two fingerbreadths, the leeches may be postponed. But if sleep has been defective, it will be wiser to apply them, and afterwards to give a hypnotic. For sleep is of great importance for the maintaining of vigour for the days which may remain.

If the patient has reached the fifth or sixth day of his illness, and neither bloodletting nor ice has been employed, the symptoms are often very severe, the distress great, and the outlook gloomy. Probably he has been sleepless for several nights, and his strength is rapidly diminishing. The call for active treatment is urgent. The prognosis depends on three factors: the age and previous health of the patient, the intensity of the infection, and the action of the medical attendant. At such a time the responsibility of the latter is great indeed. Life is trembling in the balance. His action or his inaction may decide whether or not the patient shall be deprived of many years of life, and his wife and children suffer an irreparable loss.

The first necessity is a venesection. Eight ounces of blood should be taken at once, twice as much or more if the lung be

full of *râles*. If permission for venesection cannot be obtained, place a dozen leeches over the liver and encourage the bleeding. Hypodermic injections of strychnine in 3-minim doses every four hours should follow immediately, and the systematic inhalation of oxygen for ten minutes or more every hour. Two icebags should be at once applied, soon followed by a third, and before long by a fourth. It is very probable that after this treatment the patient will fall asleep. If so, he should be undisturbed for four hours. But after this nourishment must be given and the icebags refilled every two hours. Malted milk in milk with a little brandy should be given every hour when he awakes, and all medicine by the mouth avoided. After sleep has been obtained a small enema may be given if necessary. If diarrhoea is present the rectum should be washed out with warm saline solution, and 2 oz. of starch decoction with a few drops of tincture of opium inserted.

Some improvement—often much—will certainly follow this treatment unless the patient is already very far on the downward road, or his heart be previously dilated, his lungs emphysematous, his liver cirrhotic, or his kidneys granular. Many cases are no doubt hopeless from the first; but not very rarely an apparently hopeless case recovers; and, at all events, whatever can be done to give a chance of recovery ought to be done. Here let me put in a plea for earlier consultations. Too often a "second opinion" is sought for only when death is imminent. The surgeon is right in asking that he may be allowed to see a case of perforated gastric ulcer as soon as the diagnosis is made; if twenty-four hours are allowed to elapse, the patient's chance of recovery is small indeed. It is estimated by Mr. Mayo Robson that if operated on within twelve hours after the perforation the mortality is only 16.6 per cent.; if within twenty-four hours, it is 63.0 per cent.; if within thirty-six hours, it is 87.5 per cent.; and if delayed for forty-eight hours, the operation will only rarely succeed. So may the physician plead that in pneumonia the final issue largely depends on the treatment, or want of treatment, of the first few days. In the case of an infant, or of an adult older than 30 years, the danger to life is great, and judicious treatment is required from the very first. This is not so obvious to the patient as when an operation is required, but it ought to be equally obvious to the practitioner. To delay the consultation in such a case to the fourth or fifth day is to imperil the patient's life.

Pneumonia in adults usually ends by a very rapid fall of temperature, with slowing of pulse. This "crisis" often occurs in children also, even in cases which would be designated as "broncho-pneumonia," but in children the subsidence is apt to be more gradual, and to occupy several days. Pneumonia in children sometimes lasts for three or even four weeks. It is necessary to keep a careful watch for the first indications of this quick diminution of temperature, especially in children. The icebags should be gradually removed as the temperature falls, and the last should be taken off when the thermometer marks 100° F. There is a natural tendency to collapse at the time of the crisis which must be kept in mind. Icebags over the chest at this stage would probably be injurious, though in pericarditis, as I have already mentioned, they may sometimes be used with advantage, even when the temperature is subnormal. If, in spite of care, or for want of it, the reduction of temperature is so great as to cause some collapse, it is desirable to apply warmth over the heart and abdomen, also to the feet, and to give the patient some hot water with brandy, and a draught containing ether and ammonia. By these means it is usually easy to remove any tendency to collapse.

But the crisis is often preceded by a remission of temperature which lasts only a few hours. Hence, when the ice has been removed, the temperature should still be taken every hour. If it rises to 101° F. a single icebag should be again applied, if to 103° F. at least two. Much careful observation on the part of the nurse is required at this period. If the temperature shows a persistent tendency to keep above normal after the crisis has occurred, the existence of empyema should be suspected, and an exploring needle should be passed into the dullest area. Occasionally it may be due to tuberculosis.

EMPHYEMA.

The frequent occurrence of empyema as a chronic condition following an attack of pneumonia is well recognized, but I wish to call attention to what may be termed acute empyema, in which the pneumococcal invasion mainly or exclusively involves the pleura, the lung being little affected, perhaps not at all. This seems to be more common in children, but it may

occur in adults. In a young child in the Hospital for Sick Children, who had a temperature of 106° F. and dullness over one base and who was thought to be suffering from pneumonia, the necropsy revealed a considerable amount of turbid fluid in the pleural cavity, but no consolidation of lung. In a girl recently under my care at St. Mary's Hospital, 3 oz. of pus were obtained from the left pleura as early as the eighth day of an attack, which was at first thought to be merely pneumonia, but in which the extensive dullness, feeble breath sounds, and cardiac displacement soon suggested pleural effusion. I think that in this case some pneumonia was present also for cyanosis was well marked, and the dullness of the right auricle in the fourth right space measured two and a-half finger-breadths on the third day of the illness.

When the amount of pus in the pleura is small and localized by adhesions, it is often extremely difficult of diagnosis. Probably few physicians have escaped the mortifying experience of the *post-mortem* discovery of an empyema which had been overlooked, or diagnosed as some other condition. The difficulty of diagnosis is similar to, though not quite so great as, the difficulty in suppurative pericarditis, and the two conditions are sometimes associated. There may be a small amount of pus in both pleurae as well as in the pericardium; it is rare to find any in the peritoneum.

A localized empyema may be diagnosed as a pneumonia, as a serous pleural effusion, as a bronchiectasis, as tuberculous consolidation of lung, as collapse of lung, or as a subphrenic abscess, and the diagnosis may be impossible without the exploring needle. The signs of distinction between a consolidated lung and a pleural effusion may be neatly stated and tabulated in textbooks, but there is not one of them which may not prove fallacious in practice. The temperature may be high in an acute empyema; it may be low in pneumonia. The breath sounds are sometimes audible in children over a pleural effusion, and they may be quite inaudible in an influenzal broncho-pneumonia, or in the later stages of a lobar pneumonia. Even displacement of the heart is not certain proof of a pleural effusion, at all events in children; it may sometimes be caused in them by a solid lung. This is especially true of a form of tuberculous disease of the lung in which the lower and middle lobes on the right side are attacked by tubercle spreading from the bronchial glands. This occurrence is not very rare in children, and my colleague, Dr. Batten, has shown from *post-mortem* observations that it is much more frequent on the right side than on the left. Clinically this condition may cause all the physical signs of a pleural effusion, including some displacement of the heart, as I have seen in a case in which an incision revealed a healthy pleura, with solid lung beneath.

In some cases a small localized empyema between the lobes of the lung may coexist, as in a girl under my care at Great Ormond Street. In this case Mr. Collier succeeded in draining the small, deep-seated cavity, and the patient gradually recovered.

The danger of overlooking an empyema which is small, or even of moderate size, is so great that it should be always kept in mind, and an exploring needle should be passed if there is any room for doubt. If the first puncture does not yield pus, an anaesthetic should be given, and punctures made in other sites of dullness. If pus is found an incision should at once be made, a piece of rib removed, and a drainage tube inserted.

PLEURISY.

Acute pleurisy in its earliest stages may be very successfully treated with ice to the chest. One icebag is often sufficient; it should be applied over the spot where friction is audible. It quickly relieves pain and often renders morphine unnecessary, though it might be wise to give a small amount of this drug hypodermically if the pain were very severe. Two icebags may be used if necessary. The friction often rapidly disappears after the ice has been applied for some hours, and the tendency to effusion is very decidedly checked. As an illustration of this beneficial effect, I may mention a recent case of dermoid cyst of the right thorax, under my care at St. Mary's Hospital:

A large cyst displaced the heart into the left axilla and compressed the right lung. By four distinct operations, at considerable intervals, Mr. Silcock succeeded in removing the whole of the wall of the cyst, dissecting it off the diaphragm, the left pleura, and the pericardium. On the day following the last of these operations the patient complained of pain in the left side, and the temperature rose. Over the cardiac region and to the left of it a fine crepitant friction could be heard, with the respiration only. An icebag was applied. Next day there was a definite pleural rub in the left axilla, and pain continued, though less intense.

On the day following the pain was much relieved and the rub less marked, but there was much inspiratory dyspnoea, both sternal and mastoids acting strongly with each inspiration. It was impossible to determine accurately the extent of the dullness of the right auricle, owing to the previous condition and to the operation, but it was clear that it was greatly overstrained through the occurrence of pleurisy and probably acute pulmonary congestion in the single lung that was able to perform the respiratory process. We were very reluctant to submit the patient to any further loss of blood, and fortunately we succeeded by a subcutaneous injection of morphine and atropine in escaping this necessity. Next day the danger was past, and the patient steadily recovered.

I think it is practically certain that but for the icebag this patient would have died.

In the treatment of a large serous effusion ice externally is very useful, but it should be preceded by paracentesis. As soon as it is evident that there is a considerable amount of fluid in the pleura, it is best to aspirate at once, under low pressure, and draw off as much fluid as comes easily. Then fix this side of the chest with broad bands of strapping firmly applied, and over the strapping apply two, or three, icebags. This treatment rapidly relieves the discomfort and dyspnoea, and any pain that may be present. If the ice is maintained in position for about two days there is usually little tendency to reaccumulation of fluid. But if a relapse does occur, it may be treated in the same way, and the ice applied for a longer period. The serous fluid obtained by paracentesis in such cases is usually sterile; it is probably often due to tuberculous infection of the pleura. It is frequently possible to obtain physical evidence of this by a careful examination of the *opposite* lung, for there is in many such cases more or less dullness and feebleness of breath sounds at the inner end of the first space and over the posterior part of the upper apex also below the apex of the lower lobe. This is the more striking because the rest of this lung is often hyper-resonant.

In an acute local pulmonary tuberculosis icebags applied over the dull apices often give much comfort and appear sometimes to be of distinct service in checking the rapid progress of the disease. This is especially the case when the temperature is high.

In acute laryngitis an icebag placed over the larynx gives rapid relief to the symptoms, and in a catarrhal laryngitis is curative.

In local peritonitis over gastric ulcer, with threatening perforation, ice over the epigastrium brings much comfort to the patient, and greatly aids the subsidence of symptoms.

APPENDICITIS.

In the treatment of inflammations of the appendix vermiformis the persistent application of an icebag is far more effective than fomentations or poultices; it rapidly relieves pain, and obviously diminishes the local inflammation. Clinically, cases of appendix-inflammation may be divided into three groups. The first and most important group consists of cases of perforation of the appendix, with escape of concretions or of pus into the peritoneal cavity, or of a gangrenous condition of the appendix itself. Such cases are usually of sudden onset, and the symptoms often severe. Pain, vomiting, tenderness, flexion of the thigh, and absence of descent of diaphragm during inspiration, perhaps with rapidly-increasing evidence of septic absorption, but an absence of any local swelling, are the chief indications. Such cases demand immediate operative interference; every hour is of importance. They are similar to cases of perforated gastric ulcer, or to a strangulated hernia. No question of palliative treatment must be entertained for a moment. The absorption of septic material from the peritoneum is often very rapid, and a delay of a few hours may make all the difference between life and death. No such case ought to be allowed to die without operation; not seldom, apparently desperate cases have been rescued. The operation ought never to be delayed because of the collapsed condition of the patient; it is the most effectual means of removing the collapse—often the only chance.

The cases of the second group are much more common. They are often due to a subacute inflammation of the appendix, which may be chronically thickened. The symptoms are comparatively slight, but there is usually some definite local tenderness, and often a tender local swelling can be detected, with dullness on percussion over it. This swelling may sometimes be due to a local muscular contraction in the abdominal wall, as suggested by Dr. Mackenzie, but it certainly appears in some cases to be the thickened appendix itself which is felt.

These cases will usually recover in time under any or under no treatment, but they are prone to relapse. Fomentations or poultices give some relief to discomfort, but they do not cause rapid subsidence of the swelling and of the symptoms. This, however, is often promptly effected by an icebag, the results produced by which are sometimes very striking.

Intermediate between these two groups of cases, clinically speaking, is a third group, in which the symptoms are very definite, but not so acute as to make it clear that immediate operation is required. There is usually a distinct local fullness and tenderness. The practitioner is in doubt whether pus has already formed, and whether an incision ought to be made at once or delayed. In this difficulty, which is by no means rare, and in which a correct decision is of the greatest importance, I have found that the application of an icebag for three or four hours will often give the necessary guidance. If after four hours the ice has not given distinct relief, an operation should be performed without further delay. But if there be definite, even though slight, alleviation of symptoms, the operation may be postponed for a short time and further trial made of the icebag. In many cases after twenty-four hours the relief is so definite that no further question of immediate operation need be entertained. If the improvement continue, the icebag should be kept in place persistently for two or three weeks, with absolute rest in bed. Such cases must be very carefully watched, for a small local abscess may remain and may require incision at a later period. But the condition will then be much more favourable for operation than at first, the pus will be well localized, and the surrounding inflammation quietened down. On the other hand, if the early relief is not maintained, although the icebag has been applied persistently, it will usually be wise to delay the operation no longer. Thus the icebag becomes a useful test by which we may decide the often difficult and anxious question as to whether or not an operation ought to be performed.

If recovery has occurred without operation it must be decided whether or not the appendix should be removed. If this is the first occasion on which it has offended, and the symptoms have been slight, it may surely be allowed another chance, though this is not the opinion of the most "advanced" surgeons. But if the symptoms have been severe, or if it be a second offence, probably removal after a short period of convalescence is desirable.

ACUTE NEPHRITIS.

The last form of acute visceral inflammation of which I desire to speak is acute nephritis. The method of treatment usually advised for this serious malady is often very unsatisfactory in its results. Recovery is slow, and in a large number of cases imperfect. In the worst cases, if death from acute suppression or rapid uraemia does not supervene, the patient passes into a condition of chronic albuminuria and dropsy, becoming, as my former teacher the late Dr. Moxon expressed it, "a large white man with a large white kidney." In the majority of cases more or less albuminuria remains after apparent recovery, and in many there is relapse of inflammation in the damaged organs, leading directly or indirectly to a fatal issue.

Even the etiology of acute nephritis is very imperfectly understood. Scarlet fever is, of course, responsible for a very severe form of nephritis. Diphtheria often causes transitory albuminuria, more rarely a real nephritis. The same may be said of influenza, typhoid fever, and pneumonia. The rheumatic diplococcus has been found in the urine of rheumatic patients, and it must in the future be a subject for inquiry whether an attack of acute rheumatism may possibly be responsible for slow insidious changes in the kidney, whether, for instance, it may be one of the causes of granular kidney. The "cold and wet," to which some cases of acute nephritis are ascribed, probably act, as they do in the production of pneumonia, by permitting the internal cultivation of parasitic micro-organisms, and the elimination of their toxins. Many cases of subacute nephritis in children and young adults are difficult of explanation; in many of them there has been no ascertainable exposure to cold and wet or to infection from scarlet fever. Perhaps some may be due to the injury done to the kidneys by microbes or microbic toxins from the alimentary canal eliminated by the urine. In the worst case of nephritis that I have seen in a very young child (aged 2½ years), the only fact that could be ascertained which seemed to offer any explanation was that he had suffered from an attack of vomiting and diarrhoea, which lasted about a fortnight. Soon after his recovery from this,

about three weeks from its onset, he was noticed to be "growing fat." But this "fat" was the dropsy of an acute nephritis, and his urine was solid with albumen. He was treated in the manner usually advised, and after a long and trying illness gradually made a very imperfect recovery. Considerable albuminuria remained when he was discharged from the hospital. This result is certainly not a therapeutic victory. Can we do nothing better for such patients? I think we can. The external application of icebags over the inflamed kidneys in nephritis has in several cases given me excellent results. I have less experience of the use of ice in nephritis than in pericarditis or in pneumonia, partly because the cases are less common, but mainly because of the apprehension, only gradually overcome, that the attempt might result in a fatal suppression of urine. But in this case also time has proved to me the wisdom of the old adage, "Don't think; try." For it has turned out that when used with proper precautions ice over inflamed kidneys, far from causing suppression of urine, is an excellent diuretic, rapidly increasing the amount of urine passed. No doubt there might be danger of inducing suppression if the ice were applied while the skin was dry and cold and the patient more or less collapsed. For indeed suppression may be caused by the disease itself, as I found recently in the case of a woman aged 36 whom I saw in consultation a few hours before her death on the sixth day of an acute suppression. A catheter passed into the bladder removed about an ounce of turbid urine full of albumen. No treatment had been attempted. But if a patient with nephritis is first made thoroughly warm and his skin somewhat moist by a hot-air bath, and he be kept warm by hot-water bottles, it is quite safe to apply icebags over his kidneys. Not only is it, with such precautions, quite safe, but the result is often most satisfactory.

S. S., a boy of 15, was admitted into St. Mary's Hospital under my care on May 9th, 1903, for subacute nephritis. The dropsy was slight, but the urine was reddish in colour, contained blood and epithelial and fatty casts. It was of specific gravity 1012, acid in reaction, contained albumen to the amount of one-fifth, and 1.5 per cent. of urea. The quantity passed in the twenty-four hours before treatment was begun was 34 oz. A single icebag was applied over the right kidney only, without any previous hot-air bath, as the boy's skin was warm and not dry. Directions were given that the quantity of urine passed in every successive period of eight hours should be separately measured and tested. The amount of urine on the first day of the application of the icebag was 34 oz., the same amount as on the previous day. On the second day of the ice it increased to 52 oz., on the third day it was 40 oz., on the fourth day again 52 oz. Thus the average amount of urine for the four days during which a single icebag had been applied was 44 oz., an increase of nearly 30 per cent. Two icebags were now applied, one over each kidney. A further marked increase in the quantity of urine followed immediately. On the first day of the application of two icebags the amount passed was 78 oz., on the second day 70 oz., on the third 58 oz. Thus the average amount for these three days was 68 oz., an increase of 100 per cent. This diuresis was due solely to the ice, for no medicine was given. The two bags were retained in position for ten days, with the exception of one period of eight hours, the amounts passed during these ten days being respectively 78, 70, 58, 56, 60, 53, 59, 60, 71, 70 oz., giving an average of 63.5 oz. Thus the diuresis was steadily maintained. The quality of the urine also progressively improved. It soon lost its red colour and became yellow, though some blood corpuscles could still be seen when the ice was removed after the total period of fourteen days; the casts rapidly diminished, and entirely disappeared, and the amount of albumin fell from about one-fifth to about one-tenth. The amount of urea rose to 2 per cent. After the fortnight's treatment by ice the boy was still kept in bed for five weeks, but no other treatment was used. The amount of albumen progressively diminished, and finally became only a "very faint trace."

Similar results have followed in other cases, and it is now clear that, when used with proper care, ice over inflamed kidneys has a well-marked diuretic influence by diminishing the local congestion in the inflamed organ as it lessens the congestion in an inflamed appendix or in an inflamed lung. It appears to be directly curative.

Another illustration of the diuretic effect of ice over the kidneys in nephritis may be given.

A girl aged 6, in my ward at the Hospital for Sick Children, was treated for nephritis with daily hot-air baths and diuretics. During eight days of this treatment the amount of urine averaged 16 oz. She slowly improved, and when she was sent to a convalescent home the albumen was estimated at one-third. Two weeks later she was re-admitted for fresh general oedema. The amount of urine passed on the first day after her readmission was only 2 oz.; on the second day 4 oz. A single hot-air bath was given, and then icebags were applied over the kidneys. The amount passed on the third day was 7½ oz., on the fourth day 27 oz., on the fifth 37 oz., on the sixth 36 oz., on the seventh 37 oz., on the eighth 36 oz. The effect on the albuminuria was quite as remarkable. When sent to the convalescent home the amount of albumen was estimated at one-third. When she returned it was one-fourth.

After four days of the icebag it was only one-sixth, and two days later there was only a trace of albumen present.

One more case may be related which will show how useful the application of ice may be in a very severe case of nephritis:

H. V., aged 3 years and 10 months, was admitted into the Hospital for Sick Children on November 9th, 1901, having suffered from headache, shivering, repeated vomiting for seven days, swollen face for six days, and bloody urine for two days. On admission he was pale and very restless, with a furred tongue, and temperature of 102.8°. No urine at all was passed that afternoon and evening. He vomited once.

November 10th.—Urine 2 oz., very smoky, contains blood, albumen = $\frac{1}{2}$. No casts. Vomited three times.

November 11th.—Hot-air bath, followed by two leeches over each kidney. Hot-water bottles were placed in the bed, and two icebags were applied over the kidneys. Urine about 4 oz. (not all saved). Vomited five times. In the evening the icebags were removed for four hours. Another hot-air bath was given. The icebags were afterwards replaced and kept in position all the night.

November 12th.—Temperature 101° to 102.2°. Urine about 5 oz. Vomiting troublesome. Rectal feeding adopted.

November 13th.—Child has been very drowsy since admission and restless. Very restless in the evening. Restlessness was increased by the hot-air bath. Temperature rose in the afternoon to 104°, and a patch of dullness was found at the right base, with slight pleural friction. An icebag was placed over the right base as well as over the kidneys. Four hours later the temperature had fallen from 104° to 102°, the pulse from 128 to 100, and the respirations from 50 to 40. The child had been sleeping a good deal. It was observed that he was much less restless while the icebags were in position, and became more restless when they were removed. Urine now contains a few hyaline and corpuscular casts and abundant red cells.

November 14th.—Total urine yesterday 5½ oz., albumen $\frac{1}{2}$. Still vomits anything but barley water; retains nutrients. Temperature 100.3°, pulse 100, respirations 36.

November 15th.—More than 10 oz. of urine yesterday. Friction at right base now not audible, but slight dullness remains. He is now taking malted milk by the mouth and does not vomit.

November 16th.—Better. Takes more interest in his surroundings. More than 13 oz. of urine yesterday.

November 17th.—Last evening the temperature rose to 105.4°, and the amount of urine fell to 8 oz.

November 18th.—Urine yesterday only 5 oz. Temperature 102.8°.

November 19th.—Urine yesterday more than 6 oz.

November 20th.—More urine, 9 oz. and some wet beds. The temperature fell last evening to 97.6°, and the icebags were removed for eight hours; when they were reapplied it was 100.2°; at 6 p.m. it was normal. Urine contains less blood and shows few red blood cells, but many leucocytes with abundant epithelial and granular casts.

November 21st.—Good amount of urine. Icebags left off at night, but replaced during the daytime.

November 25th.—Urine in fair amount; albumen much less. Icebags discontinued altogether, after fourteen days' use.

November 28th.—Urine in fair amount; cannot all be saved. It still contains leucocytes and granular and epithelial casts, but no blood, and only a trace of albumen.

December 5th.—Urine contains one or two granular casts, but no leucocytes, no blood, and only a trace of albumen.

December 10th.—One or two casts. Only a trace of albumen.

In conclusion, I desire to express my thanks to the Council of the Harveian Society for the honour of delivering the Harveian Lectures for 1903.

ON A DYSENTERY ANTITOXIN.

By CHARLES TODD, M.D.CAMB., D.P.H.,

Senior Assistant Bacteriologist, Serum Department, Lister Institute of Preventive Medicine.

[PRELIMINARY COMMUNICATION.]

SINCE the discovery of the dysentery bacillus in 1898 by Shiga as the cause of an epidemic of dysentery in Japan, a great deal of attention has been directed to this disease, and the work of Shiga in Japan, Kruse in Germany, Flexner and Strong in America, and others, has established the widespread distribution of the bacillus and its claim to be regarded as the causal agent. Further, the work of Duval and Bassett has shown that in the United States of America this bacillus is in all probability the cause of the summer diarrhoea of infants.

A striking point in the pathology of dysentery is that while the bacilli are found regularly in the mesenteric glands, they are not found in the spleen or other organs, and do not occur in the blood, urine, or milk. In this respect Lentz¹ points out that the disease differs markedly from the septicaemic diseases, such as typhoid, and that it must be regarded as a local infection of the intestinal mucous membrane and corresponding lymphatic glands by the bacillus, the toxin alone

passing into the circulation and giving rise to the typical clinical picture, which, as in the case of cholera, gives the impression of a severe poisoning or toxæmia.

The results of animal experiments bear out this view, as the pathological changes which result from the injection of the living bacillus may be equally produced by the use of dead cultures, showing that these changes are brought about by some toxic body, and not by any vital action of the bacillus.

THE TOXIN OF DYSENTERY.

The following experiments were commenced at the end of last year with a view to investigating the character of this toxic body, and the possibility of obtaining an antitoxin.

The filtrates from ordinary peptone broth cultures of the bacillus are only slightly toxic for animals, but it was found that by employing a more alkaline medium much more toxic filtrates could be obtained. The ordinary alkaline broth, as used for the production of diphtheria toxin (that is, alkaline broth to which normal caustic soda has been added in the proportion of 7 c.cm. per litre), was planted with the bacillus. This was grown at 37° C. for about ten days, filtered through a Pasteur-Chamberland filter, and the filtrate tested on rabbits by intravenous injection. Both Shiga's and Kruse's bacilli were tried, and it was found that the results were approximately the same. The following show some of the results:

Toxins obtained by Growing the Bacillus for Ten Days in Alkaline Broth.

No. of Toxin.	Bacillus.	M. L. D. for Large Rabbits.
No. 1	Kruse	1 c.cm.
" 2	"	"
" 3	"	"
" 4	"	"
" 5	"	"
" 6	Shiga	"

In these tests the minimal lethal dose (M. L. D.) was only approximately determined, that is, the toxin killed at 1 c.cm. and not at 0.5 c.cm.

In the subsequent experiments the bacillus of Kruse (obtained from Kral) was employed exclusively. The effect of growing the bacillus for a longer time was then tried, and it was found that by growing it for a month much more toxic filtrates were obtained—killing a rabbit at from a seventh to a tenth of a cubic centimetre (Tables I and III.)

The susceptibility of various animals to this toxin varies enormously; the rabbit, as shown above, being highly susceptible, but the guinea-pig highly resistant—5 c.cm. of the same filtrate not killing a small guinea-pig, either subcutaneously or intraperitoneally. The horse, as will be seen later, is very highly susceptible. The mouse and monkey are not highly susceptible.

The toxin is fairly stable and is not destroyed by heating at 70° C., for an hour, though 80° C., for an hour destroys it. It is not hæmolytic for the corpuscles of the rabbit. The results of the intravenous injection of a lethal dose of the toxin in a rabbit are practically the same as those obtained when the living bacillus is used; after a latent period, varying with the amount of toxin used, severe diarrhoea sets in with paralysis of the limbs, beginning sometimes in the fore and sometimes in the hind limbs, but ultimately affecting both. The animal rapidly loses weight and dies usually in from one to three days. *Post mortem* there is marked congestion of the gut, which contains a slimy mucoid fluid, the lungs shows patches of congestion and often small hæmorrhages, the other viscera, as a rule, show no marked naked-eye changes.

At the time these experiments were made the only reference to a soluble dysentery toxin which could be found in the literature was the statement by Lentz² in the article on Dysentery in Kolle and Wasserman's *Handbuch der Pathogenen-microorganismen*, "that the filtrates of dysentery cultures are toxic." More recently, however, in a paper published early this year, Conradi³ states that he was able by a process of "aseptic autolysis," to extract from the dysentery bacillus a soluble toxin, and shortly after this Neisser and Shiga⁴ were able to obtain a similar toxin by suspending the bacilli in saline, killing them by exposure to heat, keeping the suspension for two days at 37° C., and then filtering. Recently Rosenthal⁵ has found that by growing the bacillus in Martin's alkaline broth highly toxic filtrates are obtained. This agrees with the results given above.