Published in final edited form as:

Tetrahedron Lett. 2008 ; 49(27): 4306–4309.

Coupling of Vinyl Aziridines and Isocyanates

Kainan Zhang, Pramod R. Chopade[†], and Janis Louie^{*}

University of Utah, Department of Chemistry, 315 South 1400 East, Salt Lake City, UT, 84112, USA

Abstract

Thermal coupling of vinyl aziridines and phenyl isocyanate was evaluated. Although oxazolidinone products were predominant, some reactions afforded a seven-membered ring heterocycle. When Ni/IMes was employed as a catalyst, a wider array of vinyl aziridines underwent coupling reactions. The Ni catalyzed reactions generally afforded vinyl imidazolidinones as major products.

Keywords

phenyl isocyanate; vinyl aziridine; N-heterocyclic carbene; nickel; imidazolidinone; oxazolidinone; tetrahydro-[1,3]diazepin-2-one

The coupling of an isocyanate and a vinylaziridine is an effective way to access heterocyclic ring systems. Although one example of a thermal coupling has been reported,¹ these reactions are typically catalyzed by Lewis acids such as sodium iodide² and nickel iodide.³ These couplings generally afford vinyl-substituted five membered imidazolidinone products. Palladium catalysts have also been used to catalyze this coupling reaction and in some cases, these couplings display high enantioselectivity.^{4,5,6} In each of these cycloadditions, the vinyl group acts to enhance the reactivity of the aziridine. For example, in non-metal mediated reactions the vinyl group is an activating group. In Pd catalyzed coupling reactions, the vinyl group enhances binding of the aziridine substrate. However, in all of these cases, incorporation of the vinyl group *in* the heterocycle is not observed.

We present here a study detailing both thermal reactions and nickel catalyzed reactions of phenyl isocyanate and vinyl aziridines. These reactions afford interesting heterocyclic products in addition to the expected vinyl substituted imidazolidinone.

(1)

^{*} Corresponding author. Tel.: 801-581-7309; fax: 801-581-8433; e-mail: louie@chem.utah.edu.

[†] Present Address: Coker College

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Despite the lack of information regarding simple thermal coupling of isocyanates and vinyl aziridines, we found that N-benzyl-2-styryl-aziridine (**1a**) reacted readily with 3 equiv of phenyl isocyanate at elevated temperatures (equation 1). At 100 °C, the coupling proceeded smoothly and afforded 3 distinct heterocyclic products, a seven-membered product (**2a**), an oxazolidinone **3a**,⁷ and a trace amount of imidazolidinone **4a**⁷ (equation 1).

(2)

A variety of vinyl aziridines were evaluated in the thermal coupling reaction (equation 2, Table 1).⁸ Electronic factors did not have much of an effect on the reaction as both para methoxy and para CF₃ substituted substrates afforded relatively equal ratios of heterocyclic products **2** and **3** (entries 2 and 3). The reaction with **1b** also afforded a vinyl substituted imidazolidinone (**4b**) in trace amounts. Additional substitution on the olefin led to the selective formation of the oxazolidine product **3** (entries 5 and 6). In entry 6 where the substrate lacks an activating phenyl group (**1f**), the reaction was quite sluggish. Only 38% of the **3f** was isolated. When the N-benzyl group was replaced by an N-cyclohexyl group, the seven-membered ring (**2g**) was slightly favored over the oxazolidinone ring (**3g**). Other substitution patterns lead to no heterocyclic products. Terminal vinyl aziridines possessing no activating group (i.e., R=Bn, R₁,R₂=H,H, **1h** and R=Cy, R₁,R₂=H,H, **1i**) did not react. In addition, when N was substituted with a bulky *tert*-butyl group (i.e., R=*t*-Bu, R₁,R₂=H,Ph, **1j**) or when the phenyl group was replaced with an ester group (i.e., R=Bn, R₁,R₂=H,CO₂Et, **1k**), no reaction took place. N-phenyl-vinylaziridines underwent a Ni catalyzed [3,3] sigmatropic rearrangement.⁹

(3)

(4)

In an effort to differentiate the reactivity of cis and trans vinyl aziridines, the cis and trans isomers of **1b** were evaluated individually in the thermal rearrangement reaction (equations 3 and 4). Interestingly, oxazolidinone products were observed in both reactions. However, the seven membered ring product (**2b**) was only observed in the reaction of the trans vinyl aziridine.

(5)

We have recently found that the combination of Ni(COD)₂ and NHC ligands catalyze the reaction of vinyl aziridines and alkynes.¹⁰ We surmised that this catalyst combination may facilitate the coupling of a wider range of vinyl aziridines and isocyanates and in a selective manner. As such, a series of NHCs were evaluated as potential ligands (Figure 1) for the Ni (0) catalyzed cycloaddition of isocyanates and vinylaziridines. Phenyl isocyanate (0.3M)¹¹ and N-benzyl-2-styryl-aziridine (**1a**, 0.1M) were again chosen as model substrates in a reaction catalyzed by 5 mol% Ni(COD)₂ and 10 mol% NHC in toluene at 100 °C (equation 5). Again, 3 heterocyclic products were generally observed. However, unlike the thermal reactions described above, **4a** was formed as the major product in almost all Ni/NHC catalyzed reactions. The use of IPr or SIPr afforded **2a**, **3a**, and **4a** in relatively equal amounts. Vinyl imidazolidinone **4a** was the major product when either IMes or IrBu were employed; however, yields and ratios of **4a** were higher when IMes was used as the ligand.

When an unactivated vinyl aziridine (**1h**) was used as a model substrate in lieu of **1a**, only the formation of imidazolidinone **4h** was observed (equation 5). The products obtained from thermal reactions, such as the seven-membered product (**2h**) and the oxazolidinone (**3h**), were not observed in appreciable amounts.¹² Although the combination of Ni/IMes did catalyze the coupling of **1h** and PhNCO (43% NMR yield), higher yields of **4h** was formed in the presence of IrBu (83% NMR yield). Similar results were obtained when N-cyclohexyl-vinylaziridine (**1i**) was used a model substrate.

The Ni/IMes catalyzed coupling of vinyl aziridines and phenyl isocyanate proved to be a more general reaction than thermal reactions (Table 2).¹³ Activated vinyl aziridines reacted smoothly with PhNCO (entries 1-5). In most cases, an imidazolidinone (**4**) was formed preferentially. Only the reaction of **1b** and **1e** gave slightly more seven-membered ring **2b** or oxazolidinone **3e**, respectively (entries 2 and 5). In contrast to the thermal reactions shown in Table 1, unactivated vinyl aziridines were also converted to heterocyclic products. For example, the thermal reaction of **1f** was sluggish (Table 1, 38% **3f**) whereas it afforded heterocyclic products (**3f** and **4f**) in an overall 64-67% yield in the Ni/NHC catalyzed reaction (entries 6a-b). In addition, terminal vinyl aziridines gave imidazolidinone products exclusively and in generally good yields (40% and 60%, entries 8a-9a respectively). Higher yields were observed with the Ni/IrBu catalysts (70% and 92%, entries 8b and 9b respectively). Other vinyl aziridines such as **1j** and **1k**, which were unreactive under thermal conditions, did undergo Ni/IMes catalyzed coupling. Interestingly, the steric hinderance of **1j** resulted in the formation of small amounts of a unique heterocycle (**5j**). It appears that with this substrate, C-C bond cleavage competes with the more favorable C-N bond cleavage of the aziridine (entry 10). N-Phenyl substituted vinyl aziridines afforded low amounts of imidazolidinone products in addition to aza-cope products (entries 12-13).

(6)

The amount of catalyst had a profound effect on the ratio of heterocyclic products in these reactions. Specifically, we found that increasing the catalyst loading resulted in the increase of the generation of the imidazolidinone derivative **4a**. On the other hand, reducing the catalyst promoted the formation of the 7 membered ring product **2a** (Table 3). It may be possible that the Ni/IMes catalyst serves to enhance the formation of the more thermodynamically favored imidazolidinone products (**4**). A similar phenomenon has been observed in nickel iodide promoted couplings.³

Given the results in Table 3, it was unclear whether **4a** was formed from a rearrangement of **2a** and **3a** or whether it was formed directly from **1a** and PhNCO. In addition, it may be possible for **2a** and **3a** to revert back to PhNCO and vinyl aziridine before ultimately affording **4a**. To address these possibilities, **1a** and PhNCO were heated to 100 °C until a mixture of **2a** and **3a** were obtained. Two equivalents of **1b** and catalytic amounts of Ni(COD)₂ and IMes were then added to the solution and the reaction was further heated to 100 °C. Interestingly, **4a_{trans}** was the only observed product (Scheme 1). No products (i.e., **4b**) arising from the reaction of **1b** and PhNCO were observed. Similarly, no products (i.e., **4n**) arising from the reaction of **1a** and CyNCO were observed when CyNCO was added in lieu of **1b** (Scheme 1). Thus, formation of heterocyclic products (**2-4**) appears to be irreversible in both thermal and nickel catalyzed reactions.

In summary, thermal and Ni/IMes-catalyzed reactions of phenyl isocyanate and various vinyl aziridines were evaluated. Thermal reactions generally afford oxazolidinone products. In some cases, a 7-membered heterocycle is also formed. Ni catalyzed reactions generally afforded the more thermodynamically favored imidazolidinone product.

Acknowledgements

We gratefully acknowledge the University of Utah, Merck, NSF (Career Award), the Camille-Dreyfus Foundation, and the NIGMS for support of this research.

References

1. For the thermal reaction of an isocyanate and an aziridine, see: Pfeil E, Milzner K. *Angew Chem, Int Engl Ed* 1966;5:667.
2. Nadir UK, Basu N. *Tetrahedron Lett* 1992;33:7949.
3. Munegumi T, Azumaya I, Kato T, Masu H, Saito S. *Org Lett* 2006;8:379. [PubMed: 16435839]
4. Alper H. *J Am Chem Soc* 1995;117:4700.
5. Trost BM, Fandrick DR. *J Am Chem Soc* 2003;125:11836. [PubMed: 14505403]
6. Dong C, Alper H. *Tetrahedron: Asymmetry* 2004;15:1537.
7. Bulter CD, Inman AG, Alper H. *J Org Chem* 2000;65:5887. [PubMed: 10987917]
8. General procedure for the thermal reactions: A toluene solution of PhNCO (75.9 mg, 0.637 mmol, 0.3 M) and vinylaziridine **1a** (50 mg, 0.212 mmol, 0.1 M) was heated to 100 °C for 8 hours. The reaction was cooled and the solvent was removed in vacuo. The residue was purified by flash chromatography (2:3 diethyl ether: pentane) to afford imidazolidinone **4a** (6.8 mg, 9%) and the 7 membered ring **2a** (30.9mg, 41%), and oxazolidinone **3a** (29.4 mg, 39%). **1-Benzyl-3,4-diphenyl-1,3,4,7-tetrahydro-[1,3]diazepin-2-one (2a)** Light yellow oil. IR (dissolved in methylene chloride) v: 1656, 1464, 1240 cm^{-1} . ^1H NMR (300 MHz, CDCl_3 , ppm) δ : 7.42-6.94 (m, 13H), 6.96 (q, 2H), 6.11 (m, 1H), 5.86 (m, 1H), 5.40 (d, 5.7Hz, 1H), 4.632 (d, 15Hz, 1H), 4.17-4.14 (m, 2H), 3.46 (dd, 1H). ^{13}C NMR (300 MHz, CDCl_3 , ppm) δ : 47.5, 53.6, 65.0, 125.3, 125.3, 125.9, 127.3, 127.4, 127.6, 128.5, 128.7, 128.9, 129.1, 130.1, 138.0, 141.9, 146.4, 163.9. EIMS m/z : 354 (M^+), 220, 206, 144, 115, 91. HRMS(ESI/APCI) for $\text{C}_{24}\text{H}_{23}\text{N}_2\text{O}$ [$\text{M}+\text{H}$] $^+$: calcd. 355.1810, found 355.1807. **(3-Benzyl-5-styryl-oxazolidin-2-ylidene)-phenyl-amine (3a_{trans})** Light yellow oil. IR (dissolved in methylene chloride) v: 1672, 1591 cm^{-1} . ^1H NMR (300 MHz, CDCl_3 , ppm) δ : 7.42-7.17 (m, 14H), 6.97 (t, 1H), 6.65 (d, 15.9Hz, 1H), 6.18 (q, 1H), 4.68 (d, 15Hz, 1H), 4.55 (d, 15Hz, 1H), 3.56 (t, 1H), 3.19 (dd, 1H). ^{13}C NMR (300 MHz, CDCl_3 , ppm) δ : 49.5, 50.9, 77.4, 122.3, 123.8, 125.4, 127.0, 127.9, 128.6, 128.7, 128.8, 128.9, 134.8, 135.8, 147.8, 152.8. EIMS m/z : 354 (M^+), 250, 206, 91, 28. HRMS(ESI/APCI) for $\text{C}_{24}\text{H}_{23}\text{N}_2\text{O}$ [$\text{M}+\text{H}$] $^+$: calcd. 355.1810, found 355.1802. **(3-Benzyl-5-styryl-oxazolidin-2-ylidene)-phenyl-amine (3a_{cis})** Light yellow oil. IR (dissolved in methylene chloride) v: 1675, 1591, 1493 cm^{-1} . ^1H NMR (300 MHz, CDCl_3 , ppm) δ : 7.40-7.14 (m, 13H), 6.96 (t, 1H), 6.75 (d, 11.4Hz, 1H), 5.76 (dd, 1H), 5.31 (q, 1H), 4.66 (d, 15Hz, 1H), 4.54 (d, 15Hz, 1H), 3.54 (t, 1H), 3.17 (t, 1H). ^{13}C NMR (400 MHz, CDCl_3 , ppm) δ : 49.5, 51.3, 72.7, 122.1, 123.8, 127.8, 128.1, 128.6, 128.7, 128.9, 135.5, 135.7, 137.0, 147.8, 152.8. EIMS m/z : 354 (M^+), 250, 206, 91, 28. HRMS(ESI/APCI) for $\text{C}_{24}\text{H}_{23}\text{N}_2\text{O}$ [$\text{M}+\text{H}$] $^+$: calcd. 355.1810, found 355.1811.
9. (a) Scheiner P. *J Org Chem* 1967;32:2628. (b) Figeys HP. *Tetrahedron Lett* 1981;22:637.
10. (a) Zuo G, Louie J. Unpublished results (b) Wolfe JP, Ney JE. *Org Lett* 2003;5:4607. [PubMed: 14627395] For Pd-catalyzed conversion of aziridines to imines, see
11. NHC catalyze the cyclotrimerization of isocyanates. (See Duong HA, Cross MJ, Louie J. *Org Lett* 2004;6:4679. [PubMed: 15575659]). Thus, excess isocyanate provided optimal reaction conditions.
12. No reaction occurred when couplings were run in the presence of $\text{Ni}(\text{COD})_2$ and NHC alone.
13. General procedure for $\text{Ni}(\text{COD})_2$ / NHC catalyzed reaction: In glovebox, a scintillation vial equipped with a magnetic stir bar was charged with PhNCO (75.9 mg, 0.637mmol, 0.3 M) and vinylaziridine **1a** (50 mg, 0.212 mmol, 0.1 M) in toluene. A solution of $\text{Ni}(\text{COD})_2$ (2.9mg, 0.0106 mmol, 0.005M) and IMes (6.5 mg, 0.0212 mmol, 0.01M) in toluene was added. The reaction was stirred at 100 °C for 6 hours. The reaction was cooled and the solvent was removed in vacuo. The residue was purified by flash chromatography (2:3 diethyl ether : pentane) to afford imidazolidinone **4a** (41.4 mg, 55%)

and the 7 membered ring **2a** (3.77mg, 5%). **1-Benzyl-3-phenyl-4-styryl-imidazolidin-2-one (4a_{trans})** Light yellow oil. IR (dissolved in methylene chloride) ν : 1701, 1499, 1423 cm^{-1} . ^1H NMR (300 MHz, CDCl_3 , ppm) δ : 7.56-7.02 (m, 16H), 6.59 (d, 15.9Hz, 1H), 6.13 (q, 1H), 4.80 (q, 1H), 4.51 (dd, 15Hz, 2H), 3.60 (t, 1H), 3.13 (dd, 1H). ^{13}C NMR (300 MHz, CDCl_3 , ppm) δ : 48.3, 48.8, 56.5, 120.8, 123.5, 126.8, 127.8, 128.0, 128.4, 128.5, 128.8, 128.9, 133.5, 136.1, 137.0, 139.5, 158.2. EIMS m/z : 354 (M^+), 263, 250, 223, 206, 91. HRMS(ESI/APCI) for $\text{C}_{24}\text{H}_{23}\text{N}_2\text{O}$ $[\text{M}+\text{H}]^+$: calcd. 355.1810, found 355.1806.

Figure 1.
NHC Ligands

Scheme 1.
NMR analysis of Ni catalyzed rearrangement of thermal coupling products.

Table 1

Thermal coupling of PhNCO and Vinyl aziridines.^a

Entry	1	R	Vinyl Aziridine	R ₁ , R ₂	2	3 (% Yield)	4
1	1a ^b		Bn	H, Ph	2a (41) ^c	3a (39) ^c	4a (9) ^c
2	1b ^b		Bn	H, 4-C ₆ H ₄ -OMe	2b (42) ^c	3b (36) ^c	4b (3) ^c
3	1c ^b		Bn	H, 4-C ₆ H ₄ -CF ₃	2c (25) ^d	3c (28) ^d	
4	1d ^b		Bn	H, -C ₂ H ₅	2d (42) ^d	3d (40) ^d	
5	1e ^b		Bn	Me, Ph		3e (74) ^c	
6	1f ^b		Bn	Me, Me		3f (38) ^c	
7	1g ^b		Cy	H, Ph	2g (36) ^c	3g (20) ^c	

^a Reaction conditions: 0.1M vinyl aziridine, 0.3M isocyanate, toluene, 100 °C.

^b mixture of cis and trans.

^c Isolated Yields.

^d GC or NMR yields determined relative to an internal standard.

Table 2

Nickel Catalyzed Coupling of PhNCO and Vinyl aziridines.

Entry	1	R	Vinyl Aziridine	R ₁ , R ₂	2	3	(% Yield)	4	5
1	1a ^c	Bn	Bn	H, Ph	2a (5) ^{a,d}	3b (11) ^{a,d}		4a (55) ^{a,d}	
2	1b ^c	Bn	Bn	H, 4-C ₆ H ₄ -OMe	2b (30) ^{a,d}			4b (24) ^{a,d}	
3	1c ^c	Bn	Bn	H, 4-C ₆ H ₄ -CF ₃	2c (6) ^e			4c (69) ^e	
4	1d ^c	Bn	Bn	H, -C ₂ H ₅	2d (13) ^e	3d (13) ^e		4d (41) ^e	
5	1e ^c	Bn	Bn	Me, Ph		3e (60) ^{a,d}		4e (18) ^{a,d}	
6a	1f	Bn	Bn	Me, Me		3f (27) ^{a,d}		4f (37) ^{a,d}	
6b						3f (30) ^{b,d}		4f (37) ^{b,d}	
7	1g ^c	Cy	Cy	H, Ph				4g (81) ^{a,d}	
8a	1h	Bn	Bn	H, H				4h (40) ^{a,d}	
8b								4h (70) ^{b,d}	
9a	1i	Cy	Cy	H, H				4i (60) ^{a,d}	
9b								4i (92) ^{b,d}	
10	1j ^c	<i>t</i> -Bu	<i>t</i> -Bu	H, Ph	2j (7) ^{a,d}			4j (16) ^{a,d}	5j (15) ^{a,d}
11	1k ^c	Bn	Bn	H, CO ₂ Et				4k (65) ^{a,d}	
12	1l	Ph	Ph	H, H				4l (13) ^{a,d}	
13	1m ^c	Ph	Ph	H, Ph				4m (35) ^{a,d}	

^a Reaction conditions: 5 mol% Ni(COD)₂, 10 mol% IMes, 0.1M vinyl aziridine, 0.3M isocyanate, toluene, 100 °C.

^b Reaction conditions: 5 mol% Ni(COD)₂, 10 mol% IrBu, 0.1M vinyl aziridine, 0.3M isocyanate, toluene, 100 °C.

^c mixture of cis and trans.

^d Isolated Yields.

^e GC yields determined relative to an internal standard.

Table 3

Ni/IMes catalyzed coupling of PhNCO and **1a**.^a

Entry	% Ni(COD) ₂	% IMes	2a	Ratio of Products ^b 3a	4a
1	10	20	1	0	24
2	5	10	1	0	11
3	1	2	1	0	1.5
4	0.1	0.2	1.6	0	1

^aReaction conditions: 0.1M **1a**, 0.3M PhNCO, toluene.

^bDetermined by GC using naphthalene as an internal standard.