Published in final edited form as:

J Med Chem. 2006 October 19; 49(21): 6254–6263.

Conformationally-Restricted Dipeptide Amides as Potent and Selective Neuronal Nitric Oxide Synthase Inhibitors

Haitao Ji[§], José A. Gómez-Vidal^{§,†}, Pavel Martásek^{¶,‡}, Linda J. Roman^{¶,±}, and Richard B. Silverman^{*,§}

Department of Chemistry, Department of Biochemistry, Molecular Biology, and Cell Biology, and Center for Drug Discovery and Chemical Biology, Northwestern University, Evanston, Illinois 60208 –3113 Department of Biochemistry, University of Texas Health Science Center, San Antonio, Texas 78284–7760

Abstract

Four new conformationally-restricted analogues of the potent and selective neuronal nitric oxide synthase inhibitor, *L*-nitroargininyl-*L*-2,4-diaminobutyramide (**1**), have been synthesized. *N*^α-Methyl and *N*^α-benzyl derivatives (**3** and **4**, respectively) of 4*N*-(*L*-Arg^{NO2})-*trans*-4-amino-*L*-prolineamide (**2**) are also selective inhibitors, but the potency and selectivity of **3** are weak. Analogue **4** has only one-third the potency and one-half to one-third the selectivity of **2** against iNOS and eNOS, respectively. 3-*N*-(*L*-Arg^{NO2})-*trans*-3-amino-*L*-prolineamide (**6**) is as potent an inhibitor of nNOS as is **2**; selectivity for nNOS over iNOS is half of that for **2** but the selectivity for nNOS over eNOS is almost double that for **2**. The corresponding *cis*-isomer (**5**) is a weak inhibitor of nNOS. These results are supported by computer modeling.

Nitric oxide (NO) is a ubiquitous biological messenger involved in a variety of physiological processes that acts as a signal transducer but also exerts a variety of regulatory and cytostatic functions. In most instances NO mediates its biological effects by activating guanylate cyclase and increasing cyclic GMP synthesis. However, effects of nitric oxide that are independent of cyclic GMP are also known.¹

Nitric oxide synthase (NOS, EC 1.14.13.39) isoforms² are homodimers that catalyze the oxidation of *L*-arginine to *L*-citrulline and nitric oxide in a NADPH- and O₂-dependent process.³ There are at least three distinct isoforms of NOS. The constitutive endothelial isoform (eNOS) is involved in the regulation of smooth muscle relaxation and blood pressure and in the inhibition of platelet aggregation. A second constitutive isoform is the neuronal NOS (nNOS), which is important for neurotransmission. A third isozyme is the inducible NOS (iNOS), which is located in activated macrophage cells and acts as a cytotoxic agent in normal immune responses. All of the isoforms utilize NADPH, FAD, FMN, tetrahydrobiopterin, and heme as cofactors. The constitutive isoforms also require Ca²⁺ and calmodulin for activity, while the inducible isoform has tightly bound Ca²⁺ and calmodulin. The crystal structures of the oxygenase domains of murine iNOS monomer,⁴ murine and human iNOS dimer,^{5,6,7,8} human and bovine endothelial NOS dimers,⁹ and rat neuronal NOS dimer¹⁰ indicate a high degree

*Address correspondence to this author at the Department of Chemistry Phone: 1–847–491–5653 Email:

Agman@chem.northwestern.edu.

[§]Northwestern University

[¶]University of Texas

[†]Present address: Departamento de Química Farmacéutica y Orgánica, Facultad de Farmacia, Universidad de Granada, 18071 Granada, Spain

[#]Developed the eNOS overexpression system in *E. coli* and the purification of eNOS.

[±]Developed the overexpression system for nNOS in *E. coli* and the purification of the nNOS.

of structural similarity within the catalytic center and the dimer interface regions between NOS isoforms. They share only approximately 50% of primary sequence homology, suggesting that they may differ from each other in regulatory aspects.

The use of NOS inhibitors in pathologically elevated synthesis of NO has great therapeutic potential.¹¹ NO overproduction by nNOS has been associated with neurodegeneration during stroke, spinal transmission of pain, migraine headaches, Parkinson's disease, and Alzheimer's disease. Thus nNOS represents an important therapeutic target for inhibitors.¹² Selective inhibition of one isoform over the others is essential because the three isoforms of NOS have unique roles in separate tissues. Selective inhibitors may represent useful tools for investigating other biological functions of NO.¹³

Previously, we synthesized and evaluated nitroarginine-containing dipeptide esters and dipeptide amides and identified a family of compounds that had high potency and selectivity for inhibition of nNOS over both iNOS and eNOS.^{14,15} The most potent nNOS inhibitor among these compounds was *L*-Arg^{NO₂}-*L*-Dbu-NH₂ (**1**, $K_i = 130$ nM); this compound exhibited high selectivity over eNOS (>1500-fold) and iNOS (192-fold). More recently we reported **2** (nNOS $K_i = 103$ nM) as a conformationally restricted analogue of *L*-Arg^{NO₂}-*L*-Dbu-NH₂ (**1**).¹⁶ This compound was a little more potent than **1** with comparable selectivity for nNOS over eNOS and iNOS. Here we report the synthesis and *in vitro* evaluation of four new analogues of **2**, two that are *N*^α-substituted analogues (**3** and **4**) to increase lipophilicity and determine if additional binding could enhance potency and selectivity, and two that are positional isomers of the pyrrolidine ring of **2** (**5** and **6**) to determine the optimal binding orientation.

Chemistry

The solution-phase syntheses of **3** and **4** are shown in Scheme 1. Starting from commercially-available *N*-Boc-*trans*-4-*N*-Fmoc-amino-*L*-proline, only six steps were required to give the final product. All of the synthetic steps, except for the dipeptide coupling reaction, afforded quantitative yields of products. In the reductive amination reaction the starting material, *trans*-4-*N*-Fmoc-amino-*L*-proline-NH₂, has good solubility in MeOH, but the products, *N*-substituted-*trans*-4-*N*-Fmoc-amino-*L*-proline-NH₂, have poor solubility in MeOH, so they precipitated from the organic phase during the reaction. In the dipeptide coupling reaction the mixed anhydride method (isobutyl chloroformate [IBCF]/*N*-methylmorpholine [NMM]) provided a much higher yield (88%) than the active ester method (HBTU/HOBt/DIEA; 40%), especially when the starting material contained an amide group.

A solution-phase synthesis of **6**, shown in Scheme 2, started from commercially-available *trans*-3-hydroxy-*L*-proline. In the Mitsunobu reaction for the conversion of **16** to **17** the use of *p*-nitrobenzoic acid as the nucleophile afforded pure product in a 61% yield, whereas with acetic acid as the nucleophile none of the product was isolated. The hydrolysis of **17** using

sodium azide as the catalyst provided a quantitative yield of pure **18**.¹⁷ A Mitsunobu reaction was also attempted to convert alcohol **18** to amine **19**. Two different nucleophiles, diphenyl phosphoryl azide (DPPA) and phthalimide were tried, but no product was produced. That might be because the hydroxyl group and the amide group in *cis*-**18** formed a strong intramolecular hydrogen bond, which prevented the triphenyl phosphine from reaching the alcohol. When the amide was changed to a methyl ester, the Mitsunobu reaction proceeded smoothly, as was demonstrated previously.¹⁸ Alcohol **18** was activated with methanesulfonyl chloride to form the mesylate, which was converted to the azide with sodium azide in a 62% yield.

Compounds **3–6** were initially synthesized using a solid-phase synthetic method (Scheme 3 and Supporting Information Scheme 1), but **3**, **4**, and **6** were optimized by the solution-phase methods shown in Schemes 1 and 2. The solid-phase synthesis of 3-aminoproline-containing dipeptides **5** and **6** started from *N*-Boc protected *cis*- and *trans*-3-azido-*L*-proline (**22** and **23**), which were previously reported.¹⁸ Triethylammonium tris(phenylthio)stannate has been described as a very efficient reducing agent with high chemoselectivity toward azides.¹⁹ This reagent has been applied successfully on solid phase using Rink resin²⁰ or other resins²¹ and afforded amine **25** in high purity.

Results and Discussion

Compound **2** was previously found to be the most potent conformationally-restricted analogue of *L*-Arg^{NO₂}-*L*-Dbu-NH₂ (**1**), which is a highly selective inhibitor of nNOS over eNOS.¹⁵ *D*-Arg^{NO₂}-*D*-Phe-OMe is a highly selective inhibitor of nNOS over iNOS.¹⁴ Clearly, these two compounds interact differently with the NOS isozymes, as previously described.¹⁰ *N*-Substituted analogues of **2** (**3** and **4**) were synthesized to determine if having both an amino group and a hydrophobic group would enhance binding for nNOS over the other two isoforms even more selectively than one or the other. Compounds **5** and **6** (the *cis*- and *trans*-3-amino-*L*-prolinamide derivatives) also were synthesized to determine if it was better to attach the *L*-Arg^{NO₂} at the 3-position or 4-position (**2**) of the proline ring to maximize potency and selectivity for nNOS.

These compounds were assayed against all three isozymes of NOS for potency and selectivity (Table 1). When compared to non-substituted analogue **2**, the *N*-substituted compounds (**3** and **4**) showed reduced nNOS potency (82- and 3-fold, respectively) and selectivity (73- and 44-fold for **3** against iNOS and eNOS, respectively; 2- and 3-fold for **4** against iNOS and eNOS, respectively). These results support the importance of a non-substituted pyrrolidine amino group for interaction with nNOS. Compound **4** showed a two-fold decrease in potency against nNOS when compared to **1**. On the other hand, **4** was six-fold more potent than the previously reported phenylalanine-containing dipeptide (**27**) against nNOS.¹⁴ The sequence alignment and a schematic drawing of the active site of NOS were presented in an earlier paper.²² The crystal structure of **2** bound in the active site of nNOS and eNOS^{10a} shows that the pyrrolidine nitrogen is located between the heme propionate group and the carbonyl group of the tetrahydrobiopterin cofactor. Compound **27** does not have a nitrogen at that position. The loss of this charge-charge interaction between the positively charged nitrogen atom and the negatively charged heme propionate appears to be the main reason for the low nNOS potency of **27** compared to that of **2** or **6**.

Further analysis of the crystal structure of **2** bound to rat nNOS shows that the pyrrolidine nitrogen of **2**, as a secondary amine, forms a H-bond network interaction with R596, the heme propionate, and the carbonyl group of the tetrahydrobiopterin cofactor via three structural water molecules (Figure 1). Alkylation of this nitrogen atom would exclude these structural water molecules from the active site. An AutoDock calculation of **4** showed that its phenyl group has the same spatial orientation in the C1 pocket²² of nNOS as that of **27**. The superimposition of

the docked conformation of **4** and the crystallographic binding conformation of **27** with nNOS (PDB ID 1rs7) show, however, that the phenyl group of **27** binds more tightly to hydrophobic residues L337 and M336 of rat nNOS than does the phenyl group of **4** (Figure 2). The sum of the loss of the H-bond network interaction mentioned above and the partial compensation from the interaction of the benzyl group of **4** with the hydrophobic residues in the C1 pocket account for the low potency of **4** compared to **2**. Compound **3** is particularly poor because the methyl group interferes with the above H-bond network interactions and is not long enough to extend into the C1 pocket for a hydrophobic interaction (AutoDock data not shown). As described before,^{10a,22} the charge-charge interaction between the α -amino group in our dipeptide nNOS inhibitors and D597 of rat nNOS accounts for the selectivity of these inhibitors for nNOS over eNOS. The stronger the interaction that the side chain of the dipeptide inhibitors has with the active site, the more prone is the α -amino group to fix at the selective region defined by nNOS D597/eNOS N368.²² That is most likely the reason why **2** is more selective than **4** for binding to nNOS over eNOS.

The *cis*-3-amino-*L*-proline derivative (**5**) displayed very low potency against nNOS ($IC_{50} \approx 100 \mu M$). However, the *trans*-3-aminoprolinamide isomer (**6**) showed nearly the same potency against nNOS as the previously described *trans*-4-aminoprolinamide isomer (**2**) but with almost a doubling in selectivity for eNOS (Table 1). Therefore, the *trans* pyrrolidine ring is the favored configuration for the interaction of these conformationally-restricted dipeptides with nNOS regardless of whether the *L*-Arg^{NO₂} is attached at the 3- or 4-position. Molecular modeling of **5** and **6** shows that the spatial orientation of the pyrrolidine ring of **6** is almost identical to that of **2**, except that the terminal amide groups protrude in different directions (Figure 3A). Since the terminal amide group of **2** only weakly interacts directly with the residues or cofactors in the active site, the nNOS potencies of these dipeptide inhibitors are primarily determined by the *L*-Arg^{NO₂} moiety and the pyrrolidine ring part of the proline. The superimposition of the docking conformation of **5** with the crystallographic binding conformation of **2** shows that the pyrrolidine ring of the *cis* compound (**5**) is out of the binding area of the heme propionate and the tetrahydrobiopterin (Figure 3B), which explains its low potency. The corresponding calculated binding free energies (Table 2) support the results of the above docking observations. All of the scoring functions arrived at the same conclusion, except for D_Score, that the binding of **5** should be less favorable than that for **6**. Others,²³ however, have found that D_Score gives the lowest correlation between calculated and experimental binding affinities.

In summary, a non-substituted proline nitrogen is necessary for high potency against nNOS. The *L*-Arg^{NO₂} moiety must be positioned as a *trans*-3- or *trans*-4-substituent on the proline ring for enhanced potency and selectivity against nNOS over eNOS and iNOS. Attachment at the 3-position favors nNOS over eNOS selectivity greater than at the 4-position but lower nNOS over iNOS selectivity than at the 4-position.

Experimental

General methods, reagents, and materials

All reagents were purchased from Aldrich, Novabiochem, or Advanced ChemTech and were used without further purification unless stated otherwise. *trans*-3-Hydroxy-*L*-proline was purchased from Acros Organics. NADPH, calmodulin, and human ferrous hemoglobin were obtained from Sigma Chemical Co. Tetrahydrobiopterin (H₄B) was purchased from Alexis Biochemicals. HEPES, DTT, and conventional organic solvents were purchased from Fisher Scientific. Tetrahydrofuran (THF) was distilled under nitrogen from sodium/benzophenone. Sodium sulfate was used to dry organic layers during work up. ¹H NMR spectra were recorded on a Gemini 2000, Mercury 400 or Inova 500 spectrometers (75.4, 100.6, or 125.7 MHz, for ¹³C NMR spectra) in CDCl₃. Chemical shifts are reported as δ values in parts per million

with the CDCl_3 (TMS), $(\text{CD}_2\text{H})_2\text{SO}$, CD_2HOD , and HOD resonance peaks set at 0 ppm, 2.50 ppm, 3.31 ppm, and 4.80 ppm respectively. E. Merck pre-coated silica gel 60 F254 plates were used for thin-layer chromatography (TLC), with visualization accomplished with the phosphomolybdic acid or ninhydrin spray reagent or with a UV-Vis lamp. Flash chromatography was performed with E. Merck silica gel 60 (230–400 mesh) or Sorbent Technologies 250 mesh silica gel. Fractions containing pure compound were concentrated *in vacuo* and the residue lyophilized. pH measurements were conducted with an Orion Research model 701 pH meter with a general combination electrode. Elemental analyses were performed by Atlantic Microlab Inc. Mass spectra were performed on a Micromass Quattro II triple quadrupole mass spectrometer (ESI). High-resolution mass spectra were carried out at the University of Illinois at Urbana-Champaign using a Micromass Quattro tandem quadrupole/hexapole/quadrupole mass spectrometer. NOS assays were recorded on a Perkin-Elmer Lambda 10 UV/VIS spectrophotometer.

***N*^α-Boc-*trans*-4-*N*-Fmoc-amino-*L*-proline- NH_2 (7)**—To a cooled solution ($-15\text{ }^\circ\text{C}$) of *N*-Boc-*trans*-4-*N*-Fmoc-amino-*L*-proline (0.451 g, 0.001 mol) in anhydrous THF (10 mL) was added *N*-methylmorpholine (0.111 g, 0.12 mL, 0.0011 mol) and isobutyl chloroformate (0.150 g, 0.144 mL, 0.0011 mol), and stirring continued for 30 min. NH_4OH (28% aqueous solution, 0.188 g, 0.21 mL, 0.0015 mol) was added, and the reaction mixture was stirred at $-15\text{ }^\circ\text{C}$ for 30 min and then warmed to room temperature over 1 h. The solvent was evaporated under reduced pressure. The residue was suspended in EtOAc (40 mL), washed successively with 1 M KHSO_4 ($2 \times 20\text{ mL}$), saturated NaCl (20 mL), 5% NaHCO_3 ($2 \times 20\text{ mL}$), and saturated NaCl (20 mL). The solution was dried over anhydrous Na_2SO_4 . The solvent was removed under reduced pressure giving a residue, which was purified by silica gel column chromatography (CH_2Cl_2 : MeOH = 9 : 1) to afford a white solid (0.45 g, quantitative yield). ^1H NMR (500 MHz, CDCl_3) δ (7.758 + 7.748) (d, 2H, $J = 7.5\text{ Hz}$), (7.569 + 7.554) (d, 2H, $J = 7.5\text{ Hz}$), 7.389(t, 2H, $J = 7.5\text{ Hz}$), 7.301(t, 2H, $J = 7.5\text{ Hz}$), 4.421(s, 2H), 4.383–4.095 (m, 3H), 3.856–3.653 (m, 1H), 3.408–3.008 (m, 1H), 2.523–2.427(m, 1H), 2.122–1.728(m, 1H), 1.448 (s, 9H). ^{13}C NMR (125.7 MHz, CDCl_3) δ (175.0 + 173.9) (1C), 155.9 (1C), (155.6 + 154.5) (1C), 143.9 (2C), 141.5 (2C), 127.9 (2C), 127.2 (2C), 125.1 (2C), 120.2 (2C), 81.2 (1C), 66.7 (1C), 58.3 (1C), 52.3 (1C), 50.1 (1C), 47.3 (1C), (37.1 + 34.0) (1C), 28.5 (3C). APCI (CH_2Cl_2) $m/z = 452.1$ ($[\text{M} + \text{H}]^+$).

***trans*-4-*N*-Fmoc-amino-*L*-proline- NH_2 (8)**—**7** (0.451 g, 0.001 mol) was dissolved in anhydrous CH_2Cl_2 (1 mL), and trifluoroacetic acid (TFA, 1 mL) was added at $0\text{ }^\circ\text{C}$. The reaction mixture was stirred 1 h at $0\text{ }^\circ\text{C}$ and 1 h at room temperature. After the reaction was completed, the solvent was evaporated under reduced pressure to afford a colorless oil (0.465 g, quantitative yield). ^1H NMR (500 MHz, CD_3OD) δ (7.657 + 7.642) (d, 2H, $J = 7.5\text{ Hz}$), (7.500 + 7.485) (d, 2H, $J = 7.5\text{ Hz}$), 7.249 (t, 2H, $J = 7.5\text{ Hz}$), 7.170 (t, 2H, $J = 7.5\text{ Hz}$), 4.300 (m, 3H), 4.139–4.095 (m, 2H), 3.469 (m, 1H), 3.179 (m, 1H), 2.317–2.306 (m, 1H), 2.166 (m, 1H). ^{13}C NMR (125.7 MHz, CDCl_3) δ (161.1 + 160.8) (1C), 158.3 (1C), 145.3 (2C), 142.8 (2C), 129.0 (2C), 128.3 (2C), 126.1 (2C), 121.1 (2C), 67.9 (1C), 60.3 (1C), 51.9 (1C), 51.5 (1C), 48.5 (1C), 36.7 (1C).

ESI(CH_3OH) $m/z = 352.3$ ($[\text{M} + \text{H}]^+$).

***N*^α-Methyl-*trans*-4-*N*-Fmoc-amino-*L*-proline- NH_2 (9)**—To a solution of **8** (0.465 g, 0.001 mol) dissolved in MeOH (20 mL) at $0\text{ }^\circ\text{C}$ was added a 37% formaldehyde solution (0.812 g, 0.745 mL, 0.01 mol), NaBH_3CN (0.314 g, 0.005 mol) and acetic acid (0.120 g, 0.115 mL, 0.002 mol). The reaction mixture was stirred at room temperature overnight. The solvent was evaporated under reduced pressure. The residue was dissolved in 5% NaHCO_3 (10 mL) and extracted with EtOAc (10 mL $\times$ 3). The combined organic layers were washed with brine and evaporated *in vacuo*. The residue was purified by silica gel column chromatography

(CH₂Cl₂ : MeOH = 9.5 : 0.5) to afford a white solid (0.365 g, quantitative yield). ¹H NMR (500 MHz, DMSO-*d*₆) δ (7.897 + 7.881) (d, 2H, J = 8Hz), (7.694 + 7.680) (d, 2H, J = 7Hz), 7.412 (t, 2H, J = 8Hz), 7.331 (t, 2H, J = 7.5Hz), (4.316 + 4.302) (d, 2H, J = 7Hz), 4.204 (t, 1H, J = 7Hz), 4.108–3.960 (m, 1H), 3.191 (t, 1H, J = 7Hz), 2.781 (t, 1H, J = 8Hz), 2.236 (s, 3H), 2.140 (t, 1H, J = 8.5Hz), 1.986–1.896 (m, 2H). ¹³C NMR (125.7 MHz, DMSO-*d*₆) δ 174.9 (1C), 155.7 (1C), 143.9 (2C), 140.8 (2C), 127.6 (2C), 127.1 (2C), 125.2 (2C), 120.2 (2C), 67.6 (1C), 65.2 (1C), 61.1 (1C), 48.9 (1C), 46.7 (1C), 40.9 (1C), 36.7 (1C). ESI (CH₃CN) *m/z* = 366.6([M + H]⁺).

N^α-Benzyl-*trans*-4-*N*-Fmoc-amino-*L*-proline-NH₂ (10)—To a solution of **8** (0.465 g, 0.001 mol) dissolved in MeOH (20 mL) at 0 °C was added benzaldehyde (0.531 g, 0.505 mL, 0.005 mol), NaBH₃CN (0.314 g, 0.005 mol) and acetic acid (0.120 g, 0.115 mL, 0.002 mol). The reaction mixture was stirred at room temperature overnight. The solvent was evaporated under reduced pressure. The residue was dissolved in 5% NaHCO₃ (10 mL) and extracted with EtOAc (10 mL × 3). The combined organic layers were washed with brine and evaporated *in vacuo*. The residue was purified by silica gel column chromatography (CH₂Cl₂ : MeOH = 9.5 : 0.5) to afford a white solid (0.441 g, quantitative yield). ¹H NMR (500 MHz, DMSO-*d*₆) δ (7.887 + 7.872) (d, 2H, J = 7.5Hz), (7.672 + 7.657) (d, 2H, J = 7.5Hz), 7.462–7.241 (m, 9H), (4.288 + 4.275) (d, 2H, J = 6.5Hz), 4.182 (t, 1H, J = 6.5Hz), 3.984–3.940 (m, 1H), 3.398–3.358 (m, 2H), 3.103 (t, 1H, J = 7.5Hz), 3.014 (t, 1H, J = 8Hz), 2.092 (t, 1H, J = 8.5Hz), 1.986–1.951 (m, 2H). ¹³C NMR (125.7 MHz, DMSO-*d*₆) δ 175.2 (1C), 155.7 (1C), 143.9 (2C), 140.8 (2C), 138.3 (1C), 128.8 (2C), 128.2 (2C), 127.6 (2C), 127.1 (3C), 125.1 (2C), 120.1 (2C), 67.7 (1C), 65.2 (1C), 58.2 (1C), 57.8 (1C), 48.9 (1C), 46.7 (1C), 36.7 (1C). ESI(CH₃OH) *m/z* = 442.6([M + H]⁺).

N^α-Methyl-*trans*-4-*N*-amino-*L*-proline-NH₂ dihydrochloride salt (11)—**9** (0.365 g, 0.001 mol) was treated with 20% piperidine (10 mL) in anhydrous DMF and stirred under a nitrogen atmosphere for 2 h. TLC monitored the complete conversion of the starting material to product. The reaction mixture was then concentrated under reduced pressure. The resulting residue was dissolved in a dilute HCl aqueous solution (four drops of concentrated HCl in 20 mL of H₂O) and washed with hexane to remove the piperidine-dibenzofulvene adduct and the free dibenzofulvene. The aqueous layer was concentrated *in vacuo* to yield the desired white solid (0.216 g, quantitative yield). ¹H NMR (500 MHz, D₂O) δ 3.921–3.861 (m, 1H), (3.557 + 3.542, 3.536 + 3.522) (dd, 1H, J = 7.5Hz, J = 7Hz), 3.340–3.306 (m, 1H), 2.566–2.529 (m, 1H), 2.405 (s, 3H), 2.368–2.236 (m, 2H). ¹³C NMR (125.7 MHz, D₂O) δ 176.4 (1C), 66.6 (1C), 58.7 (1C), 48.0 (1C), 39.7 (1C), 34.3 (1C). ESI (CH₃OH) *m/z* = 144.2([M + H]⁺).

N^α-Benzyl-*trans*-4-*N*-amino-*L*-proline-NH₂ dihydrochloride (12)—**10** (0.441 g, 0.001 mol) was treated with 20% piperidine (10 mL) in anhydrous DMF and stirred under a nitrogen atmosphere for 2 h. After complete conversion of the starting material to product (by TLC) the reaction mixture was concentrated under reduced pressure. The resulting residue was dissolved in the dilute aqueous HCl solution (four drops of concentrated HCl in 20 mL of H₂O) and washed with CH₂Cl₂ to remove the piperidine-dibenzofulvene adduct and the free dibenzofulvene. The aqueous layer was concentrated *in vacuo* to yield the desired white solid (0.292 g, quantitative yield). ¹H NMR (500 MHz, D₂O) δ 7.471–7.414 (m, 5H), 4.650–4.614 (m, 1H), (4.504 + 4.479) (d, 1H, J = 7.5Hz), (4.459 + 4.434) (d, 1H, J = 7.5Hz), 4.121–4.092 (m, 2H), 3.592–3.540 (m, 1H), 2.757–2.708 (m, 1H), 2.595–2.533 (m, 1H). ¹³C NMR (125.7 MHz, D₂O) δ 168.7 (1C), 131.1 (2C), 130.8 (1C), 129.5 (2C), 128.6 (1C), 64.6 (1C), 58.7 (1C), 55.6 (1C), 46.9 (1C), 33.1 (1C). ESI (CH₃OH) *m/z* = 220.2 ([M + H]⁺).

4-*N*-(N^α-Boc-*L*-Arg^{NO2})-N^α-methyl-*trans*-4-amino-*L*-proline-NH₂ (13)—**11** (0.324 g, 0.0015 mol) was dissolved in 0.5 M NaOH aqueous solution (10 mL) and then was extracted

with CH₂Cl₂ (10 mL × 2). The combined organic layers were washed with water (10 mL) to generate the free base form of **11**. A solution of *N*^α-Boc-*L*-Arg^{NO₂}-OH (0.319 g, 0.001 mol) in dry THF (20 mL) was cooled to −15 °C, and 4-methylmorpholine (0.111 g, 0.121 mL, 0.0011 mol) and isobutyl chloroformate (0.150 g, 0.144 mL, 0.0011 mol) were added. The reaction mixture was stirred for 30 min, and the free base form of **11** (0.157 mL, 0.0011 mol) in anhydrous THF (5 mL) was added via a cannula. The solution was stirred for 30 min at −15 °C and was then allowed to warm to room temperature over 30 min. The reaction mixture was stirred at room temperature for 1 h, and the solvent was evaporated *in vacuo*. The residue was purified directly by silica gel column chromatography (CH₂Cl₂ : MeOH = 8.5 : 1.5) to afford 0.39 g of a colorless oil (88%). ¹H NMR (500 MHz, CD₃OD) δ 4.364–4.336 (m, 1H), 4.080–4.015 (m, 1H), 3.407–3.376 (m, 1H), 3.311–3.271 (m, 2H), 3.036–3.003 (m, 1H), 2.380 (s, 3H), 2.311 (t, 1H, J = 8.5 Hz), 2.209–2.152 (m, 1H), 2.094–2.055 (m, 1H), 1.800–1.636 (m, 4H), 1.437 (s, 9H). ¹³C NMR (125.7 MHz, CD₃OD) δ 179.1 (1C), 174.8 (1C), 161.0 (1C), 158.0 (1C), 80.8 (1C), 68.9 (1C), 61.9 (1C), 55.6 (1C), 41.8 (1C), 41.5 (2C), 38.3 (1C), 30.8 (1C), 28.8 (3C), 26.2 (1C). ESI (CH₃CN) *m/z* = 445.4 ([M + H]⁺); *m/z* = 467.3 ([M + Na]⁺).

4-*N*-(*N*^α-Boc-*L*-Arg^{NO₂})-*N*^α-benzyl-*trans*-4-amino-*L*-proline-NH₂ (14**)—**12**** (0.438 g, 0.0015 mol) was dissolved in 0.5 M NaOH aqueous solution (10 mL) and then extracted with CH₂Cl₂ (10 mL × 2). The combined organic layers were washed with water (10 mL) to generate the free base form of **12**. A solution of *N*^α-Boc-*L*-Arg^{NO₂}-OH (0.319 g, 0.001 mol) in dry THF (20 mL) was cooled to −15 °C, and 4-methylmorpholine (0.111 g, 0.121 mL, 0.0011 mol) and isobutyl chloroformate (0.150 g, 0.144 mL, 0.0011 mol) were added. The reaction mixture was stirred for 30 min, and the free base form of **12** (0.241 g, 0.0011 mol) in anhydrous THF (5 mL) was added via a cannula. The solution was stirred for 30 min at −15 °C and then allowed to warm to room temperature over 30 min. The reaction mixture was stirred at room temperature for 1 h, and the solvent was evaporated *in vacuo*. The residue was suspended in EtOAc (20 mL) and washed successively with 5% NaHCO₃ (2 × 20 mL) and saturated NaCl (20 mL). The solution was dried over anhydrous MgSO₄, and the solvent was removed *in vacuo*. The residue was purified by silica gel column chromatography (CH₂Cl₂ : MeOH = 9 : 1) to afford 0.36 g of a colorless oil (70%). ¹H NMR (500 MHz, CD₃OD) δ 7.344–7.214 (m, 5H), 4.295–4.266 (m, 1H), 4.000–3.800 (m, 1H), (3.878 + 3.852) (d, 1H, J = 13 Hz), (3.521 + 3.495) (d, 1H, J = 13 Hz), 3.288–3.206 (m, 4H), 2.280 (t, 1H, J = 8.5 Hz), 2.177–2.123 (m, 1H), 2.087–2.026 (m, 1H), 1.696–1.525 (m, 4H), 1.400 (s, 9H). ¹³C NMR (125.7 MHz, CD₃OD) δ 179.8 (1C), 174.8 (1C), 161.1 (1C), 157.9 (1C), 139.2 (1C), 130.3 (2C), 129.6 (2C), 128.6 (1C), 80.8 (1C), 67.0 (1C), 60.2 (1C), 58.9 (1C), 55.6 (1C), 41.8 (2C), 37.8 (1C), 30.8 (1C), 28.8 (3C), 26.1 (1C). ESI (CH₃CN) *m/z* = 521.1 ([M + H]⁺); *m/z* = 543.1 ([M + Na]⁺).

4-*N*-(*L*-Arg^{NO₂})-*N*^α-methyl-*trans*-4-amino-*L*-proline-NH₂ ditrifluoroacetate (3**)—**13**** (0.222 g, 0.001 mol) was dissolved in anhydrous CH₂Cl₂ (1 mL), and TFA (1 mL) was added at 0 °C. The reaction mixture was stirred for 1 h at 0 °C and 1 h at room temperature. After the reaction was done, the solvent was evaporated *in vacuo*. The residue was partitioned between water (10 mL) and ethyl acetate (10 mL). The aqueous layer was then washed with ethyl acetate (2 × 10 mL). The combined aqueous layers were evaporated by high-vacuum rotary evaporation to afford a white foamy solid (0.286 g, quantitative yield). ¹H NMR (500 MHz, D₂O) δ 4.459–4.446 (m, 1H), 4.382–4.354 (m, 1H), 4.040–4.018 (m, 1H), 3.933–3.922 (m, 1H), 3.220–3.151 (m, 3H), 2.914 (s, 3H), 2.460 (m, 2H), 1.855 (m, 2H), 1.597 (m, 2H). ¹³C NMR (125.7 MHz, D₂O) δ 169.6 (1C), 169.4 (1C), (162.9 + 162.6) (2C, TFA-COOH), 158.9 (1C), (119.6 + 117.5 + 115.1 + 112.8) (2C, TFA-CF₃), 67.3 (1C), 59.0 (1C), 52.8 (1C), 47.4 (1C), 40.9 (1C), 40.3 (1C), 34.6 (1C), 27.9 (1C), 23.0 (1C). Comb. Anal. (C₁₂H₂₄N₈O₄·3TFA) C, H, N.

HPLC conditions for the purification of 4-*N*-(*L*-Arg^{NO2})-*N*^α-methyl-*trans*-4-amino-*L*-proline-NH₂ ditrifluoroacetate (3)—Column: Phenomenex Gemini 5μ C18 110A preparative column, column size is 250 × 21.2 mm 5μ micron; injection concentration was 20 mg/mL; flow rate: 12 mL/min; detection was at 254 nm; the mobile phase was H₂O (0.1% TFA) : 60% CH₃CN-40% H₂O (0.08% TFA) = 97 : 3, and an isocratic method was used. The retention time was 5 min.

4-*N*-(*L*-Arg^{NO2})-*N*^α-benzyl-*trans*-4-amino-*L*-proline-NH₂ ditrifluoroacetate (4)—**14** (0.260 g, 0.5 mmol) was dissolved in anhydrous CH₂Cl₂ (1 mL), and TFA (1 mL) was added at 0 °C. The reaction mixture was stirred for 1 h at 0 °C, and for 1 h at room temperature. After the reaction was completed, the solvent was evaporated *in vacuo*, and the residue was dissolved in water (10 mL). The color impurity was extracted with ethyl acetate (10 mL). The aqueous layer was then washed with ethyl acetate (2 × 10 mL), and the combined aqueous layers were evaporated by high-vacuum rotary evaporation to afford a white foamy solid (0.324 g, quantitative yield). ¹H NMR (500 MHz, D₂O) δ 7.279–7.242 (m, 5H), 4.329–4.183 (m, 4H), 3.807–3.772 (m, 2H), 3.151–3.074 (m, 3H), 2.338–2.215 (m, 2H), 1.725 (m, 2H), 1.466 (m, 2H). ¹³C NMR (125.7 MHz, D₂O) δ 169.5 (1C), 169.4 (1C), (162.8 + 162.5 + 162.3 + 162.0) (2C, TFA-COOH), 158.6 (1C), 130.8 (2C), 130.4 (1C), 129.3 (2C), 128.7 (1C), (119.5 + 117.2 + 114.9 + 112.6) (2C, TFA-CF₃), (64.8 + 64.7) (1C), 58.9 (1C), 57.1 (1C), (52.8 + 52.7) (1C), (47.14 + 47.09) (1C), 40.2 (1C), 34.3 (1C), 27.9 (1C), 22.9 (1C). ESI (CH₃OH) *m/z* = 421.3 ([M + H]⁺); *m/z* = 443.3 ([M + Na]⁺). HRMS (ES⁺, CH₃OH): Calc.: 421.2312; Found : 421.2319. Comb. Anal. (C₁₈H₂₈N₈O₄·3TFA) C, H, N.

HPLC conditions for the purification of 4-*N*-(*L*-Arg^{NO2})-*N*^α-benzyl-*trans*-4-amino-*L*-proline-NH₂ ditrifluoroacetate (4)—Column: a Phenomenex Gemini 5μ C18 110A preparative column, column size is 250 × 21.2 mm 5μ micron; injection concentration was 20 mg/mL; flow rate: 12 mL/min; detection was at 254 nm; the mobile phase was H₂O (0.1% TFA) : 60% CH₃CN-40% H₂O (0.08% TFA) = 70 : 30, and an isocratic method was used. The retention time was 10 min.

***N*^α-Boc-*trans*-3-hydroxy-*L*-proline (15)**—A solution of *trans*-3-hydroxy-*L*-proline (1.311 g, 0.01 mol) in a 2 : 1 mixture of THF/H₂O (20 mL) was treated first with 10% aqueous NaOH (4.5 mL) and then with di-*tert*-butyl dicarbonate (3.274 g, 0.015 mol). The reaction mixture was stirred at room temperature overnight, and THF was evaporated *in vacuo*. The residue was adjusted to pH = 2 by the addition of 10% aqueous KHSO₄. The acidic solution was extracted several times with ethyl acetate. The combined organic extracts were washed with H₂O and brine and then dried over anhydrous Na₂SO₄. Removal of the desiccant and evaporation of the solvent *in vacuo* gave the product as a solid, which was purified by silica gel column chromatography (EtOAc : MeOH : AcOH = 200 : 0.5 : 0.5) to afford a white solid (2.31 g, quantitative yield). ¹H NMR (500 MHz, CD₃OD) δ 4.371–4.330 (m, 1H), 4.181–4.029 (m, 1H), 3.542–3.439 (m, 2H), 2.002–1.968 (m, 1H), 1.862 (m, 1H), 1.458–1.333 (m, 9H). ¹³C NMR (125.7 MHz, CD₃OD) δ (174.2 + 173.9) (1C), (156.3 + 155.9) (1C), (81.5 + 81.3) (1C), (75.7 + 74.9) (1C), (69.3 + 68.9) (1C), (45.8 + 45.3) (1C), (33.4 + 32.8) (1C), (28.9 + 28.7) (1C). ESI (CH₃OH) *m/z* = 232.1 ([M + H]⁺); *m/z* = 463.1 ([2M + H]⁺).

***N*^α-Boc-*trans*-3-hydroxy-*L*-proline-NH₂ (16)**—To a cooled solution (−15 °C) of **15** (0.231 g, 0.001 mol) in anhydrous THF (10 mL) was added 4-methylmorpholine (0.111 g, 0.121 mL, 0.0011 mol) and isobutyl chloroformate (0.150 g, 0.144 mL, 0.0011 mol); stirring continued for 30 min. NH₄OH (28% aqueous solution, 0.188 g, 0.206 mL, 0.0015 mol) was added, and the mixture was stirred at −15 °C for 30 min and then warmed to room temperature during 1 h. The solvent was removed under reduced pressure, and the residue was purified directly by silica gel column chromatography (CH₂Cl₂ : MeOH = 9 : 1) to afford a white foamy

solid (0.230 g, quantitative yield). ^1H NMR (500 MHz, CD_3OD) δ 4.315–4.285 (m, 1H), 4.103–4.000 (m, 1H), 3.569–3.517 (m, 2H), 2.062–1.996 (m, 1H), 1.851–1.842 (m, 1H), 1.474–1.401 (m, 9H). ^{13}C NMR (125.7 MHz, CD_3OD) δ (176.2 + 175.8) (1C), (156.7 + 156.2) (1C), (81.7 + 81.5) (1C), (76.5 + 75.5) (1C), (70.6 + 70.3) (1C), (46.2 + 45.7) (1C), (33.7 + 33.1) (1C), (28.82 + 28.76) (1C). ESI (CH_3OH) m/z = 231.2 ($[\text{M} + \text{H}]^+$); m/z = 253.2 ($[\text{M} + \text{Na}]^+$); m/z = 483.3 ($[\text{2M} + \text{Na}]^+$).

***N*^α-Boc-*cis*-4-(4-nitro-benzoyloxy)-*L*-proline- NH_2 (17)**—Diisopropyl

azodicarboxylate (0.230 g, 0.227 mL, 0.0015 mol) was added slowly via syringe to a stirred ice-cooled solution of 16 (0.230 g, 0.001 mol), *p*-nitrobenzoic acid (0.250 g, 0.0015 mol), and triphenyl phosphine (0.393 g, 0.0015 mol) in dry THF (10 mL) under nitrogen. The reaction temperature was allowed to rise to room temperature. After 36 h the solvent was evaporated under reduced pressure, and the residue was purified directly by silica gel column chromatography (hexanes : EtOAc = 3 : 7) to afford a pale yellowish oil (0.23 g, 61%). ^1H NMR (500 MHz, CDCl_3) δ (8.252 + 8.235) (d, 2H, J = 8.5 Hz), (8.166 + 8.149) (d, 2H, J = 8.5 Hz), 6.422 (s, 1H), 6.322 (s, 1H), 5.718 (m, 1H), 4.566 (m, 1H), 3.691 (m, 2H), 2.303 (m, 2H), 1.441 (s, 9H). ^{13}C NMR (125.7 MHz, CDCl_3) δ 171.2 (1C), 164.0 (1C), 154.5 (1C), 150.8 (1C), 135.0 (1C), 131.0 (2C), 123.7 (2C), 81.3 (1C), (75.3 + 74.2) (1C), (63.5 + 62.2) (1C), 44.5 (1C), 30.2 (1C), 28.3 (1C). ESI (CH_3OH) m/z = 380.1 ($[\text{M} + \text{H}]^+$); m/z = 402.3 ($[\text{M} + \text{Na}]^+$); m/z = 780.9 ($[\text{2M} + \text{Na}]^+$).

***N*^α-Boc-*cis*-3-hydroxy-*L*-proline- NH_2 (18)**—A solution of 17 (0.379 g, 0.001 mol) and

NaN_3 (0.195 g, 0.003 mol) in dry MeOH (10 mL) was warmed to 40 °C for 14 h under nitrogen. The solvent was removed under reduced pressure. The residue was purified by silica gel column chromatography (CH_2Cl_2 : MeOH = 9 : 1) to afford a white solid (0.23 g, quantitative yield). ^1H NMR (500 MHz, CD_3OD) δ 4.497–4.485 (m, 1H), 4.172–4.159 (m, 1H), 3.586–3.538 (m, 1H), 3.415–3.351 (m, 1H), 2.016–1.897 (m, 2H), 1.416–1.393 (m, 9H). ^{13}C NMR (125.7 MHz, CD_3OD) δ (175.3 + 175.0) (1C), (156.4 + 156.2) (1C), (81.5 + 81.3) (1C), (72.9 + 72.2) (1C), (66.0 + 65.4) (1C), (45.8 + 45.3) (1C), (33.6 + 33.1) (1C), (28.8 + 28.7) (1C). ESI (CH_3OH) m/z = 231.2 ($[\text{M} + \text{H}]^+$); m/z = 253.3 ($[\text{M} + \text{Na}]^+$); m/z = 483.1 ($[\text{2M} + \text{Na}]^+$).

***N*^α-Boc-*trans*-3-azido-*L*-proline- NH_2 (19)**—To an ice-cooled solution of 18 (0.230 g, 0.001 mol) in dry pyridine (5 mL) was added dropwise methanesulfonyl chloride (0.172 g, 0.117 mL, 0.0015 mol). The reaction was stirred at 0 °C for 2 h, and then kept in a refrigerator overnight. Pyridine was evaporated under reduced pressure, and the residue was partitioned between water (10 mL) and EtOAc (10 mL). The aqueous layer was extracted with EtOAc (10 mL $\times$ 3). The combined organic layers were washed with brine, dried over MgSO_4 , filtered, and evaporated to give the mesylate intermediate, which was used without further purification.

The dry mesylate intermediate was dissolved in 3 mL of dry DMF in the presence of NaN_3 (0.098 g, 0.0015 mol). The resulting mixture was gradually heated to 70 °C over 2 h and then stirred at 70 °C for 21 h. The solvent was removed under reduced pressure. The residue was partitioned between water (10 mL) and EtOAc (10 mL), and the aqueous layer was extracted with EtOAc (10 mL $\times$ 2). The combined organic layers were then washed with brine, dried over MgSO_4 , filtered, and evaporated. The resulting crude oil was then purified by silica gel column chromatography (hexanes : EtOAc = 3 : 7) to provide a thick colorless oil (0.255 g, 62%). ^1H NMR (500 MHz, CDCl_3) δ 4.479 (m, 1H), 4.310 (m, 1H), 3.556–3.500 (m, 2H), 2.223–2.007 (m, 2H), 1.478 (m, 9H). ^{13}C NMR (125.7 MHz, CDCl_3) δ (172.9 + 172.6) (1C), (155.7 + 154.4) (1C), (81.5 + 81.3) (1C), (66.5 + 65.1) (1C), (64.6 + 62.6) (1C), (45.2 + 44.8) (1C), (30.3 + 29.4) (1C), 28.4 (1C). ESI (CH_3OH) m/z = 256.2 ($[\text{M} + \text{H}]^+$); m/z = 278.5 ($[\text{M} + \text{Na}]^+$); m/z = 533.1 ($[\text{2M} + \text{Na}]^+$).

***N*^α-Boc-*trans*-3-amino-*L*-proline-NH₂ (20)—19** (0.225 g, 0.001 mol) was dissolved in MeOH (10 mL) containing a catalytic amount of 10% Pd/C. The solution was stirred under a hydrogen atmosphere at room temperature for 21 h. The reaction mixture was filtered through Celite, and the filtrate was concentrated *in vacuo* to provide a thick colorless oil (0.230 g, quantitative yield). ¹H NMR (500 MHz, CD₃OD) δ 4.745 (brs, 4H), 3.928–3.853 (m, 1H), 3.607–3.546 (m, 1H), 3.531–3.489 (m, 2H), 2.134–2.081 (m, 1H), 1.751–1.707 (m, 1H), 1.471–1.435 (m, 9H). ¹³C NMR (125.7 MHz, CD₃OD) δ (177.0 + 176.6) (1C), (156.5 + 156.1) (1C), (81.6 + 81.3) (1C), (70.3 + 70.0) (1C), (58.4 + 57.4) (1C), (46.4 + 46.0) (1C), (33.5 + 33.0) (1C), (28.9 + 28.8) (1C). ESI (CH₃OH) *m/z* = 230.2 ([M + H]⁺); *m/z* = 252.4 ([M + Na]⁺); *m/z* = 459.3 ([2M + H]⁺); *m/z* = 481.2 ([2M + Na]⁺)

3-*N*-(*N*^α-Boc-*L*-Arg^{NO2})-*N*^α-Boc-*trans*-3-amino-*L*-proline-NH₂ (21)—A solution of *N*^α-Boc-*L*-Arg^{NO2}-OH (0.319 g, 0.001 mol) in dry THF (20 mL) was cooled to –15 °C, and *N*-methylmorpholine (0.111 g, 0.121 mL, 0.0011 mol) and isobutyl chloroformate (0.150 g, 0.144 mL, 0.0011 mol) were added. The reaction mixture was continued to stir for 30 min at –15 °C. **20** (0.252 g, 0.0011 mol) in anhydrous THF (5 mL) was added to the reaction mixture through a cannula. The solution was stirred for 30 min at –15 °C and then allowed to warm to room temperature over 30 min. The reaction mixture was stirred at room temperature for 1 h, and the solvent was evaporated under reduced pressure. The residue was purified by silica gel column chromatography (CH₂Cl₂ : MeOH = 8.5 : 1.5) to afford a colorless oil (0.43 g, 81%). ¹H NMR (500 MHz, CD₃OD) δ 4.364 (m, 1H), 4.083–3.992 (m, 2H), 3.530–3.517 (m, 2H), 3.270–3.234 (m, 2H), 2.154–2.129 (m, 1H), 1.903–1.880 (m, 1H), 1.735–1.518 (m, 4H), 1.518–1.341 (m, 18H). ¹³C NMR (125.7 MHz, CD₃OD) δ 176.1 (1C), (175.6 + 174.8) (1C), 161.0 (1C), 157.9 (1C), (156.3 + 155.9) (1C), (81.9 + 81.6) (1C), 80.8 (1C), (67.1 + 66.9) (1C), (56.1 + 55.8) (1C), (55.5 + 55.4) (1C), (46.4 + 46.0) (1C), 41.9 (1C), (31.2 + 30.8) (1C), (30.6 + 30.5) (1C), (28.9 + 28.7) (6C), 26.0 (1C). ESI (CH₃OH) *m/z* = 531.5 ([M + H]⁺); *m/z* = 553.5 ([M + Na]⁺); *m/z* = 1061.4 ([2M + H]⁺); *m/z* = 1083.2 ([2M + Na]⁺).

3-*N*-(*L*-Arg^{NO2})-*trans*-3-amino-*L*-proline-NH₂ (6)—21 (0.265 g, 0.001 mol) was dissolved in anhydrous CH₂Cl₂ (1 mL), and TFA (1 mL) was added at 0 °C. The reaction mixture was stirred for 1 h at 0 °C and 1 h more at room temperature. After the reaction was completed, the solvent was evaporated *in vacuo*, and the residue was dissolved in water (10 mL). The color impurity was extracted with ethyl acetate (10 mL × 2). The combined aqueous layers were evaporated under high-vacuum rotary evaporator to provide a white foamy solid (0.28 g, quantitative yield). ¹H NMR (500 MHz, D₂O) δ 4.465–4.429 (m, 1H), (4.210 + 4.200) (d, 1H, J = 5Hz), 3.931 (t, 1H, J = 6.5Hz), 3.451 (t, 2H, J = 7Hz), 3.170 (m, 2H), 2.301–2.259 (m, 1H), 2.071–2.031 (m, 2H), 1.864–1.808 (m, 2H), 1.589 (m, 2H). ¹³C NMR (125.7 MHz, D₂O) δ 170.3 (1C), (169.3 + 169.3) (1C), (163.0 + 162.7 + 162.4 + 162.1) (2C, TFA-COOH), 158.8 (1C), (119.7 + 117.4 + 115.1 + 112.8) (2C, TFA-CF₃), 64.2 (1C), (54.4 + 54.3) (1C), (52.9 + 52.8) (1C), 45.3 (1C), 40.8 (1C), 29.3 (1C), 28.0 (1C), 23.2 (1C). ESI (CH₃OH) *m/z* = 331.4 ([M + H]⁺); *m/z* = 661.2 ([2M + H]⁺); HRMS (ES⁺, CH₃OH) Calc.: 331.1829, Found : 331.1841. Comb. Anal. (C₁₁H₂₂N₈O₄·3TFA) C, H, N.

HPLC purification of 3-*N*-(*L*-Arg^{NO2})-*trans*-3-amino-*L*-proline-NH₂ (6)—Column: a Phenomenex Gemini 5μ C18 110A preparative column, column size is 250 × 21.2 mm 5μ micron; injection concentration was 20 mg/mL; flow rate: 12 mL/min; detection was at 254 nm; the mobile phase was H₂O (0.1% TFA): 60% CH₃CN-40% H₂O (0.08% TFA) = 97 : 3, and an isocratic method was used. The retention time was 5 minutes.

Solid-phase synthesis of dipeptide analogue 5—Rink amide resin (0.1 mmol) was treated twice with 20% piperidine in DMF (5 mL) at room temperature for 5 min and 20 min. The resin was washed with DMF (3x), DCM (2x), and MeOH (3x). Deprotected Rink amide

resin (0.1 mmol) was suspended in DMF (1 mL). DIEA (0.2 mmol) and a solution of *N*-Boc-*cis*-3-azido-*L*-proline¹⁹ (**22**, 0.2 mmol) and HBTU (0.2 mmol) in DMF (4 mL) were added sequentially. The resulting mixture was shaken at room temperature for 6 h and filtered. The resin was washed with DMF (3x), DCM (2x), and MeOH (3x). The resulting resin-bound azido proline *cis*-**24** was dried under vacuum for 2 h. Rink amide resin loaded with the azido proline derivative (*cis*-**24**, 0.1 mmol) was suspended in dry THF under nitrogen. A yellow solution of preformed $[\text{Et}_3\text{NH}]^+[\text{Sn}(\text{PhS})_3]^-$ (0.2 mmol) was injected, and the mixture was stirred for 1 h at room temperature. The reaction was quenched with a saturated solution of K_2CO_3 in water. The resin was filtered and washed with water (3x), and THF (3x). Then the resin was treated twice with a mixture of THF/sat. solution of K_2CO_3 (2:1) for 15 min. The resin was filtered and washed with water (3x), methanol (3x), DCM (3x), and methanol (3x). Rink amide resin loaded with the amino proline derivative (*cis*-**25**, 0.1 mmol) was suspended in DMF (1 mL). DIEA (0.4 mmol) and a solution of Boc-*L*-Arg^{NO₂} (0.4 mmol) and HBTU (0.4 mmol) in DMF (4 mL) were added sequentially. The resulting mixture was shaken at room temperature for 3 h and filtered. The resin was washed with DMF (3x), DCM (2x), and MeOH (3x). This coupling reaction was repeated a second time. The resulting resin-bound dipeptide *cis*-**26** was dried under vacuum for 2 h. The resulting Rink amide resin loaded with the dipeptide (*cis*-**26**, 0.1 mmol) was swollen in DCM and filtered. A solution of TFA and DCM (50:50, 6 mL) containing 1% SiHET₃ (TES) was added, and the resulting mixture was shaken for 30 min at room temperature. The suspension was filtered, and the resin was washed with DCM/TFA (8:2, 3x). The solvent was removed on the rotary evaporator, and the residue was treated with DCM and concentrated to dryness. This last step was repeated twice. The obtained dipeptide **5** was pre-purified by solid-phase extraction (SPE) using a C-18 pre-conditioned column in water. The dipeptide was purified by HPLC using a semiprep column [Phenomenex, Luna, 250x10 mm, C18(2)] with a pre-column [Phenomenex, Luna 50x10 mm, C18(2)]. A 100 mL solution of 5 mg/100–200 mL was injected into a Beckman HPLC instrument (System Gold 125, UV detector 166) and chromatographed at a flow rate of 4 mL/min, detecting at 214 nm. Program: [A: water + 0.1% TFA; B: ACN + 0.1% TFA] for 5 min using 1% of B, then 10% of B over 15 min., then 5 min using 10% of B, then 1% of B over 5 min. The total run time was 30 min. Fractions containing the pure dipeptide were concentrated to dryness, and the residue was lyophilized. ¹H NMR (500 MHz, D₂O) δ 4.576–4.561 (d, 1H, J = 7.5 Hz), 4.024–4.000 (t, 1H, J = 6 Hz), 3.705–3.653 (m, 1H), 3.457–3.360 (m, 2H), 3.289 (m, 2H), 2.538–2.467 (m, 1H), 2.173–2.064 (m, 2H), 1.967–1.807 (m, 2H), 1.659 (m, 2H); ¹³C NMR (100.7 MHz, D₂O) δ 169.9 (1C), 167.8 (1C), 159.1 (1C), 62.5 (1C), 52.8 (1C), 51.0 (1C), 44.4 (1C), 40.3 (1C), 29.7 (1C), 28.0 (1C), 23.9 (1C); ESI(CH₃OH) *m/z* = 331.5([M + H]⁺); *m/z* = 661.3([2M + H]⁺). Compound **6** also was made by this same solid-phase method except starting with **23**. The compound obtained was identical in all respects to that synthesized by the solution-phase method described above.

Method for the preparation of triethylammonium tris(phenylthio)stannate

Preparation of $[\text{Et}_3\text{NH}]^+[\text{Sn}(\text{PhS})_3]^-$ (Caution: stench): SnCl_2 (0.2 mmol, 2 equiv.) was solubilized in dry THF by sonicating the suspension under nitrogen. Thiophenol (0.8 mmol, 8 equiv), and TEA (1 mmol, 10 equiv.) were sequentially injected under nitrogen. The obtained yellow suspension was filtered into a dry syringe using a 0.45 μm filter on line. This solution was used immediately.

General method for the solid-phase synthesis of *N*-substituted derivatives of 4-*N*-(*L*-Arg^{NO₂})-*trans*-4-amino-*L*-proline-NH₂ (**3**, **4**) (see Supporting Information Scheme 1)

Rink amide resin (0.1 mmol) was treated twice with 20% piperidine in DMF (5 mL) at room temperature for 5 min and 20 min. The resin was washed with DMF (3x), DCM (2x), and MeOH (3x). The deprotected Rink amide resin (0.1 mmol) was suspended in DMF (1 mL). DIEA (0.2 mmol) and a solution of *N*-Boc-*trans*-4-azido-*L*-proline (0.2 mmol) and HBTU (0.2

mmol) in DMF (4 mL) were added sequentially. The resulting mixture was shaken at room temperature for 6 h and filtered. The resin was washed with DMF (3x), DCM (2x), and MeOH (3x). The resulting resin-bound amino acid **28** was dried under vacuum for 2 h. Rink amide resin loaded with the azido proline derivative **28** (0.1 mmol) was suspended in dry THF under nitrogen. A yellow solution of preformed $[\text{Et}_3\text{NH}]^+[\text{Sn}(\text{PhS})_3]^-$ (0.2 mmol) was injected, and the mixture was stirred for 1 h at room temperature. The reaction was quenched with a saturated solution of K_2CO_3 in water. The resin was filtered and washed with water (3x) and THF (3x). Then the resin was treated twice with a mixture of THF/sat. solution of K_2CO_3 (2:1) for 15 min. The resin was filtered and washed with water (3x), methanol (3x), DCM (3x), and methanol (3x). Rink amide resin loaded with the amino proline derivative **29** (0.1 mmol) was suspended in NMP (1 mL). DIEA (0.4 mmol) and a solution of Fmoc-*L*-Arg^{NO₂} (0.4 mmol) and HBTU (0.4 mmol) in NMP (4 mL) were added sequentially. The resulting mixture was shaken at room temperature for 3 h and filtered. The resin was washed with DMF (3x), DCM (2x), and MeOH (3x). This coupling reaction was repeated for a second time. The resulting resin-bound dipeptide **30** was dried under vacuum for 2 h. The resulting Rink amide resin loaded with the dipeptide (**30**, 0.1 mmol) was treated twice with 20% piperidine in DMF (5 mL) at room temperature for 5 min and 20 min. The resin was washed with DMF (3x), DCM (2x), and MeOH (3x). Rink amide resin **31** (0.1 mmol) was pre-swollen with dry THF and a solution of (trifluoroacetyl)benzotriazole (1 mmol) in dry THF was added. The suspension was shaken at room temperature for 2 h. The resin was washed with THF (3x), MeOH (3x), DCM (3x), and MeOH (3x). The resulting resin-bound protected dipeptide **32** was dried under vacuum for 2 h. The Boc-protected proline-containing dipeptide loaded resin **32** was dried overnight under nitrogen and then suspended in dry DCM (0.1 mmol in 3 mL of dry DCM). The suspension was cooled in an ice-bath, and 2,6-lutidine (0.6 mmol) and TMSOTf (0.4 mmol) were sequentially injected dropwise under argon. The suspension was warmed to room temperature after 10 min and left stirring for 2 h under argon. The reaction was quenched using a saturated solution of ammonium chloride in water. The resin **33** was washed with DCM (3x), MeOH (3x), water (3x), MeOH (3x), and DCM (3x). Deprotected intermediate **33** (0.1 mmol) was dried under vacuum, and a freshly prepared solution of formaldehyde (for **3**) or benzaldehyde (for **4**) (1 mmol) in trimethylorthoformate (4 mL) was added under nitrogen. The same solution (2 mL) was added again after 1 h at room temperature. The suspension was stirred for 1 h at room temperature, and NaBH_3CN (1 mmol) was added in 1 mL of TMOF. The suspension was washed after 15 h with MeOH (3x), DCM (3x), MeOH (3x), H_2O (2x), and MeOH (2x). Resin **34** was dried under vacuum for 2 h. The *N*^α-substituted loaded resin (**34**, 0.1 mmol) was pre-swollen with THF, and a solution of LiOH (1 mmol) in THF/ H_2O (5:1) was added. The resin was shaken at room temperature for 3 h and washed with THF (3x), MeOH (3x), H_2O (3x), and MeOH (3x). The deprotected dipeptide loaded resin **35** was dried under vacuum for 2 h. The resulting Rink amide resin loaded with the dipeptide (**35**, 0.1 mmol) was swollen in DCM and filtered. A solution of TFA and DCM (50:50, 6 mL) containing 1% SiHEt_3 (TES) was added, and the resulting mixture was shaken for 30 min. at room temperature. The suspension was filtered, and the resin was washed with DCM/TFA (8:2, 3x). The solvent was removed on a rotary evaporator, and the residue was treated with DCM and concentrated to dryness. This last step was repeated twice. The obtained dipeptide **3** or **4** was further purified by solid-phase extraction (SPE) using a C-18 pre-conditioned column in water.

Enzyme and assay

All of the NOS isoforms used were recombinant enzymes overexpressed in *E.coli*. The murine macrophage iNOS was expressed and isolated according to the procedure of Hevel et al.²⁴ The rat nNOS was expressed²⁵ and purified as described. The bovine eNOS was isolated as reported.²⁶ Nitric oxide formation from NOS was monitored by the hemoglobin capture assay as described.²⁷

Determination of K_i values

The apparent K_i values were obtained by measuring percent inhibition in the presence of 10 μM *L*-arginine with at least three concentrations of inhibitor. The parameters of the following inhibition equation²⁸ were fitted to the initial velocity data: % inhibition = $100[I]/\{[I] + K_i(1 + [S]/K_m)\}$. K_m values for *L*-arginine were 1.3 μM (nNOS), 8.2 μM (iNOS), and 1.7 μM (eNOS). The selectivity of an inhibitor was defined as the ratio of the respective K_i values.

Molecular modeling

Docking simulations were carried out with AutoDock 3.0.5 as described by Ji et al.²² The protein crystal structure used in the docking studies is that of (4*S*)-*N*-(4-amino-5-[aminoethyl]aminopentyl)-*N'*-nitroguanidine bound to nNOS (PDB ID: 1p6i). The ligands were built in SYBYL version 6.8²⁹ by modifying the molecular structure of **2**, which was extracted from the crystal structure of 1p6j (PDB ID). Energy minimizations were performed following the addition of hydrogen atoms and partial atom charge calculations by the Gasteiger-Marsili method.³⁰ Appropriate atom types were specified and labeled for physiological conditions, and ligand chirality was inspected. The grid maps were calculated using AutoGrid. The dimensions of the grid box was $31 \times 28 \times 31$ Å. Default parameters as described in detail previously were used, and 100 docked conformations were yielded. A visual comparison of the superimposition of the guanidino moiety of the calculated conformations and that of **2** served as a preliminary identification of appropriate structures. The binding affinity of the inhibitors to the protein were then evaluated by the total AutoDock docking energies and further rescored using the CScore module of Sybyl 6.8 comprising the following scoring functions: F_Score, D_Score, PMF_Score, G_Score and ChemScore. For rescoring docked poses, atomic coordinates of the target protein were unchanged, and relaxation of the bound ligand was performed.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to the National Institutes of Health (GM 49725 to R.B.S. and GM52419 and HL30050 to Prof. Bettie Sue Masters, in whose laboratory P.M. and L.J.R. work) for financial support of this work. Grants from NATO and the Spanish Ministry of Science and Technology to J.A.G.V. also are gratefully acknowledged.

References and Notes

1. Mayer B. Nitric Oxide. Handbook of Experimental Pharmacology 2000;143
2. Moncada S, Higgs A, Furchgott R. XIV. International union of pharmacology nomenclature in nitric oxide research. Pharmacol. Rev 1997;49:137–142. [PubMed: 9228663]
3. Hobbs AJ, Higgs A, Moncada S. Inhibition of nitric oxide synthase as a potential therapeutic target. Annu. Rev. Pharmacol. Toxicol 1999;39:191–220. [PubMed: 10331082]
4. Crane BR, Arvai AS, Gachhui R, Wu C, Ghosh DK, Getzoff ED, Stuehr DJ, Tainer JA. The structure of nitric oxide synthase oxygenase domain and inhibitor complexes Science 1997;278:425–431.
5. Crane BR, Arvai AS, Ghosh S, Getzoff ED, Stuehr DJ, Tainer JA. Structures of the N^7 -hydroxy-*L*-arginine complex of inducible nitric oxide synthase oxygenase dimer with active and inactive pterins. Biochemistry 2000;39:4608–4621. [PubMed: 10769116]
6. Crane BR, Arvai AS, Ghosh DK, Wu C, Getzoff ED, Stuehr DJ, Tainer JA. Structure of nitric oxide synthase oxygenase dimer with pterin and substrate Science 1998;279:2121–2126.
7. Fischmann TO, Hruza A, Niu XD, Fossetta JD, Lunn CA, Dolphin E, Prongay AJ, Reichert P, Lundell DJ, Narula SK, Weber P. Structural characterization of nitric oxide synthase isoforms reveals striking active-site conservation. Nature Struct. Biol 1999;6:233–242. [PubMed: 10074942]

8. Li H, Raman CS, Glaser CB, Blasko E, Young TA, Parkinson JF, Whitlow M, Poulos TL. Crystal structures of zinc-free and -bound heme domain of human inducible nitric-oxide synthase. *J. Biol. Chem* 1999;274:21276–21284. [PubMed: 10409685]
9. Raman CS, Li H, Martásek P, Král V, Masters BSS, Poulos TL. Crystal structure of constitutive endothelial nitric oxide synthase: A paradigm for pterin function involving a novel metal center. *Cell* 1998;95:939–950. [PubMed: 9875848]
10. (a) Flinspach ML, Li H, Jamal J, Yang W, Huang H, Hah J-M, Gomez-Vidal JA, Litzinger EA, Silverman RB, Poulos TL. Structural basis for dipeptide amide isoform-selective inhibition of neuronal nitric oxide synthase. *Nature (Struct. Mol. Biol.)* 2004;11:54–59. [PubMed: 14718923] (b) Flinspach M, Li H, Jamal J, Yang W, Huang H, Silverman RB, Poulos TL. Structures of the Neuronal and Endothelial NOS Heme Domain with D-Nitroarginine-containing Dipeptide Inhibitors Bound. *Biochemistry* 2004;43:5181–5187. [PubMed: 15122883] (c) Li H, Flinspach ML, Igarashi J, Jamal J, Yang W, Gómez-Vidal JA, Litzinger EA, Huang H, Erdal EP, Silverman RB, Poulos TP. Exploring the Binding Conformations of Bulkier Dipeptide Amide Inhibitors in Constitutive Nitric Oxide Synthases. *Biochemistry* 2005;44:15222–15229. [PubMed: 16285725]
11. Alcaraz MJ, Guillen MI. Nitric oxide related therapeutic phenomenon: A challenging task. *Curr. Pharm. Des* 2002;8:215–231. [PubMed: 11812268]
12. Lowe JA III. Nitric oxide synthase inhibitors: Recent patent activity. *IDrugs* 2000;3:63–72.
13. Salerno L, Sorrenti V, Di Giacomo C, Romeo G, Siracusa MA. Progress in the development of selective nitric oxide synthase (NOS) inhibitors. *Curr. Pharm. Des* 2002;8:177–200. [PubMed: 11812267]
14. Silverman RB, Huang H, Marletta MA, Martásek P. Selective inhibition of neuronal nitric oxide synthase by N⁰-nitroarginine- and phenylalanine-containing dipeptides and dipeptide esters. *J. Med. Chem* 1997;40:2813–2817. [PubMed: 9288162]
15. Huang H, Martásek P, Roman LJ, Masters BSS, Silverman RB. N⁰-nitroarginine-containing dipeptide amides. Potent and highly selective inhibitors of neuronal nitric oxide synthase. *J. Med. Chem* 1999;42:3147–3153. [PubMed: 10447959]
16. Gomez-Vidal JA, Martasek P, Roman LJ, Silverman RB. *J. Med. Chem* 2004;47:703–710. [PubMed: 14736250]
17. Gómez-Vidal JA, Forrester MT, Silverman RB. Mild and selective sodium azide mediated cleavage of p-nitrobenzoic esters. *Org. Lett* 2001;3:2477–2479. [PubMed: 11483039]
18. Gómez-Vidal JA, Silverman RB. Short, highly efficient syntheses of protected 3-azido- and 4-azidoproline and their precursors. *Org. Lett* 2001;3:2481–2484. [PubMed: 11483040]
19. Bartra M, Romea P, Urpí F, Vilarrasa J. A Fast Procedure for the Reduction of Azides and Nitro Compounds Based on the Reducing Ability of Sn(SR)₃[−] Species. *Tetrahedron* 1990;46:587–594.
20. Kim J, Bi Y, Paikoff SJ, Schultz PG. The Solid Phase Synthesis of Oligoureas. *Tetrahedron Lett* 1996;37:5305–5308.
21. a) Kick E, Ellman JA. Expedient Method for the Solid-Phase Synthesis of Aspartic Acid Protease Inhibitors Directed toward the Generation of Libraries. *J. Med. Chem* 1995;38:1427–1430. [PubMed: 7739000] b) Tortolani DR, Biller SA. A Solid Phase Synthesis of Miconazole Analogs via an Iodoetherification Reaction. *Tetrahedron Lett* 1996;37:5687–5690.
22. Ji H, Li H, Flinspach M, Poulos TL, Silverman RB. Computer Modeling of Selective Regions in the Active Site of Nitric Oxide Synthases: Implication for the Design of Isoform-Selective Inhibitors. *J. Med. Chem* 2003;46:5700–5711. [PubMed: 14667223]
23. (a) Terp GE, Johansen BN, Christensen IT, Jorgensen FS. A new concept for multidimensional selection of ligand conformations (MultiSelect) and multidimensional scoring (MultiScore) of protein-ligand binding affinities. *J. Med. Chem* 2001;44:2333–2343. [PubMed: 11428927] (b) Wang R, Lu Y, Wang S. Comparative Evaluation of 11 Scoring Functions for Molecular Docking. *J. Med. Chem* 2003;46:2287–2303. [PubMed: 12773034]
24. Hevel JM, White KA, Marletta M. A purification of the inducible murine macrophage nitric oxide synthase. *J. Biol. Chem* 1991;266:22789–22791. [PubMed: 1720773]
25. Roman LJ, Sheta EA, Martásek P, Gross SS, Liu Q, Masters BSS. High-level expression of functional rat neuronal nitric oxide synthase in escherichia coli. *Proc.Natl. Acad. Sci. U.S.A* 1995;92:8428–8432. [PubMed: 7545302]

26. Martasek P, Liu Q, Roman LJ, Gross SS, Sessa WC, Masters BSS. Characterization of bovine endothelial nitric oxide synthase expressed in *Escherichia coli*. *Biochem. Biophys. Res. Commun* 1996;219:359–365. [PubMed: 8604992]
27. Hevel JM, Marletta MA. Nitric oxide synthase assays. *Methods Enzymol* 1994;133:250–258. [PubMed: 7516999]
28. Segel, IH. *Enzyme Kinetics*. John Wiley and Sons; NewYork: 1975. p. 105
29. SYBYL Molecular Modelling Package. version 6.8. Tripos; St. Louis, MO: 2001.
30. Gasteiger J, Marsili M. Iterative partial equalization of orbital electronegativity-a rapid access to atomic charges. *Tetrahedron* 1980;36:3219–3228.

Scheme 1.
Solution-phase synthesis of **3** and **4**

Scheme 2.
Solution-phase synthesis of **6**

Scheme 3.
Solid-phase synthesis of **5** and **6**

Figure 1.
The H-bond network of the pyrrolidine nitrogen atom of **2** bound to rat nNOS (PDB id 1p6j).
The distances of the hydrogen bonds are shown in Angstroms (Å).

Figure 2. Superimposition of docked conformation of **4** and the crystallographic conformation of **27** in rat nNOS (PDB id 1rs7). The blue conformation is the crystallographic binding conformation of **27**; the green structure is the docking conformation of **4**.

A **B****Figure 3.**

Superimposition of the crystallographic conformation of **2** (PDB id 1p6j) and the docked conformations of **5** and **6** in rat nNOS. A: The blue conformation is the crystallographic binding conformation of **2**; the green structure is the docking conformation of **6**. B: The blue conformation is the crystallographic binding conformation of **2**; the green structure is the docking conformation of **5**.

Table 1

In vitro Assay Data for Dipeptide Inhibitors.

Compound	nNOS (IC ₅₀) ^a	nNOS (K _i) ^a	iNOS (IC ₅₀) ^d	iNOS (K _i) ^d	eNOS (IC ₅₀) ^d	eNOS (K _i) ^d	iNOS/nNOS ^g	eNOS/nNOS ^g
1	1.1	0.13 ^c	55.5	25 ^c	1375	200	190	1540
2 ^b	0.64	0.10 ^e	65	29	880	128	290	1280
3 ^h	73.5 ± 3.5	8.46 ± 0.40	83.5 ± 3.5	37.9 ± 1.6	1690 ± 100	246 ± 14	4.5	29
4 ^h	2.85 ± 0.21	0.328 ± 0.024	87.5 ± 2.1	39.7 ± 1.0	1100 ± 90	159 ± 13	121	485
5	100	-	-	-	-	-	-	-
6 ^h	0.75 ± 0.03	0.087 ± 0.004	12.7 ± 1.1	5.76 ± 0.61	1380 ± 80	200 ± 10	66	2299
27 ^f	17.4	2 ^c	8 mM	3600 ^c	-	-	1800	-

^a μM;

^b from ref. 15;

^c experimental K_i;

^d from ref. 16;

^e experimental K_i = 103 nM;

^f from ref. 14;

^g selectivity of inhibition in favor of nNOS inhibition obtained from the ratio of the K_i for iNOS or eNOS over that for nNOS.

^h The K_i values for **3–6** represent two separate measurements with 5–6 data points each and correlation coefficients of 0.947–0.992.

Table 2

Predicted binding free energies (Kcal/mol) of **2**, **5** and **6** with nNOS using AutoDock, Chemscore, F_Score, PMF_Score, E_docked, G_Score, and D_Score scoring functions.

	Chemscore	F_Score	PMF_Score	E_docked	G_Score	D_Score
2	-17.86	-18.38	-143.53	-11.94	-283.45	-23.95
5	-12.51	-7.30	-121.31	-9.26	-226.56	-25.21
6	-19.06	-19.18	-134.04	-10.14	-239.99	-21.25