Published in final edited form as:

Org Lett. 2007 August 2; 9(16): 3093–3096.

Stereoselective [2,3]-Sigmatropic Rearrangements of Unstabilized Nitrogen Ylides

Robert E. Gawley and Kwangyul Moon

Department of Chemistry and Biochemistry, University of Arkansas, Fayetteville, AR 72701, bgawley@uark.edu

Abstract

The steric course of the [2,3]-rearrangement of several unstabilized nitrogen ylides has been investigated. The reactions proceed cleanly through an *anti* transition state, affording modest to good yields of a single diastereomer of the product. In two examples containing an *N*-cinnamyl group, a competing [1,2]-rearrangement affords a minor product.

Electrophilic substitution reactions of α -alkoxy- and α -aminoorganolithium compounds (Eq 1) have been extensively studied, and constitute an important class of synthetic methods, with hundreds of applications reported so far.^{1,2} A second class of carbon-carbon bond forming reactions utilizing these reactive intermediates is sigmatropic rearrangements.^{3,4} In 1978, Still and Mitra reported the first example of a [2,3]-rearrangement of an organolithium generated by tin-lithium exchange,⁵ which was later shown to be invertive at the lithium-bearing carbon (Eq 2).^{6–8}

(1)

Correspondence to: Robert E. Gawley.

Warmly dedicated to Professor Miguel Yus, on the occasion of his 60th birthday

The nitrogen (aza) analog of the Still-Wittig [2,3]-rearrangement (Eq 3) is less facile, due primarily to the lower stability of the lithium amide compared to the lithium alkoxide product of the oxygen version.^{3,4,9,10} Once generated, the *N*-allyl α -aminoorganolithium often follows competing [1,2]- and [2,3]-rearrangement pathways (Eq 3).^{11–13} Quaternization of the nitrogen (Eq 4) with a trimethylsilyl group,¹⁴ boron trifluoride,¹⁵ or an alkyl group^{13, 16} accelerates the concerted [2,3]-rearrangement over the radical-mediated [1,2]-rearrangement of the ylides. The steric course of the anionic and ylide aza-[2,3]-rearrangements is invertive if the metal bearing carbon is stereogenic (Eqs 3 and 5),¹³ but the stereoselectivity at the migration terminus of nitrogen ylides has only been explored in stabilized ylides such as the enolate shown in Eq 4. In most cases, the diastereoselectivity is modest,^{16–18} with only a few examples of high selectivity in auxiliary mediated processes.^{19,20} We now report the results of our studies on the diastereoselectivity at the migration terminus of the [2,3]-rearrangement of several unstabilized, lithionitrogen ylides (Eq 5), in which we find that the rearrangement is highly stereoselective. *Note:* Strictly speaking, the ylide is the zwitterion having a positive nitrogen and a negative carbon. Since the anionic carbon is stereogenic, we draw the carbon-lithium ion pair as a bond.

Stannane **1** is a challenging test of the methodology for two reasons. Firstly, stannane **1** fails to undergo tin-lithium exchange, so it was of interest to determine whether quaternization of the nitrogen would facilitate the exchange. Secondly, we wanted to evaluate the diastereoselectivity of the [2,3]-rearrangement in the context of acyclic stereoselection. To test the first point, **1** was simply quaternized with methyl iodide. Treatment of the methiodide salt

with butyllithium in THF at $-78\text{ }^{\circ}\text{C}$ for 1 h, quenching with methanol, and silica gel chromatography afforded a 69% yield of tetrabutyltin, indicating that transmetalation to the ylide is facile in this system. Quaternization of racemic **1** with *E*-crotyl bromide, and *E*- and *Z*-cinnamyl bromide affords ammonium salts **2a–c**, as shown in Scheme 1. For some compounds, anion exchange from halide to hexafluorophosphate facilitated handling, provided a salt that was less hygroscopic, and increased solubility in organic solvents (see Supporting Information).

As illustrated in Scheme 2, stannylammonium ions **2a–c** were treated with butyllithium in THF at $-60\text{ }^{\circ}\text{C}$ to effect transmetalation and stirred at that temperature for 20–24 h. After workup and column chromatography, amines **5a–c** were obtained in the yields shown in Table 1, entries 1–3.

Of interest in these rearrangements is the possibility of two transition state conformers, *syn* and *anti* **4a–c**, which would afford diastereomeric products **5** or **6**, and possibly [1,2]-rearrangement product **7**. In all cases, only one diastereomer was isolated. Independent synthesis of **5b** and **5c** established the relative configurations (see Supporting Information). It appears that the [2,3]-rearrangement prefers the *anti* transition state **4**, independent of alkene geometry, even though *anti* **4c** having a *Z* double bond appears somewhat more congested. The lower yield from the *Z*-cinnamyl intermediate **3c** may indicate that steric crowding in the *anti* transition state slows the rearrangement, allowing pathways toward nonproductive decomposition to become more competitive; in this case, there were numerous unidentified polar byproducts. It is noteworthy that under these conditions, none of the [1,2]-products **7a–c** were detected.

If R^2 and R^3 in Eq 5 are different, then the nitrogen is a stereocenter and the stereoselectivity of the rearrangement will rely on the stereoselective quaternization of the nitrogen. To probe the diastereoselectivity of the rearrangement in cyclic systems, we chose racemic *N*-methyl-2-(tributylstannyl)-piperidine, **8**. Based on precedent from a single example in an earlier report,¹³ we anticipated that alkylation of **8** would occur *trans* to the tin.

In the event, alkylation of **8** with *E*-crotyl bromide and with *E*- and *Z*-cinnamyl bromide afforded ammonium ions **9a–c**, with 90–94% diastereoselectivity, as indicated by integration of the *N*-methyl peaks in the NMR (Scheme 3). Again, exchange of the halide anion for PF_6^- often facilitated handling of the salt, increased its solubility in organic solvents, and made it less hygroscopic.

Transmetalation of **9a–c** afforded ylides **10a–c**, and ultimately their rearrangement products, as summarized in Table 1, entries 4–6, and illustrated in Scheme 4. The *E*-crotyl ylide **10a** rearranged to a mixture of isomers, of which piperidine **12a** was the major (92%) component. The isomers were not obtained in sufficient quantity to identify; one may be the [1,2]-product **14a**, while others are diastereomers that could have arisen from transition structure *syn*-**11**, or from minor contaminants of *cis* alkene ylide (**10**, $\text{R}^2=\text{Me}$, $\text{R}^1=\text{H}$). *Trans*-cinnamyl ylide **10b** afforded a 3.6:1 mixture of [2,3]-product **12b** and [1,2]-product **14b**, while *cis*-cinnamyl ylide **10c** afforded a 6.9:1 mixture of [2,3]-product **12c** and [1,2]-product **14c**. None of the diastereomeric products **13b–c** were detected by GC-MS, and the structure of **12b** was confirmed by independent synthesis (See Supporting Information). To further confirm these assignments, stannane **8** was transmetalated with BuLi at $-78\text{ }^{\circ}\text{C}$ in THF, and alkylated with *trans* cinnamyl bromide according to our established procedure.²¹ Analysis of the product mixture revealed a mixture of $\text{S}_{\text{N}}2$ product **14b**, along with $\text{S}_{\text{N}}2'$ products **12b** and **13b**. In the [2,3]-rearrangement of **10b–c**, the extra stabilization afforded by the phenyl group probably facilitates homolytic cleavage of the allylic C–N bond to produce **14b–c** after radical recombination. In summary, nitrogen ylides **3a–c** and **10a–c** rearrange preferentially through

transition structures *anti*-4 and *anti*-11, creating two adjacent stereocenters with a high degree of stereoselectivity. Further applications of the [2,3]-rearrangement of unstabilized nitrogen ylides are under active investigation and will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

Acknowledgement is made to the donors of the Petroleum Research Fund, administered by the American Chemical Society, for partial support of this work. We also thank NIH for partial support (GM 56271). Core facilities were funded by NIH (P20 R15569) and the Arkansas Biosciences Institute. We are grateful to Dr Donna Lula for exploratory experiments.

References

1. Hoppe D, Marr F, Brüggemann M. *Top Organomet Chem* 2003;5:61–138.
2. Beak P, Johnson TA, Kim DD, Lim SH. *Top Organomet Chem* 2003;5:139–176.
3. Markó, I. *Comprehensive Organic Synthesis*. In: Trost, BM.; Fleming, I., editors. *Selectivity, Strategy and Efficiency in Modern Organic Chemistry*. 3. Pergamon; Oxford: 1991. p. 913-974.
4. Tomooka, K. *The chemistry of organolithium compounds*. Rappoport, Z.; Marek, I., editors. John Wiley & Sons, Ltd.; Chichester: 2004. p. 749-828.
5. Still WC, Mitra A. *J Am Chem Soc* 1978;100:1927–1928.
6. Verner EJ, Cohen T. *J Am Chem Soc* 1992;114:375–377.
7. Hoffmann R, Brückner R. *Angew Chem Int Ed Engl* 1992;31:647–649.
8. Tomooka K, Igarashi T, Watanabe M, Nakai T. *Tetrahedron Lett* 1992;33:5795–5798.
9. Nakai T, Tomooka K. *Pure Appl Chem* 1997;69:595–600.
10. Gawley, RE.; Coldham, I. *The Chemistry of Organolithium Compounds (Patai Series)*. Rappoport, Z.; Marek, I., editors. Wiley; Chichester: 2004. p. 997-1053.
11. Heard GL, Yates BF. *J Org Chem* 1996;61:7276–7284. [PubMed: 11667650]
12. Coldham I. *J Chem Soc Perkin Trans* 1993;1:1275–1276.
13. Gawley RE, Zhang Q, Campagna S. *J Am Chem Soc* 1995;117:11817–11818.
14. Murata Y, Nakai T. *Chemistry Letters* 1990:2069–2072.
15. Kessar SV, Singh P, Singh KN, Kaul VK, Kumar G. *Tetrahedron Lett* 1995;36:8481–8484.
16. Coldham I, Middleton ML, Taylor PL. *J Chem Soc, Perkin Trans 1* 1998:2817–2821.
17. Roberts E, Sançon JP, Sweeney JB, Workman JA. *Org Lett* 2003;5:4775–4777. [PubMed: 14653671]
18. Zhou CY, Yu WY, Chan PWH, Che CM. *J Org Chem* 2004;69:7072–7082. [PubMed: 15471455]
19. Aggarwal VK, Fang Gy, Charmant JPH, Meek G. *Org Lett* 2003;5:1757–1760. [PubMed: 12735770]
20. Workman JA, Garrido NP, Sançon J, Roberts E, Wessel HP, Sweeney JB. *J Am Chem Soc* 2005;127:1066–1067. [PubMed: 15669822]
21. Gawley RE, Zhang Q. *J Org Chem* 1995;60:5763–5769.

Scheme 1.

Scheme 2.

Scheme 3.

Scheme 4.

Table 1
Rearrangements of ylides (Stannane + BuLi, -60 °C, unless otherwise noted).

Entry	Ylide	R ¹	R ²	X ⁻	Product(s)	Yield
1	2a	Me	H	I ⁻	5a	57%
2	2b	Ph	H	PF ₆ ⁻	5b	54%
3	2c	H	Ph	PF ₆ ⁻	5c	29%
4	10a	Me	H	PF ₆ ⁻	12a	60%
5 [*]	10b	Ph	H	Br ⁻	12b+14b (4:1)	55%
6 [*]	10c	H	Ph	Br ⁻	12c+14c (9:1)	35%

^{*}T = -83 °C; MeLi instead of BuLi for transmetalation.