Published in final edited form as:

J Am Chem Soc. 2007 May 30; 129(21): 6684–6685.

A modular approach for facile biosynthesis of labdane-related diterpenes

Anthony Cyr[†], P. Ross Wilderman, Mara Determan, and Reuben J. Peters

Iowa State University, Department of Biochemistry, Biophysics, and Molecular Biology, Ames, IA 50011

Abstract

Labdane-related diterpenoids are a large group of over 5,000 natural products whose biosynthesis typically proceeds through a labdadienyl/copalyl diphosphate (CPP) intermediate to a further cyclized and/or rearranged hydrocarbon diterpene en route to more elaborated compounds. Here we report a modular approach for facile biosynthesis of labdane-related diterpenes wherein base pGGxC vectors capable of introducing bacterial production of any one of the three common stereoisomers of CPP can be co-introduced with diterpene synthases that convert these CPP intermediates to specific diterpene hydrocarbon skeletal structures. The utility of this approach is demonstrated by individually engineering *E. coli* to produce any one of eight different diterpene skeletal structures, which collectively serve as precursors to literally thousands of distinct natural products.

Labdane-related diterpenoids comprise a large group of over 5,000 known natural products defined as minimally containing the fused bicyclic hydrocarbon structure found in the labdane family of diterpenoids. This characteristic core structure results from the unusual biosynthetic origins of these compounds, which is uniquely initiated by a sequential pair of terpene synthase catalyzed reactions. In particular, cyclization of the universal diterpenoid precursor (*E,E,E*)-geranylgeranyl diphosphate (GGPP, **1**) to a specific stereoisomer of labdadienyl/copalyl diphosphate (CPP, e.g. **2-4**) in a carbon-carbon double bond protonation-initiated reaction catalyzed by class II diterpene cyclases. This core bicyclic structure is then further cyclized and/or rearranged in diphosphate ionization initiated reactions catalyzed by more typical class I, but generally CPP stereospecific, labdane-related diterpene synthases, as in the related/derived structural families (e.g. kauranes, abietanes, and pimaranes).¹

Included in the group of labdane-related diterpenoids are the gibberellin phytohormones required for normal growth and development in all higher plants, which has provided a ready reservoir of biosynthetic genes whose duplication has led to the widespread occurrence of such natural products throughout the plant kingdom. However, the vast majority of these compounds are secondary metabolites only found in a subset of species, in limited quantities, and often only under certain conditions. Thus, although several such compounds are utilized in industry, research, and medicine (e.g. resin acids and forskolin), similar uses for the bulk of this group of natural products have not been well explored. The use of recombinant microbial hosts for production of specific isoprenoid/terpenoid natural products has been recently reviewed.² To provide a foundation for broader investigations of labdane-related diterpenoid natural products we report here development of a modular approach that facilitates recombinant bacterial expression of various combinations of the sequentially acting class II and class I diterpene

E-mail: rjpeters@iastate.edu.

[†]Current address: University of Iowa, Medical Scientist Training Program

synthases, to produce hydrocarbon diterpene precursors to literally thousands of known natural products.

Because diterpene biosynthesis in plants occurs in plastids, recombinant microbial expression of the associated enzymes requires construction of pseudo-mature genes missing the N-terminal plastid targeting peptide sequence that interferes with proper protein folding.³ This was recently accomplished for the *ent*-CPP synthase from *Arabidopsis thaliana* (AtCPS),⁴ providing a guide for construction of pseudo-mature versions of other strictly class II diterpene cyclases and, thus, enabling the studies reported here.

While all organisms produce terpenoids/isoprenoids, *E. coli* does not typically produce GGPP, hence it was necessary to introduce a GGPP synthase (GGPS), which was done through the use of a previously reported recombinant pseudo-mature GGPS from grand fir (*Abies grandis*), rAgGGPS.⁵ To minimize the number of plasmids required for these metabolic engineering efforts, rAgGGPS was introduced into the dual gene expression vector pACYCDuet, creating a base pGG plasmid (see Supporting Information). For initial proof-of-concept studies a previously reported pseudo-mature version of the bifunctional class II/I diterpene cyclase from grand fir, abietadiene synthase (rAgAS)⁶ was added to pGG to create pGGAS. Transformation with pGGAS yielded recombinant bacteria that produce the expected abietadiene (**5**) diterpene upon induction of the encoded enzymatic genes (Scheme 1).

From this initial success we were encouraged to attempt the expression of more typical separate class II and class I labdane-related diterpene synthases. In particular, because we have recently reported characterization of a pseudo-mature version of AtCPS, this gene was incorporated into pGG to create pGGeC, which should enable bacterial production of *ent*-CPP (**2**). We also have reported functional recombinant expression of tagged (e.g. with glutathione-S-transferase, GST) versions of class I labdane-related diterpene synthases,⁷ including that for the *ent*-CPP specific kaurene synthase (KS) from *Arabidopsis thaliana* (AtKS),⁸ using pET based expression vectors (e.g. pDEST), which can be co-transformed with pACYCDuet and, hence, the derived pGG and pGGeC. Co-transformation of pGGeC with a pDEST15/AtKS vector encoding a GST-AtKS fusion protein yielded recombinant bacteria that produce the expected *ent*-kaur-16-ene (**6**) (Figure 1).

More detailed analysis of this kaurene (**6**) producing strain was carried out to optimize diterpene production. Somewhat surprisingly, ~90% of the hydrocarbon product was secreted to the media, which had not been previously reported.⁹ Similar secretion was observed with all the labdane-related diterpenes we have tested. Accordingly, it is possible to trap these hydrocarbon compounds on hydrophobic (Diaion HP-20) beads under mixed phase liquid-solid (i.e. media-beads) growth conditions,¹⁰ and easily isolate the produced diterpenes (e.g. Figure 1). With the optimized conditions it was possible to obtain ~100 μ g of kaurene (**6**)/L mixed phase bacterial culture, similar to previously reported terpenoid yield from *E. coli* that has not been engineered for increased isoprenoid precursor supply.¹¹

The vast majority of labdane-related diterpenoid natural products are derived from three of the four possible stereoisomers of CPP, specifically, *ent*- (**2**), *syn*- (**3**), or normal (**4**) CPP.¹ The pGGeC construct provides access to *ent*-CPP derived diterpenes through co-expression of class I labdane-related diterpene synthases. To provide similar access to *syn*-CPP (**3**) derived diterpenes we utilized the *syn*-CPP diterpene cyclase from rice (*Oryza sativa*) we have previously identified (OsCPS4)¹² and, based on our work with rAtCPS,⁴ constructed a pseudo-mature version suitable for recombinant bacterial expression (rOsCPS4, missing amino acids 2-70). Insertion of rOsCPS4 into pGG created pGGsC. To provide access to normal CPP (**4**) derived diterpenes, we returned to abietadiene synthase, which produces normal CPP (**4**) as an intermediate, and for which a D621A mutant (rAgAS:D621A) that no longer has class I activity

[i.e. only produces **4** from GGPP (**1**)] has been reported.¹³ Thus, rAgAS:D621A was inserted into pGG to create pGGnC.

To demonstrate the ability of these three pGGxC constructs to provide access to various stereoisomers of CPP for the production of a range of CPP derived diterpenes, pGGeC and pGGsC were individually co-transformed with previously described class I enzymes specific for the corresponding stereoisomer of CPP,⁷ while pGGnC was co-transformed with a D404A mutant of rAgAS (rAgAS:D404A) that only catalyzes the production of abietadiene (**5**) from **4** (i.e. only has class I activity).¹³ In each case this resulted in recombinant bacteria that produce the expected labdane-related diterpene (Table 1). Eight different hydrocarbon skeletal structures, with one as distinct double bond isomers, were produced, proving the utility of our modular approach for combinatorial biosynthesis of labdane-related diterpenes (Scheme 1). Notably, more than 1,000 natural products are derived from kaurene, with almost as many being derived from abietadiene, as well as ~500 from various pimaradienes.¹⁴ Thus, the diterpenes produced in this study are precursors to literally thousands of known compounds, and these results establish a foundation for biosynthetic production of elaborated labdane-related diterpenoid natural products. Both in providing substrates for characterization of downstream biosynthetic steps and for further extension of metabolic engineering efforts. In addition, the three pGGxC constructs reported here will be useful for characterization of novel class I labdane-related diterpene synthases. Finally, we expect that a similar approach will provide access to labdane-related diterpenes and diterpenoid natural products derived from other class II intermediates such as clerodadienyl diphosphate (i.e. by incorporating class II diterpene cyclases that produce such compounds¹⁵ into pGG), which will further broaden the reach of the modular approach to labdane-related diterpene biosynthesis demonstrated here.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

We thank Professor Robert M. Coates (Univ. Illinois) for authentic standards, Professor Rodney B. Croteau (Washington State Univ.) for the rAgGGPS and rAgAS clones, and Luke Lowry, Dana Morrone, and Meimei Xu for technical support. This work was funded by grants from the NSF (MCB-0416948) and NIH (GM076324) to R.J.P.

References

1. MacMillan, J.; Beale, MH. Diterpene biosynthesis. In: Cane, DE., editor. *Isoprenoids Including Carotenoids and Steroids*. 1st ed.. 2. Elsevier Science Ltd.; Oxford: 1999. p. 217-243.
2. Chang CYM, Keasling JD. *Nat. Chem. Biol* 2006;2:674–681. [PubMed: 17108985]
3. Croteau, R.; Kutchan, TM.; Lewis, NG. Natural products (secondary metabolites). In: Buchanan, B.; Grissem, W.; Jones, R., editors. *Biochemistry & Molecular Biology of Plants*. Am. Soc. Plant Biologists; Rockville, MD, USA: 2000. p. 1250-1318.
4. Pristic S, Peters RJ. *Plant Physiol*. 2007in press
5. Burke C, Croteau R. *J. Biol. Chem* 2002;277:3141–3149. [PubMed: 11733504]
6. Peters RJ, Flory JE, Jetter R, Ravn MM, Lee H-J, Coates RM, Croteau RB. *Biochemistry* 2000;39:15592–15602. [PubMed: 11112547]
- 7(a). Wilderman PR, Xu M, Jin Y, Coates RM, Peters RJ. *Plant Physiol* 2004;135:2098–2105. [PubMed: 15299118] (b) Morrone D, Jin Y, Xu M, Choi S-Y, Coates RM, Peters RJ. *Arch. Biochem. Biophys* 2006;448:133–140. [PubMed: 16256063] (c) Xu M, Wilderman PR, Morrone D, Xu J, Roy A, Margis-Pinheiro M, Upadhyaya N, Coates RM, Peters RJ. *Phytochemistry* 2007;68:312–326. [PubMed: 17141283]
8. Roy A, Roberts FG, Wilderman PR, Peters RJ, Coates RM. *J. Am. Chem. Soc.* 2007submitted

9. Reiling KK, Yoshikuni Y, Martin VJJ, Newman J, Bohlmann J, Keasling JD. *Biotechnol. Bioeng* 2004;87:200–212. [PubMed: 15236249]
10. Jackson BE, Hart-Wells EA, Matsuda SPT. *Org. Lett* 2003;5:1629–1632. [PubMed: 12735738]
11. Martin VJJ, Pitera DJ, Withers ST, Newman JD, Keasling JD. *Nature Biotech* 2003;21:796–802.
12. Xu M, Hillwig ML, Prisic S, Coates RM, Peters RJ. *Plant J* 2004;39:309–318. [PubMed: 15255861]
13. Peters RJ, Ravn MM, Coates RM, Croteau RB. *J. Am. Chem. Soc* 2001;123:8974–8978. [PubMed: 11552804]
14. Buckingham, J. *Dictionary of Natural Products* (on-line web edition). Chapman & Hall/CRC Press; 2002.
15. Hamano Y, Kuzuyama Y, Itoh N, Furihata K, Seto H, Dairi T. *J. Biol. Chem* 2002;277:37098–37104. [PubMed: 12138123]

Figure 1. Production of *ent*-kaur-16-ene (**6**) by *E. coli* transformed with pGGeC and pDEST15/rAtKS. Total ion chromatogram from GC-MS analysis of organic solvent eluant of Diaion HP-20 beads from a mixed phase culture (see Supporting Information).

Scheme 1.
Modular approach to labdane-related diterpene biosynthesis.

Table 1

Recombinant bacteria characterized in this study

Plasmid(s)	Product
pGGAS	abieta-7,13-diene (5)
pGGeC + pDEST15/rAtKS	ent-kaur-16-ene (6)
pGGeC + pDEST15/OsKSL5j	ent-pimara-8(14),15-diene (7)
pGGeC + pTH1/OsKSL6	ent-isokaur-15-ene (8)
pGGeC + pDEST15/OsKSL7	ent-cassa-12,15-diene (9)
pGGeC + pDEST15/OsKSL10	ent-sandaracopimaradiene (10)
pGGsC + pDEST14/OsKSL4	syn-pimara-7,15-diene (11)
pGGsC + pTH8/OsKSL8	syn-stemar-13-ene (12)
pGGsC + pDEST15/OsKSL11	syn-stemod-13(17)-ene (13)
PGGnC + pDEST14/rAgAS:D404A	abieta-7,13-diene (5)

E. coli containing the indicated plasmid(s) produce the indicated labdane-related diterpene, as identified by GC-MS based comparison to authentic standards, with production levels ranging from 5-100 μ g diterpene/L mixed phase culture (see Supporting Information).