What is the cause of this arteriolar contraction? Diathesis, psychic factors, overwork, sedentary occupations, obesity, endocrine changes, circulating pressor bases, have all had their advocates. Their very number is a confession of our ignorance. There is no doubt that high blood pressure is often familial. Lord Dawson found a percentage of school children had pressures above the normal,² but it is as yet too early to say whether any of them will eventually develop true hyperpiesia. Nervous causes and overwork certainly raise the pressure, but many men overwork and never suffer from hyperpiesia, and neurotic people often have low pressures. Sedentary occupations are blamed for many things: hyperpiesia is only one of them. Hyperpiesia and overweight may coexist, but it is doubtful whether the majority of obese persons have high pressures. There is no reason why circulating pressor bases should not be eliminated, since it is known that the kidneys remain normal until their arterioles become affected with those of the rest of the body.

Hyperpiesia will probably be found to be a deficiency disease. As noted by Ismail³ it has something in common with diabetes. In both diseases the usual time of onset is in early middle age; both run a prolonged course, have a certain familial tendency, and are frequently found together. As in diabetes, hyperpiesia is much more severe when it occurs in young people, and its course is more rapid: the so-called malignant hypertension of Keith and Wagener.⁴

A certain tension of the arterioles is necessary for everyone. This tension is unlikely to be maintained by the sympathetic nervous system, which is usually reserved for the rapid changes called for by the external environment. By analogy with other systems of the body,

it is probable that a chemical agent is responsible. It may well be that this substance, or substances, derived from protein, is normally present in excess, and the fine adjustment is made by a hormone, which converts it into something that can be excreted by the kidney. This view would accord with the work of Bain⁵ and Major,⁶ who have both found in normal urine substances with a pressor action; these were absent, or reduced, in the urine of those suffering from high blood pressure. If the hormone is inadequate, the arterioles will be over-stimulated, and the blood pressure will have to be raised to maintain the circulation through the capillaries. According to the extent of the hormone deficiency, various grades of hyperpiesia will result. Overwork and worry are important secondary factors, which will aggravate still further a state which is already abnormal.

Whether this view is correct, time will show. Hyperpiesia is a malady for which we have at present no cure; it will inevitably run its course. The potential hyperpietic, however, demands active treatment to prevent, if possible, the condition becoming established. In the later stages, also, much can be done to mitigate the disease, both by countering the secondary factors of overwork and worry, and by intensive courses aimed at relieving temporarily the stress upon the overburdened cardio-vascular system.

REFERENCES

- ¹ Weiss, S., Robb, G. P., and Blumgart, H. L.: *Amer. Heart Journ.*, 1928-9, iv, 664.
- ² Dawson, Lord: *British Medical Journal*, 1925, ii, 1161.
- ³ Ismail, A.: *Lancet*, 1928, ii, 275.
- ⁴ Keith, M.: Wagener, H. P., and Kernohan, J. W.: *Arch. Int. Med.*, 1928, xli, 141.
- ⁵ Bain, W.: *Quart. Journ. Exper. Physiol.*, 1914, viii, 229.
- ⁶ Major, R. H.: *Journ. Amer. Med. Assoc.*, 1924, lxxviii, 81.

TREATMENT OF HYPERPIESIA*

BY

W. EDGECOMBE, M.D., F.R.C.P., F.R.C.S.

PHYSICIAN, HARROGATE INFIRMARY.

The treatment of hyperpiesia must be based on our conception of the cause and the mode of production of the disease. Of the causes, diathesis—the inherent, often familial, tendency to the affection—cannot be removed. Psychical causes, such as overwork, anxiety, mental stress, and worry, are more or less amenable to modification, and must be remedied when possible. Inactive metabolism accompanying obesity and sedentary occupations is susceptible to suitable treatment. Endocrine factors—hyperadrenalism, subthyroidism, and disturbances of ovarian secretion, as at the menopause—can be partially controlled, though not with any great certainty, except in the case of myxoedema. Of the pressor bases concerned in the production of hyperpiesia we have little exact knowledge. Believing them to arise in the alimentary canal and to be derived from protein, we endeavour to control them by reducing the protein intake, and by promoting active elimination of waste products by the bowel, kidneys, and skin. Aperient and diuretic mineral waters and intestinal lavage by the Plombières douches are valuable means to this end. In illustration of the relation between intestinal infection and hyperpiesia the following two cases are of interest.

CASE I

Male, aged 49, when seen in 1913 had a blood pressure of 140/90, and was found on examination to have a large swelling in the region of the ascending colon which was dia-

gnosed as a diverticulitis. In January of the following year he was operated upon for suspected cancer of the bowel; the growth was found inoperable, and the abdomen was closed. In June his blood pressure was 145/95. In 1915 it had risen slightly to 150/95; in 1916 it was 160/100, reduced after a course of baths to 150/95. He was lost sight of until August, 1923, when he reappeared with a blood pressure of 230/130, and symptoms of slight weakness and dragging of the left leg. The swelling in the abdomen had almost disappeared, but was still just palpable. In seven years his blood pressure had risen from 160/100 to 230/130. A course of baths reduced it to 200/124. In 1924 his blood pressure was 220/125, reduced after a course to 200/105, and he died of cerebral haemorrhage in 1925. It is not unreasonable to infer that the chronic infection of the bowel was the cause of the hyperpiesia.

CASE II

Male, aged 48 in 1915, now aged 64. The records of his blood pressure date from 1916, and are seen to have gradually mounted year by year without any cause being discovered. Each year he had a short course of spa treatment. Between 1928 and 1929 there was a sharp rise, for which no explanation could be found. His blood pressure in May, 1929, was 230/110 before and 180/110 after a course of baths. In June, 1929, he had an acute perforation of the bowel, which was found on operation to be from a ruptured diverticulum. He made a good recovery. In December, 1929, his blood pressure was 190/100. An x-ray examination after a barium enema disclosed extensive diverticulitis of the large bowel. Since then he has remained in good health.

Consideration of the mode of production of hyperpiesia points the way to physical methods of treatment. The primary fault lies in the peripheral circulation; it is the narrowing of the arterioles and capillaries, and the consequent diminution of the total channel area available to the blood stream, that determines the high blood pressure; the enlargement of the heart is a secondary resultant phenomenon. Any measures that dilate the vessels and

* Read to the Section of Hydrology and Climatology at the Annual Meeting of the British Medical Association, Eastbourne, 1931.

widen the collective stream will delay the progress of the incipient disease and relieve temporarily when it is fully developed. For this purpose drugs such as the nitrites, iodine, acetylcholine, veratrin, the sulphocyanates, luminal, etc., are evanescent in their action, and though valuable as temporary measures and in the emergency of the hyperpietic crises, are of little avail as permanent remedies. There is no known drug that will effectively prevent the development of the disease. Vaso-dilatation of a more pronounced and enduring character can best be effected by physical modes of treatment. Immersion baths of mineral water at or above body temperature, effervescent carbonic acid baths, massage douches, Aix or Vichy, dry massage, electrical procedures such as the high-frequency current, and especially diathermy, will produce a peripheral vaso-dilatation and a fall of blood pressure. The result is not merely transitory, but endures after each bath, and is cumulative over a series of baths. To explain this effect it is assumed that not only are existing narrowed channels opened up in the peripheral circulation, but channels, as it were disused are called again into play, and it may be that new anastomoses are developed. Whatever the action, the effect is certainly more prolonged than that produced by drugs.

Early potential cases of hyperpiesia can be cured or arrested by timely treatment by physical methods. It is, of course, impossible to say whether a given case of suspected hyperpiesia would have completely developed had no treatment been adopted. Here is a case, typical of many, in which it is not unreasonable to assume that a cure was effected. (All the cases quoted below are believed to be examples of true essential hyperpiesia; that is to say, they were not cases of primary kidney disease with secondary high blood pressure; all of them have been under observation for a considerable period of years.)

CASE III

A young man, aged 35, in 1912 had a blood pressure of 160/110; his kidneys were sound, and there was no gross thickening of the vessels or enlargement of the heart. The causes were thought to be overwork, mental stress, and a sedentary life. He was given a course of spa treatment, with Aix massage douches and effervescing carbonated baths. In the following year his pressure was 150/105, and he had another course. In 1915, after a further course, his figures were 145/75. Since then he was seen at intervals of one or two years up to 1930, during which time his pressure remained steady at 120 to 130 systolic, and 70 to 80 diastolic. He is now perfectly well at the age of 55.

Another case, which has been under observation for twenty-five years, may be cited as an example of arrest of the disease by physical methods.

CASE IV

A woman aged 45 had, in 1906, a sustained blood pressure of 160/115, reduced after a course of Nauheim baths to 130/105. In 1907 the readings were 135/105 before and 115/95 after her course. Thereafter she was seen yearly up to 1930, having occasional courses of treatment by massage douches and Nauheim baths. Her blood pressure ranged from 140 to 200 systolic, and 90 to 100 diastolic, the last reading being 200/100 in 1930, reduced after a course of baths to 165/95. She is now 70, in good health, and active and energetic for her age. We believe that repeated courses of bath treatment have been instrumental in delaying the development of hyperpiesia.

As an example of the delaying effects of repeated courses of spa treatment on developing hyperpiesia, the following case may be mentioned.

CASE V

A woman aged 47 was first seen in 1919 with a blood pressure of 130/98—the systolic pressure not high, but the

diastolic raised so definitely as to create the suspicion of hyperpiesia. Her progress is epitomized in the following table.

Year	Blood Pressure	Treatment	Blood Pressure after Treatment
1919	130/98	Aix douches	130/90
1923	150/105	" .. Massage. Thyroid	130/90
1925	140/100	Vichy douche. Massage	—
1926, June ...	170/110	" .. "	150/108
October	160/100	" .. "	160/100
1927, April ..	180/100	" .. "	170/110
September	180/100	" .. "	170/108
1928	210/125	" .. "	190/110

Patient died in July, 1931, from cerebral haemorrhage.

That an annual course of spa treatment will afford temporary relief in advanced cases is shown by the two following examples. The patients were women, aged 64 and 58 respectively.

CASE VI

Year	Blood Pressure	Treatment	Blood Pressure after Treatment
1922	220/120	Aix douches. Liver packs	205/105
1923	240/135	" .. Nauheim baths	220/110
1924	240/120	Vichy douche. "	225/115
1925	225/120	" .. Massage	225/115
1926	240/120	" .. "	210/120

Patient died from cerebral haemorrhage in March, 1927.

CASE VII

Year	Blood Pressure	Treatment	Blood Pressure after Treatment
1927	220/110	Aix douches. Nauheim baths.	190/100
1928	220/140	Thyroid Aix douches. "	200/100
1929	230/140	" Diathermy	190/105
1930	240/120	" Nauheim baths	220/110

This patient is still living and in reasonably good condition. Both these patients felt themselves greatly relieved of their symptoms of fullness, headache, giddiness, and breathlessness after each course of baths, the improvement being maintained for some months.

The following is a far advanced case where life, we believe, was prolonged by repeated courses of bath treatment.

CASE VIII

A man, aged 60, had a slight left hemiplegia in March, 1927, from which he recovered almost completely. The following figures summarize his case.

Year	Blood Pressure	Treatment	Blood Pressure after Treatment
1927, April ...	220/160	Nauheim baths. Diathermy	240/130
1928, May ...	250/140	" .. "	240/130
September	240/140	Aix douches. "	210/140
1929, April ...	265/160	Nauheim baths. "	245/140

Patient died of cerebral haemorrhage in June, 1929.

Advanced cases require careful watching while under bath treatment. If the pressure falls rapidly and considerably, symptoms of lassitude, fatigue, and breathlessness may supervene, and the patient feels worse rather than better. This is an indication that the reduced head of pressure is insufficient to maintain an adequate circulation, that heart failure is impending, and the treatment should be stopped at once. If, as the pressure falls, the heart rate slows, or does not increase, and the patient feels relieved, all is well, and the treatment may be continued. A rise in heart rate with a fall in pressure is a danger sign. Also a considerable fall in the systolic without a corresponding fall in diastolic pressure—that is, a reduced pulse pressure—is an ominous sign of a failing left ventricle.

Memoranda

MEDICAL, SURGICAL, OBSTETRICAL

CATARRHAL INFECTION AND FORMALIN VAPOUR

The prevalence of catarrhal conditions in schools, especially during the early months of the year, is, I presume, one of the main problems which confront every medical officer. The loss of school time, the overcrowding of the school sanatoriums, and the strain imposed upon the nursing arrangements are harassing to masters and doctors alike. I would suggest that systematic spraying, punctiliously carried out on the plan laid down, has a marked effect in cutting down the incidence of catarrhal conditions, and in mitigating the virulence of the infection.

Some four years ago, with a badly infected boarding school of 140 boys, it was decided that the school should be sprayed, and a technique was evolved by the sister-in-charge (for which I can take little or no credit), which has been systematically carried out ever since. The spray solution is 10 per cent. formalin, used in a sovereign metal spray. The method, which is as follows, is carried out daily. All class rooms and studies are sprayed, including the tops of the desks, the backs of all pews in the chapel, the heads of all the beds and spaces between the beds in the dormitories, tooth-mugs, bathrooms, corridors, door handles, and dining tables. The spray, which is a fine one, is of course carried over a far larger area than the actual object on which it is directed. The time taken over this work is about one and a half hours a day.

I will now give the comparative figures of the numbers admitted to the sanatorium with catarrhal infection before and after this routine was established. In this group of

Before Spraying		After Spraying	
Year	Admissions	Year	Admissions
1925	90	1929	30
1926	59	1930	15
1927	78	1931	31
3 years ...	227	3 years ...	76

The year 1928 has been omitted, since spraying was not instituted at the beginning of the year, and the technique was not then standardized.

infections I have included feverish colds, influenza, bronchitis, and the pneumonias, and have excluded admissions from other causes.

The figures tend rather to mislead in that one is far more prone to admit the slighter cases into the sanatorium

when there is no pressure for space. I have not given the number of days occupied in hospital for a similar reason—namely, that when there is little pressure the tendency is always to retain cases longer. These figures are incomplete in a very important particular—that of the numbers who only attend the school surgery for the treatment of these conditions. Such figures I do not possess, but their number is also stated to have been markedly reduced.

There is no doubt in my mind nor in those of the sister and of the school authorities that the formalin spraying has been of very marked benefit. Further, this improvement has coincided with the institution of spraying and has continued for too long a period to be a mere coincidence. An interesting point is that for a short time the work was deputed by the sister to others. Soon after, an outbreak of colds and coughs started to spread through the school. It was discovered that the work had been very perfunctorily carried out.

I am less convinced of the efficiency of the spray in certain other infective conditions, such as follicular tonsillitis, which would appear not to be controlled. As to the exanthemata, I am not in a position to express any definite opinion. Their occurrences have only been slight during the period under review.

In conclusion, I would emphasize the apparent necessity for a punctilious routine, and I should be delighted to receive information of the experience of others along similar lines.

Banbury.

C. J. L. WELLS, M.B., B.Ch.Oxon.

British Medical Association

CLINICAL AND SCIENTIFIC PROCEEDINGS

NORTH-EAST ULSTER DIVISION

Tuberculosis of the Abdominal Lymphatic Glands

At a meeting of the North-East Ulster Division, on December 11th, 1931, with the chairman, Dr. BOYLAN, presiding, Mr. S. T. IRWIN, M.Ch., F.R.C.S., read a paper on tuberculosis of the abdominal lymphatic glands.

Mr. Irwin discussed first the incidence of tuberculous infection, and the arguments for and against the main portal of entry being the abdominal lymphatic glands. He described the work of Calmette in this connexion, and emphasized the importance of this probability as regards the diagnosis and prognosis in abdominal diseases. In the category of acute emergencies, especially in childhood, lymphadenitis occupied a prominent place, but he was not sure that this term was quite satisfactory, for all the types of this disease were not of definitely inflammatory origin, and in many cases no palpable swelling was present. Mr. Irwin then submitted a comprehensive summary of the literature, which indicated that these cases of lymphadenitis were due to the absorption of low-grade infections, caused not only by tubercle bacilli, but also by other organisms.

Anatomical investigations showed that the lymph vessels of the abdomen were grouped round the main arterial trunks, with glands situated at intervals, and that normally in all cases the largest glands in any group were placed proximally. Three groups were of special pathological importance: the one which accompanied the ileo-caecal artery and drained the lower six inches of the ileum, the ileo-caecal valve, the caecum, and the appendix; the one surrounding the superior mesenteric artery, and draining the jejunum and the upper part of the ileum; and the group which converged upon the coeliac axis,