from the use of pneumatic drills," thirty-one were compensated as disabled after the lapse of the twenty-six weeks' treatment received under the Sickness Insurance Act. According to Dr. Maria Seyring (*British Medical Journal*, 1930, ii, 1013), the maladies affecting pneumatic drill workers are angioneurosis of the capillary vessels of the fingers, leading to Raynaud's disease; atrophy of the muscles of the thenar and hypothenar eminences and interossei supplied by the median, radial, and ulnar nerves, the biceps, and infraspinatus; and injury to the elbow-joints and shoulder-joints. *Mutatis mutandis*, similar conditions are found among the comparatively few men employed on pounding-up machines (for removing creases in the leather when the upper is fixed to the insole, the machine vibrating 16,000 times to the minute). The German figures make out a case for inquiry into this form of injury, not only to ascertain the extent of the disability, but also to stimulate engineers to find a means external to the workman for carrying out this process.

The Home Office Committee of 1907 heard evidence from Middlesbrough about "slag cough," and described it as a "slight chronic inflammation of the bronchial tubes induced by the extremely fine state of division into which basic slag has to be ground. This illness has some distinguishing features, and may be held to be specific to the employment, but it cannot be included in the schedule for the reason that the evidence shows that it does not incapacitate for more than one or two days." No further evidence in this country has been advanced questioning the correctness of this view, which is, however, in striking contrast to the experience in Germany, both before and since the above opinion was expressed. The reason no doubt is that, in comparison with Germany, the preparation of basic slag in this country is on a very small scale. In years past I had been mystified by the number of cases of pneumonia referred to in the German Factory Inspectors' Reports from inhalation of basic slag dust. Thus, a report on basic slag works made by an official of the Imperial Health Office stated that, in 1892, in eighteen works 91.1 per cent. of the workers became ill; 56.4 per cent. of the cases were diseases of the respiratory organs, and of twenty-four deaths, nineteen were due to respiratory diseases. Again, in 1893, in twenty-one works sickness amounted to 108.9 per cent.; 54.5 per cent. of the cases were due to respiratory diseases, and of twenty-two deaths, eighteen were due to such diseases. As a consequence statutory regulations requiring monthly examination of all persons employed in basic slag works were enacted. Improvement by meticulous, locally applied, exhaust ventilation must have followed, as the figures of cases reported, thirty-six, and of those first compensated, nine, in the table for 1929 are reassuring.

Boilermakers' deafness is referred to by the 1907 Committee as "unquestionably an injury due to employment, and as widely prevalent among men working in incessant noise of the ship-building yard or the boilermakers' shop. It does not, however, prevent a man from continuing at his trade, and cannot, therefore, give rise to claims for compensation on the grounds of incapacitation"; but may it not lead to inability to secure work on the reopening of yards or shops after a temporary closing down? And here the German list shows one hundred and forty-eight referred, and fourteen included in the list of the more or less permanently incapacitated. Dr. Thomas Barr, from an examination of one hundred boilermakers, and a comparison of their power of hearing with that of a similar number of iron moulders and postmen in Glasgow, found that no one engaged in boilermaking for any length of time escapes impairment of hearing, and that three-fourths of the number could either not hear at all at a public meeting, or only with difficulty. The deafness is due to atrophy of the sensitive membrane in the cochlea and the organs of Corti, which constitute the receiving board of impressions of high-pitched notes.

Tropical diseases, spotted fever, and scurvy contracted by sailors is a remarkable inclusion, and, though the numbers (181 and 3 respectively) are small, they show it to have been worth while. Seamen are altogether excluded from our code, as the machinery set up (a certificate from

the certifying surgeon of the district in which the person was employed) is inapplicable.

Lastly, infectious ailments contracted by nurses, laboratory assistants, disinfectors, and the like, stand third and fourth respectively in the German list of cases reported and compensated for the first time. The possibility of including such cases was discussed in the 1907 Committee's report. It was held that while in individual cases—as, for example, that of a disinfecter contracting small-pox—the evidence in favour might be strong, yet "neither the wording of the Act, nor the character of the Third Schedule in its present form, nor the discussion of the Bill in Parliament, gives any indication that our legislation on workmen's compensation contemplates cases of ordinary illnesses contracted by chance through exposure to infection while in an employer's service."

The boldness, therefore, of the German Code, despite its apparent significant shortcomings, justifies my title of "New light on workmen's compensation." A word—even an apology—may be expected from me as one who signed the 1907 Committee's report; the Committee was tied, as it thought, by the quotation above, and the corollary that, before a malady could be scheduled it must last more than a week, must be gradual in its onset and not an accident, and must be so specific to the employment as to enable the disease to be established in individual cases. Intimate acquaintance with industrial disease since then makes me think these tenets should be revised, and that, instead of demanding as essential only some clear-cut distinguishable sign, more weight should be given to well-recognized cumulative effects on workpeople in the occupations.

The German figures, with the exception of silicosis, are likely to increase for some years as knowledge about the maladies becomes more widely disseminated. Already a steady rise each year has occurred in practically all those diseases which have been scheduled for the four years. The increase in the list was deferred, so it has been suggested, owing to the ignorance of medical practitioners about industrial diseases; the truth of this is shown by a decrease year by year in the number of cases marked "wrong diagnosis." This reproduces the experience of a gradual increase from year to year of not a few of the scheduled diseases in our list with the increasing knowledge of the conditions of compensation gained by workpeople and medical practitioners.

T. M. LEGGE.

ALCOHOL AND MENTAL DEFECT

ADDRESS BY DR. NORWOOD EAST

The following is an abstract of a paper read by Dr. W. Norwood East, one of H.M. Commissioners of Prisons, in opening a discussion on mental defectiveness and alcohol and drug addiction at a meeting of the Society for the Study of Inebriety, held on January 12th.

The distinction between the mentally defective person and the subnormal inefficient is empirical, and mental disorder may become superimposed upon mental defect; it has therefore seemed undesirable to restrict these observations to defectives who are certifiable under the Mental Deficiency Acts. It has been suggested that an increase in the number of mental defectives has taken place during the twenty years preceding 1929. Official statistics show that since 1915 the convictions for drunkenness have greatly decreased, and in 1929 (51,966) were less than half the convictions in 1915 (135,811). It appears, therefore, that the diminution in drunkenness is not yet reflected in the number of defectives. It may be that our experience in this matter is comparable to that of Norway, where it was found, many years ago, that a great diminution in drunkenness was reported without any decrease in the number of idiots.

There is little evidence to show that alcoholism is a direct cause of mental defectiveness in this country, if the latter term is used in the statutory sense. It is recognized that the effects of alcohol are most disastrous

in young people, but they seldom become so addicted to alcohol before reaching the age of 18 years as to produce the arrested or incomplete development of mind referred to in the Mental Deficiency Act, 1927. Nevertheless, intoxication in this country is not uncommon in adolescence, and in the three months October to December, 1925, 487 cases of drunkenness in persons under 21 were dealt with by courts of summary jurisdiction in England and Wales, and 114 of these were in persons under 18.

It is clear from the Report of the Royal Commission on the Care and Control of the Feeble-minded, the Departmental Committee on the Inebriate Acts, and other publications, that the term "mental defectiveness," prior to the Act of 1913, was frequently used in a general sense only. From 30 to 70 per cent. of the inebriates were said to be mentally defective. Recently, I found that of 698 persons convicted of various offences during the ten years 1921-30, and certified in prison under the Mental Deficiency Acts, 7.4 per cent. were, or had been, convicted of drunkenness. Dr. Douglas Turner found, at the Royal Eastern Counties Institution, less than 2.25 per cent. of adult defectives were intemperate; Dr. Rees Thomas, at the Rampton State Institution, found only 3.5 per cent. of the defectives could have been called at any time heavy drinkers; and Dr. A. E. Carver, approaching the subject as superintendent of an institution for alcoholism and drug addiction, found that in 432 adult male inebriates less than 1 per cent. were certifiable under the Mental Deficiency Acts. The Central Association for Mental Welfare also made inquiries for me of the local associations throughout the country. Of twenty-seven areas which sent in replies, eighteen were unaware of any cases of alcoholism among the ascertained defectives, and in other areas the number was negligible. The Central Association itself reported only one case. Clearly these results are in marked contrast to the earlier conclusions.

The Mental Deficiency Act, 1927, extended the conception of mental defect defined in the 1913 Act, and some workers consider it extended the use of the term "mental defectiveness" too far. There are others, however, who desire to extend the use of the term still further; I suggest that to do so would effect a return to the confusion of thought which was apparent before 1913, and also hamper medico-social research.

If it is agreed that the association of alcoholism and mental defectiveness is infrequent, it becomes a matter of interest to ascertain the reasons. Defectives are often seclusive—if not, their companions generally are younger than themselves and as impecunious; both factors tend to diminish convivial drinking. Intemperance is often the result of a complex habit, but many defectives, although suggestible, lack the powers of attention, receptivity, retention, and association of ideas which enable a particular stimulus to be followed by an elaborate habitual reaction. Defectives may be aware of their inferiority, and may drink to rid themselves of this disharmony, but it is probably true that the outlook of the adult defective towards the community leads him to consider that his units are superior to himself rather than to emphasize his inferiority in regard to them.

Defectives live, for the most part, in the present, free from past disappointments, unencumbered by imagination, and untroubled by the prospect of future hardships. They neither censure themselves nor anticipate the disapproval of others. The main reason for their sobriety appears to lie in the fact that their personalities are little affected by antagonistic complexes. Further, the observations of many years on delinquent defectives lead me to believe that, as a class, they are less homosexual than their fellows, although sexual offences are next in frequency to crimes of acquisition. If, in fact, the alcoholic releases repressed homosexual tendencies by his addiction, the defective may be seldom an alcohol addict for this reason.

Experience of delinquent defectives, delinquent inebriates, and inebriates who are not delinquents, leaves me in no doubt at all that the mental defective, as we see him in this country to-day, is seldom an alcoholic; that if he becomes an addict this is incidental to the

mental defect, and that the past estimates of the association of alcoholism and mental defect were excessive.

The part played by alcohol as a direct cause of entry to the social problem group of subnormals is complicated by the fact that the alcoholic is frequently the subject of other inherent causes of social inefficiency—a psychopathic make-up or asocial or temperamental limitation. Only broad suppositions seem justified in our present state of uncertainty concerning the types of case to be included in the group.

Prostitutes are a fairly well-defined subgroup of the social problem group, and Sanger has shown that 23 per cent. of the parents of 2,000 cases were intemperate. It is reasonable to assume that many of the daughters were mentally unstable, and perhaps their careers were influenced in consequence. Alcoholism may be also a direct cause of prostitution, but I have no reason to doubt the statements made to me by many prostitutes, during eight years' work among this class of prisoner, that they were entirely abstemious before they became public women, and of others that they were obliged to drink before they had sufficient assurance to accost possible clients.

A large number of subnormals lack will power and force of character, are temperamentally unstable, indifferent, incompetent, and peculiarly liable to solve their conflicts by alcohol. A tendency to addiction is also manifest in the cycloid temperament, and Kretschmer, discussing the schizoid temperament, considers its indifference is a common variant of affective insensibility, and that from the ranks of the indifferents are recruited the unstable, the gamblers, drinkers, harmless criminals, prostitutes, tramps, and others. It is particularly true of this group that in many of its units a vicious circle is easily induced. Alcohol restricts their inherent mental limitations and impels them to counteract this by further intoxication. It is also true that from among this group are recruited the majority of the alcoholists who appear repeatedly before the criminal courts charged with offences connected with drunkenness; many exhibit both intolerance to the action of alcohol and constitutional imbalance of the instincts and emotions.

There seems to be general agreement that it is a difficult matter, in our present state of knowledge, to assess the importance of alcohol as a direct cause of insanity. The general figure given, 10 per cent., according to some authorities, may be less. The Alcohol Investigation Committee of the Medical Research Council presented figures which indicate that the changes in the prevalence of alcoholism in women during the years 1913-18 had no observable relation to the number of cases of certifiable insanity. The convictions for drunkenness among women showed a continuous fall from 35,765 in 1913 to 7,222 in 1918, but the number of first admissions of women to lunatic asylums showed no such change; in 1913 it was 9,372, and in 1918 there was a slight rise to 9,726.

Alcoholism is both a symptom and a cause of mental disorder in many patients. General paralysis of the insane is a notable exception, and many paralytics drink to excess after the disease is declared and before they come under care. Alcohol taken in excess also by delusional patients, by manic-depressives in the exalted or depressed phases of the disorder, by epileptics, and by demented persons, sometimes leads to charges of drunkenness in the criminal courts. Although it is perhaps true to say of many insane persons that they are easily affected by alcohol, I am inclined to believe that some of the insane and introverted cases which come before the criminal courts show considerable tolerance of the drug.

INFLUENCE OF DRUG ADDICTION

The literature suggests that the estimation of the number of drug addicts is almost impossible to determine with any degree of accuracy in many countries. Perhaps mainly because the narcotic is administered in secret its effects may escape public notice, and facts are likely to be obscured, at least temporarily, if a different method of using the same drug comes into vogue, and if one drug is replaced by another. It is also extremely difficult to ascertain, with any degree of precision, the importance of ancestral drug addiction on the mental condition of the offspring. Chopra has come to the conclusion that heredity

does not play any part in inducing drug addiction. Sceleth and Kuh, as a result of experience gained in the treatment of 5,000 cases of morphinism, state that they have never seen an addict born an addict because his mother was an addict.

In this country drug addiction is admittedly infrequent, and its influence as a direct cause of mental defectiveness, subnormality, or insanity, for all practical purposes, is negligible. This appears to be the case, also, in Egypt, India, and British Malaya. The Indian Hemp Drug Commission, as the result of a searching inquiry, considered that only 7.3 per cent. of the admissions to mental hospitals in British India were attributable to hashish. Chopra was unable to obtain any evidence to show that opium by itself was responsible for mental derangements in India, and Samuel, at the Central Mental Hospital at Tanjong Rambutan, considered that opium, as a cause of insanity, was negligible.

There is, however, a large body of opinion in America, England, France, Germany, and elsewhere which considers that drug addiction is a sign of an inherent neurotic personality. The cases may be divided into two main classes: accidental addicts who develop the habit after the drug has been taken medicinally, and those, forming a larger group, who take narcotics because they possess psychopathic personalities. In some cases one drug is followed by another, or by other extravagances, during the course of the disorder.

New Zealand

[FROM OUR CORRESPONDENT IN WELLINGTON]

Vital Statistics

For the past year the general death rate was 8.56 per 1,000 mean population, which is believed to be the lowest in the world. The birth rate was higher than in the previous year, but the natural increase was only 1 per cent. Diseases of the heart and arteries were responsible for the highest number of deaths, approximately one-third of the total being due to such diseases. Cancer was the next principal cause of death. The number of deaths from tuberculosis continued to decrease, New Zealand, in comparison with other countries, having the lowest death rate from this cause. While infant mortality had slightly increased, the increase was to be found in the number of deaths of infants under 1 month. Septicaemia was the chief cause of maternal mortality.

Hospital Administration

Mainly as a result of the criticism of Dr. Campbell Begg the board of trustees of Wellington Hospital has adopted the principle of decentralizing the administration by appointing four standing committees, with subcommittees, to attend to matters of detail. Each standing committee has a clearly defined sphere of responsibility, but the four are co-ordinated on a basis that should prevent them from developing into water-tight compartments and should promote efficiency and economy. If the scheme operates successfully it may be given a wider application, and so result in national, as well as local, economy. Apart from the General Purposes Committee and the Fees and Relief Committee, there are a Medical Committee and a Nursing Committee. The Medical Committee, subject to the approval of the board, defines the sphere, privileges, and responsibilities of the different medical and surgical divisions and subdivisions, considers reports from their representatives, and co-ordinates their requirements. It considers requisitions for new medical and surgical equipment, and arranges staff meetings at which the professional work of the hospital, post-mortem findings, specimens removed at operations, and clinical cases presenting difficulty are considered. The Medical Committee also advises the board in regard to the standard of nursing service and

service given by house-surgeons, and, if requested, advises upon the suitability of applicants for medical and surgical appointments. In conjunction with the House Committee and the medical superintendent, it controls the clinical work of the hospital, chiefly with a view to the harmonious co-operation of the senior and assistant honorary staff with the stipendiary and nursing staff. It is a question whether a detailed constitution such as the foregoing, with rules and regulations, is to be preferred to reliance on a good understanding as between the sections of the staff of a large hospital, but there is no evidence that the two are incompatible, and it can be said with truth, so far as New Zealand hospitals are concerned, that hitherto the honorary staffs have had no proper status, and little or no direct contact with the board of management. The medical superintendents in the larger hospitals are now to be mainly administrative officers.

New Zealand Branch, British Empire Cancer Campaign

Research work is being carried on in the laboratories of the Otago University under Dr. Begg, and a biochemist has recently been included in the staff. The cancer clinics at the larger hospitals are doing excellent work, and valuable records are being obtained. Radium has been purchased in various instances where necessary and in suitable form, and the branch is co-operating with the Department of Health to ensure that radium is available only at suitable centres, and its use confined to those who are properly qualified in this branch of radio-therapeutics. The response of the people generally to appeals for financial help has been good, despite bad times and many other claims, and £50,000 is in hand; to this has to be added a further sum of £16,500. The intention of the branch is to build up a capital sum of £100,000 as an endowment for the cancer campaign in New Zealand. This branch was established at the time of Mr. Sampson Handley's visit to this country, and has done much already to improve the diagnosis and treatment of cancer cases, and to cause the public to take an intelligent, but not an alarmist, interest in the campaign against cancer.

Scotland

James Mackenzie Institute

The twelfth annual report of the James Mackenzie Institute for Clinical Research, St. Andrews, expresses indebtedness to the Medical Research Council for continuing the grant of £400 towards the upkeep of children's records, and also its grant for special research work. Expenditure during the year exceeded income by £78, and a strong appeal is therefore made for the raising of an endowment fund if the financial position of the Institute is to be secured and its important work continued. During the past year 97 new cases and 1,150 additional notes were added to the files, 795 specimens were examined in the bacteriological and chemical laboratories, and 472 films were exposed in the radiology department. Further progress was made in the inspection and analysis of the records of the Institute. At the end of the year 64.5 per cent. of the records were found to be complete, in that notes had been added in steady sequence; 13 per cent. needed to be brought up to date; and 22.5 per cent. were incomplete, largely because patients had been lost sight of. From a review of 584 records much valuable information was extracted, and the committee charged with this duty has been impressed by the abundance of material available in the files of the Institute. Its survey of the records being now, for the most part, complete, the committee proposes to investigate certain groups of