DISCUSSION OF A RECENT CASE

The above summary of etiology and diagnosis is given, since there will now be described a patient to whom none of the mentioned causes of origin appear to apply, and in whom diagnosis was made only during operation.

A man, aged 66, a road sweeper by occupation, presented himself in January, 1929, with the symptoms and signs of benign enlargement of the prostate. This was enucleated in the usual manner by the suprapubic route in a one-stage operation, use being made of a Pilcher bag, which was removed forty-eight hours later. This bag was maintained in position by a cord passing over a pulley to a weight of 1 lb. Five days later severe bleeding occurred, which was controlled by reintroducing the bag for several hours. The patient made a good recovery, and returned to his work as a road sweeper.

During the next two years he had slight dribbling when sitting down, and during this period developed a right inguinal hernia, which filled half the scrotum when standing.

In May, 1931, there suddenly appeared a swelling the size of a lemon, in the left side of the scrotum. It was painless, but three days after it appeared he presented himself for examination. This showed an ordinary right-sided inguinal hernia, and the left-sided fluid swelling, which was apparently attached to the cord and taken to be a hydrocele. His complaint of dribbling was noted, but assumed to be an unfortunate complication of the prostatectomy of two years before. Urethroscopy and cystoscopy were not done, but he was admitted for scrotal exploration and repair of the hernia and hydrocele.

At the operation the hernia was dealt with in the orthodox manner and the left-sided swelling approached by an inguinal incision. The swelling was seen to be no part of the cord, but was a fluid swelling arising by a pedicle one and a half inches wide from the corpus spongiosum, just in front of the bulb. A diagnosis of urethral diverticulum was made, and it was supposed that some distal obstruction accounted for it. A flexible bougie was therefore passed into the external meatus. It passed along the urethra with ease, and coiled itself up in the diverticulum. There was no urethral obstruction. The bladder was now opened suprapubically. Its neck presented the usual appearance at this time after prostatectomy. A sound entered at the internal meatus passed easily through, and could be manoeuvred either into the diverticulum or onward to the external meatus. A soft rubber catheter was now introduced along the length of the urethra and the diverticulum opened. The neck seen from inside looked just like the neck of a diverticulum of the bladder. It was lined by smooth epithelium, and the orifice was one inch long and one-third of an inch wide, being aligned with the urethra, in which could be seen lying the catheter. The diverticulum was cut away, enough being left to close the pedicle over the catheter by double breasting. The catheter was tied *in situ*. Into the bladder was placed a Marion's tube. The left inguinal incision, through which the above urethral operation was performed, was closed. Apart from a scrotal haematocoele the convalescence was uninterrupted, and the patient resumed his work as a road sweeper, on which he has since been engaged. When seen recently his only complaint was slight dribbling of urine, of which he loses several drops an hour.

OBSERVATIONS ON THE ABOVE CASE

Attention is drawn to the facility with which such a condition may be dealt with through a low inguinal incision. The lax scrotum resulting from the existing right-sided hernia made the exposure much easier. Further, this, like most other urethral operations, is best undertaken in association with suprapubic cystostomy, both for the additional angle of approach thus afforded to the lesion and for the suprapubic drainage during convalescence, which allows the urethra to remain dry throughout healing.

It appears advisable that disorders of micturition after prostatectomy should be investigated by urethroscopy and a urethrogram, as well as by cystoscopy.

When we come to consider the etiology of urethral diverticula in the light of this case it is by no means easy

to arrive at any conclusion. But it may be said outright that the case affords no support for the theory of congenital origin. Why should this man, aged 66, suddenly develop a urethral bulge because on the first day of his life there was a temporary rise of pressure in the bulbar urethra? Next, there was no doubt that the urethra distal to the diverticulum was completely patent at the time of the operation, and the obstruction of stricture or stone, which comes first to the mind, may be entirely dismissed.

It is a natural thing to try to associate this diverticulum with the prostatectomy he had undergone two years before. But it is very hard to do so. The prostatic cavity appeared normal, and was in no way connected with the diverticulum. Had it been, then the diverticulum would have presented in the perineum and not in the scrotum. There was no residual post-operative urethritis. Now it might have been that the passage of sounds after the prostatectomy had injured the urethra. But none of these had been passed for over two years.

I tentatively put forward the suggestion that there was urethral obstruction due to the right-sided inguinal hernia. The hernia pressed on the urethra just distal to the bulb. What would be the effect of this on a man with post-prostatectomy dribbling due to a damaged sphincter? It would help him to control the dribbling. But if he lay down or sat down, and the hernia went back up the inguinal canal, this additional control of micturition would cease. This is precisely what this man complained of. He could control his water when at work as a street sweeper, but not when at meals or in bed. Later, the pressure of the hernia on the urethra just in front of the bulb would cause obstruction, and behind this point the diverticulum appeared.

If this be the case, then we must regard urethral diverticulum as another of the possible complications of an inguinal hernia, as well as one of the differential diagnoses of hydrocele.

Finally, I would express my indebtedness to Mr. James McClure, F.R.C.S., of the London Temperance Hospital, by whose kindness I had the opportunity of taking care of, and operating upon, this patient, and with whose permission these notes are published.

HUMAN AND BOVINE TUBERCULOSIS

THE QUESTION OF MUTATION OF THE
TUBERCLE BACILLUS

BY

NATHAN RAW, C.M.G., M.D., M.R.C.P.

LORD CHANCELLOR'S VISITOR; LATE BRITISH MEMBER OF THE
INTERNATIONAL COMMITTEE ON TUBERCULOSIS

In my original paper in the *British Medical Journal* of 1903 I ventured on the following conclusions:

1. That there are two distinct varieties of tuberculosis affecting the human body, one produced by human tubercle, the other by bovine tubercle.
2. That human and bovine tubercle are separate and distinct.
3. That phthisis is caused by the human bacillus.
4. That gland, bone, and joint tuberculosis and lupus are of bovine origin and conveyed by milk.

The general opinion of bacteriologists with regard to specific micro-organisms is that they retain their characteristics under ordinary conditions, and are not subject to mutation. If, however, these micro-organisms are subjected to a new environment, in some instances a change takes place, which is usually called transmutation. I have for many years been of the opinion that bovine tubercle bacilli introduced into the human body in infancy or early childhood may, after many years' growth in new

tissues, undergo mutation and become human bacilli, capable of producing the usual lesions in the human body generally attributable to the human bacillus. Up to the present time I have had no opportunity of proving my theory, but the following case may throw some light on the subject. The case was investigated in great detail, and is as follows:

A man, aged 50, was seen by me at hospital in 1929, and gave the following history. No evidence of tuberculosis in family; was brought up on a farm, and when 5 years old developed glands in the neck, which were removed at the age of 7 by a distinguished surgeon, who said they were tuberculous. On one side of the neck the glands suppurated, and some small glands had remained indurated on the other side all his life. Two years ago the patient developed pulmonary tuberculosis, and at the same time the glands became active and swollen, finally suppurating. He had tuberculosis in both lungs of two years' duration, with profuse expectoration, which on bacteriological examination showed typical human bacilli. The pus from the neck glands was also carefully examined, both by culture and by animal inoculation. Guinea-pigs developed extensive generalized tuberculosis, whilst rabbits all gave negative results, and gained in weight. Cultures on 4 per cent. glycerin agar from the guinea-pigs showed a typical human growth.

DISCUSSION

The whole experiment showed that the bacilli found in such large numbers were of the human type, but the question of supreme importance is, were they originally human, or was the first infection at the age of 5 caused by milk containing bovine bacilli, which, after a long residence in a new host, had been transformed into a human bacillus? Tubercle bacilli recovered from the human cannot always be definitely classified as human or bovine. Many intermediate strains are met with, which I suggest may be stages of mutation in a new environment. After examination of hundreds of cases of tuberculosis in the laboratory I have never found human and bovine bacilli in the same body. I am certain that no change takes place in the three types of bacilli—namely, human, bovine, and avian—on ordinary media, as I have subcultured these three types continuously for twenty-two years, and though they have lost all virulence and toxicity, and are non-tuberculigenic, they still retain their distinct cultural characteristics. It is only when the tubercle bacilli are placed in a new host, as in the human body, that mutation occurs.

It is now generally accepted by clinicians that all cervical gland infections are caused by the bovine bacillus conveyed by milk, and the figures quoted by that eminent worker Stanley Griffiths are as follows. In children under 5 years of age he found about 90 per cent. due to bovine infection, but in cases between 20 and 40 years of age he found a much higher percentage due to human bacilli. Adults are not, as a rule, susceptible to bovine infections, and I suggest as an explanation of these figures that the original bovine bacillus is being slowly changed into an intermediate or even a true human bacillus.

PROBLEM OF PULMONARY TUBERCULOSIS

We have not yet sufficient knowledge to answer with any certainty two vital questions: (1) When does infection occur? (2) How does it take place?

We are generally agreed in thinking that infection takes place in early childhood, but many authorities deny the possibility of direct respiratory infection, and affirm that the bacilli are conveyed from the alimentary system to the bronchial glands, and so to the lungs. These questions are not decided. I have examined many thousands of cases of pulmonary tuberculosis, but have never been able to understand why they generally occurred between the ages of 18 and 30. If the infection has occurred in childhood it must be latent in the tissues. I would like

to suggest as a possibility that bovine bacilli introduced into the alimentary system in childhood may remain dormant in the mesenteric glands for a period of ten to fifteen years, in the meantime slowly assuming in human tissues the characteristics of human bacilli. By direct lymphatic extension, or by the lacteals, these bacilli may be conveyed to the bronchial glands, and so set up pulmonary tuberculosis.

If there is any substance in this supposition, it is obvious that, by directing our energies to the prevention of bovine infection through milk, we will not only eradicate all forms of surgical tuberculosis, but may also be assisting in the prevention of the more serious and fatal form of the disease popularly called "consumption."

BIBLIOGRAPHY

- Ferran, F.: Les mutations du bacille de Koch dans ses rapports avec la théorie de Mendel sur l'hérédité biologique et avec la prophylaxie de la tuberculose naturelle, *Monde médical*, 1929, xxxix, 77.
 Petroff, S. A.: Variability of Tubercle Bacillus, *New England Journ. Med.*, 1929, cc, 1148.
 Karwacki, L.: Morphologie des streptothrix obtenus par la transformation des bacilles tuberculeux, *C. R. Soc. de Biologie*, 1929, c, 214.
 Idem: Saprophytisme des bacilles tuberculeux, *Ibid.*, 1929, c, 1152.
 Idem: Conséquences pratiques du saprophytisme des bacilles tuberculeux, *Ibid.*, 1929, c, 1154.
 Felsenfeld, O.: Die Mutation des Tuberkulosevirus, *Med. Klin.*, 1929, xxv, 830.

Memoranda

MEDICAL, SURGICAL, OBSTETRICAL

CONGENITAL EPICANTHUS AND PTOSIS TRANSMITTED THROUGH FOUR GENERATIONS

Epicanthus is a subject on which little has been written during recent years. Foggin (*Ophthalmic Review*, vol. xx, 1901) classified congenital epicanthus as: (a) transient or temporary; (b) permanent (very unusual), whilst it is recognized that incomplete development of the nasal bones is the main factor in its production (thus acquired epicanthus may occur in necrosis of the nasal bridge from whatever cause). Usually, as the nose grows forwards and the face develops, the condition disappears.

The case in point, a boy aged 5, was very definite, and was associated with a marked congenital ptosis—in order to look at the ceiling it was necessary for the child to flex his neck on the back to the utmost. The condition was definitely traced back, through each generation, to his great-grandmother, and, further, in no case had the condition entirely disappeared in adult life—an uncommon event.

The members of the family affected are marked with an asterisk.

There has been no history of intermarriage in the family, and no evidence of mental defect. In the present case the only disturbing element to the child seems to be a liability to conjunctivitis. There is no defect in visual acuity.