fat overlying Sibson's fascia, and this provides an excellent plane in which the dissection can proceed once the opening in the deep cervical fascia has been enlarged.

These observations are founded upon twenty-six examples of "collar-stud" abscess treated by excision. Union by first intention occurred in all save two, both of these being in the supraclavicular region, and having advanced skin involvement. All the end-results have been satisfactory. I desire to make clear that in each case the glandular involvement was strictly limited. Indeed, it has been my experience that the tuberculous process is usually circumscribed in this condition. In all cases artificial sunlight treatment was commenced soon after operation, and where there were dental caries and diseased tonsils these received appropriate attention.

Finally, it is submitted that for those who must be educated and rendered fit for the battle of life with as little delay as possible, complete excision of a "collar-stud" abscess is a form of treatment which cannot be bettered.

REFERENCES

- ¹ Bailey, Hamilton: *British Medical Journal*, 1929.
- ² Calot: *Indispensable Orthopaedics*.

CEREBRAL ANEURYSM WITH RECURRENT LEAKAGE

BY

R. M. J. HARPER, M.A., M.D. OXON.

HONORARY MEDICAL OFFICER, NORTH DEVON DISPENSARY

(With Special Plate)

The patient, a single woman of 41, had enjoyed such exceptional health that, except for an appendicectomy during the war, she had never had to consult a doctor. Since October, 1929, she had complained of headaches before her menstrual periods, for which she took large doses of aspirin. Although it is said by her family that her manner was "strange" at these times, she carried on an active business life and never received medical attention.

March 7th, 1931.—Being in the pre-menstrual state, she complained of severe headache; later in the day her relatives, hearing a fall, found her unconscious on the floor. Seen by her doctor shortly afterwards, she was semi-conscious, pale, and collapsed. The power of speech was markedly affected, but there were no other physical signs. The reflexes were normal; the plantar response was flexor on both sides; the blood pressure was 150/90. A large fibroid was present in the lower abdomen. The patient made steady progress, and seemed likely to make a complete recovery.

April 1st.—She was getting out of bed and walking; her speech had improved greatly, but gave the impression that she talked with affectation.

April 4th.—She complained of severe headache, but as she was again in the pre-menstrual state this was not unusual. She described the headache as being frontal and occipital in distribution. She had been sitting in her bedroom, and later in the day she again fell; when found, she had vomited and was lying unconscious on the floor. She was removed to a nursing home.

The patient, on examination, was found to be stuporous and unable to co-operate in any way. She had lost the power of speech, and she held her head with her left hand. In the central nervous system the outstanding physical signs were rigidity of the neck muscles, and a right brachial monoplegia. In addition, it was noted that the arm reflexes were normal, the legs equal in tone, the knee-jerks absent, the ankle-jerks present, the plantar response flexor on both sides, and all abdominal reflexes absent. Kernig's sign was positive. The optic disks were within normal limits. The signs in the respiratory system suggested that a considerable quantity of vomitus had been inhaled into the right side of the chest. The temperature, after being subnormal, rose a few hours later to 99.8° F.

In view of the possibility of the condition being due to subarachnoid haemorrhage, a lumbar puncture was performed. The cerebro-spinal fluid, which was under pressure, was uniformly stained with fresh blood; after standing, the supernatant fluid was distinctly yellow in colour. These characteristics of the fluid, taken in conjunction with the above physical signs, led to a diagnosis of leaking cerebral aneurysm; and such localizing signs as there were pointed to the probability that the aneurysm was situated in the left Sylvian fissure. On the assumption that the attack of March 7th had a similar origin, the yellow staining of the supernatant fluid, indicative of altered blood pigments, would be consistent with this explanation.

During the next few days the condition of the patient improved; she was able to understand simple commands, but the power of speech and the right brachial monoplegia showed no improvement. The disks remained within normal limits, although the veins seemed to be slightly engorged. The pulmonary condition was actively progressive, and, in spite of the condition of the central nervous system becoming more normal, the patient died from broncho-pneumonia on April 10th, 1931.

FINDINGS AT NECROPSY

A post-mortem examination of the brain was carried out a few hours after death. There was some fresh blood in the main sulci, and about a dessertspoonful round the base of the brain. Over the left Sylvian fissure could be seen a clot of blood about the size of a shilling. Without further examination, the brain was allowed to harden in 10 per cent. formol saline. On dissection, the circle of Willis was followed out along the left middle cerebral artery, which led into a clot of fresh blood about the size of a walnut. When this was scraped away, there was revealed another tougher and more organized clot about a quarter the size of the other—obviously the result of an earlier haemorrhage. This second clot contained a small aneurysm at the origin of one of the branches from the middle cerebral artery. This aneurysm, shown slightly magnified, reproduced on the Special Plate, was grape-like in formation, and most probably of congenital type. A dissection of the other arteries of the brain showed that they were in a healthy condition, and revealed no further aneurysms. In the plate, indicated by a horse-hair, will be seen in the wall of the aneurysm a small opening (not an artefact) through which the leakage evidently took place.

Memoranda

MEDICAL, SURGICAL, OBSTETRICAL

A "CONTRACEPTIVE"

(With Special Plate)

It is contended by some birth-control authorities that the insertion of a Gräfenberg ring precludes all chance of pregnancy, except, perhaps, that of extrauterine gestation. An account of the following case may be of interest, and serve as a warning.

In February, 1931, Mrs. X, aged 36, married eight years, but with no children, consulted a London physician who is a protagonist of birth control, and was assured that if she had a ring inserted into the uterus no other means of contraception were necessary. Under gas and oxygen, a full-size Gräfenberg ring was planted in the uterus, and a few weeks later the patient proceeded to India with her husband. Throughout the early months of the year she suffered from irregularities, menorrhagia, and backache, with mastitis during her menstrual periods. In August and September no periods were seen, and there was constant low fever, with much right-sided pain in the abdomen. There was no vomiting, but a constant brown discharge. The question of pregnancy had never entered her mind, nor that of the medical attendant, but as her

R. M. J. HARPER: CEREBRAL ANEURYSM
WITH RECURRENT LEAKAGE

J. A. HENDERSON & F. PETTIGREW: HAEMORRHAGE
INTO SUPRARENAL CAPSULE

A. H. MORLEY, ARTERIO-VENOUS
ANEURYSM OF SUPERFICIAL TEMPORAL VEIN

H. G. WYATT: STERNAL FISSURE
IN A CHINESE