These experimental principles are of decided importance for the elucidation of some of the problems of hepatic cirrhosis in man, but the experiments themselves also afford an excellent lesson in the difficulty of directly applying results obtained in animals to human disease. A large number of cases of thrombosis of the portal vein are on record, in which, at necropsy, the liver appeared to be normal in every respect. In some of these the old thrombus had been canalized and the portal circulation to some extent restored. In others, in which this phenomenon had not occurred, the hepatic artery had apparently been able to take over the complete blood supply of the liver cells. If any atrophy of the human liver occurs following portal thrombosis, as after the ligation experiments in animals, the atrophic area must soon disappear without leaving a trace.

Hepatic Artery

The effects of experimental ligation of the hepatic artery or its main branches in animals are entirely different from those obtained with the portal vein, immediate necrosis of hepatic cells being the result, and not gradual atrophy. Cameron and Mayes (1930), who give a full historical account of the effects of occlusion of the hepatic artery in man and in animals, point out (as was perhaps not sufficiently realized by Rous and Larimore) that results depend to a considerable extent on the anatomical arrangement of the vessels, which may vary greatly in different animal species and even in man. Rapid occlusion of the hepatic artery in man may be associated with swift and fatal necrosis of the liver, whereas with the slower obstruction due to aneurysm no changes may ensue after ligation of the artery.

Hepatic Veins

The effects of obstruction of the efferent vessels of the liver have been studied by Bolton (1914), and by Bolton and Barnard (1931). Sudden total occlusion of the inferior vena cava between the liver and the heart as a rule causes death of the animal within a few hours at most. By an ingenious experimental method Bolton has been able to obstruct partially the inferior vena cava with a constricting band between the liver and the heart, thus inducing great venous congestion of the liver, ascites, and other signs of increased portal pressure. In his first paper Bolton describes the course of events in a monkey, which, after being subjected to a partial obstruction of the inferior vena cava, survived in good health for a year. When the animal was then examined a complete obliteration of the lumen of the inferior vena cava was found, resulting in widespread degeneration of the liver parenchyma, especially towards the centre of the lobules. In these degenerated areas fibrosis was absent. In Bolton's second paper (with Barnard), cats were used and killed at intervals of 4 to 113 days after partial obstruction of the inferior vena cava in the same situation. It was found in these experiments of widely differing duration that the essential lesion consisted of necrosis of the liver cells in the central part of the lobules, leaving islets of intact cells around the portal canals. The necrotic cells soon disappear, their place being taken by dilated sinusoids; no trace of fibrosis is ever observed. The absence of fibrosis in all of these experiments which involve obstruction to one or other part of the blood supply to the liver is a point of very great importance.

BIBLIOGRAPHY

- Abderhalden and Roske: Quoted by Ludwig, H., *Zeit. f. d. ges. exp. Med.*, 1931, lxxx, 36.
 Barcroft, J., and others: *Lancet*, 1925, i, 319; 1926, i, 544; *Journ. Physiol.*, 1925, lx, 443; 1927, lxxiv, 1.
 Barcroft, J.: *Ergebn. der Physiol.*, 1926, xxv, 818.
 Idem: *Veterinary Journ.*, 1931, lxxxvii, 466, 504.
 Barcroft, J., Nisimaru, Y., and Ray, G. B.: *Journ. Physiol.*, 1932, lxxiv, 44.
 Barcroft, J., Nisimaru, Y., and Puri, S. R.: *Ibid.*, 1932, lxxiv, 321.

- Bartlett, F. K., Corper, H. J., and Long, E. R.: *Amer. Journ. Physiol.*, 1914, xxxv, 36.
 Bauer, A.: *Journ. d'Anat. et de Physiol.*, 1909, xlv, 1.
 Binet, L.: *La Rate: Organe Réservoir*, Masson, Paris, 1930.
 Bradley, C. C.: *Journ. Anat. and Physiol.*, 1908, xliii, 1.
 Burton-Opitz, R.: *Pflüger's Archiv. f. d. ges. Physiol.*, 1909, cxxix, 189.
 Cantlie, J.: *Journ. Anat. and Physiol.*, 1898, xxxii (*Proc. Anat. Soc. Great Britain and Ireland*, p. 4).
 Cohnheim, J.: *Virchow's Archiv.*, 1867, xl, 1.
 Copher and Dick, B. M.: *Arch. Surg.*, 1928.
 Cullen, T. S.: *Ibid.*, 1925, xi, 718.
 Dévé, F.: *Normandie Médicale*, Rouen, March 1st, 1908.
 Dick, B. M.: *Edin. Med. Journ.*, 1928, N.S., xxxv, 533.
 Dock, G., and Warthin, A. S.: *Amer. Journ. Med. Sci.*, 1904, cxxxvii, 24.
 Flower: *Med. Times and Gaz.*, 1872; *Brit. Assoc. Rep.*, 1872, p. 150 (quoted by Bradley, C. C.).
 Gilbert, A., and Villaret, M.: *Arch. de Méd. exper. et d'Anat. pathol.*, 1906, xxi, 373.
 Glénard, F.: *Lyon Méd.*, 1890, xlv, 5, 80, 115, 189, 259; *Bull. et Mém. Soc. méd. des Hôp. de Paris*, 1901, S. 3, xviii, 386.
 Grab, W., Janssen, S., and Rein, H.: *Zeit. f. Biol.*, 1929-30, lxxxix, 324.
 Herringham, W. P.: *St. Bart's Hosp. Report.*, 1905, xli, 15.
 Loeb, O.: *Arch. f. exp. Pathol. u. Pharmacol.*, 1907, lvi, 320.
 Looten, J.: *Journ. de l'Anat. et de la Physiol. norm. et path.*, 1908, lxxxvii, 110.
 McIndoe, A. H., and Counseller, V. S.: *Arch. Surg.*, 1927, xv, 589.
 McNee, J. W.: *Lancet*, 1931, i, 951, 1009, 1063.
 Nisimaru, Y., and Steggerda, F. R.: *Journ. Physiol.*, 1932, lxxiv, 327.
 Pick, E. P.: *Wien. klin. Woch.*, 1931, xlv, 985.
 Rolleston, G.: *Brit. Assoc. Rep.*, Notes and Abstracts, 1861, p. 174.
 Roy, C. S.: *Journ. Physiol.*, 1881, xii, 203.
 Rous, P., and Larimore, L. D.: *Journ. Exper. Med.*, 1920, xxxi, 609.
 Ruysch, F., quoted by Gray, H.: *On the Structure and Use of the Spleen*, Parker, London, 1854.
 Schäfer, E. A., and Moore, B.: *Journ. Physiol.*, 1896, xx, 1.
 Schiff, M.: *Leçons sur la physiologie de la digestion*, II, 1867.
 Sérégé, H.: *Journ. de Méd. de Bordeaux*, 1901, xxxi, 271, 291, 321.
 Idem: *Compt. Rend. Soc. Biol.*, (Série II), 1902, liv, 201, 300.
 Steenhuis, T. S.: *Proefschr. Rijks-Universiteit te Groningen*, 1911 (quoted by Rous and Larimore).
 Stukeley, W.: *On the Spleen, its Description and History, Uses and Diseases*, London, 1722.
 Tait, J., and Cashin, M. F.: *Quart. Journ. Exper. Physiol.*, 1925, xv, 421.
 Toldt, C., and Zuckerkandl, E.: 1876 (quoted by McIndoe and Counseller).
 Warthin, A. S.: *International Clinics*, 1910, iv (20th Series), 189.
 Wells, H., Gideon, and Hedenburg, O. F.: *Journ. Infect. Dis.*, 1912, xi, 349.
 Wertheimer, E., and Le Page, L.: *Compt. Rend. Soc. Biol.*, 1896, x, 950.
 Winternitz, M. C.: *Johns Hopkins Hosp. Bull.*, 1911, xxii, 396.

RECENT ADVANCES IN THE TREATMENT OF FRACTURES*

BY

R. WATSON JONES, B.Sc., F.R.C.S., M.Ch.Orth.

LECTURER IN PATHOLOGY OF ORTHOPAEDIC CONDITIONS, LIVERPOOL UNIVERSITY; HONORARY ASSISTANT SURGEON, LIVERPOOL ROYAL INFIRMARY AND SHROPSHIRE ORTHOPAEDIC HOSPITAL

(With Special Plate)

It is well recognized that in the treatment of fractures faulty alignment, over-riding, and all forms of malunion give rise to serious disability. But other equally serious and equally common causes of disability have not received the recognition they deserve. Too commonly the surgeon's attention has been focused on the bones, to the exclusion of other parts of the limb. In the determination to avoid malunion, joints have been unnecessarily immobilized, and irrecoverable stiffness has been allowed to develop. Colles's fractures may still be seen, perfectly reduced and united in excellent position, but associated with serious and frequently permanent stiffness of the metacarpo-phalangeal joints. Very frequently these disabilities, which are not due to the primary injury, but appear during treatment and in consequence of it, are even more serious than the disability which the treatment sets out to cure.

MAIN OBJECTS OF TREATMENT

It is of course essential, not only that a fracture shall be perfectly reduced, but that the fragments shall be completely immobilized for a period of sufficient duration

* Paper read at the Liverpool Medical Institution, March, 1932.

to secure consolidation. There is no more constant factor in the establishment of non-union than inadequate fixation, or fixation for too short a period. But it is quite wrong to assume that the immobilization must of necessity give rise to muscle wasting, decalcification of bone, stiffness of joints, and circulatory disturbances with recurring oedema. These are the consequences of disuse, and are to be avoided by instituting functional treatment from the very first days of the fracture. There are therefore three aspects of the treatment of a fracture, all of equal importance, and the neglect of any of the three will give rise to permanent disability:

1. Reduction of displacement.
2. Complete immobilization of the fragments.
3. Functional activity.

In order to maintain functional activity it is important that fixation should involve no more of the limb than is necessary. Joints which need not be fixed must be definitely moved, and muscles, even in the part of the limb which is immobilized, should be in constant activity. Carr's splint was long ago discarded in the treatment of Colles's fracture because the palmar bar prevented flexion movements of the fingers. But even when gutter splints were used with the fingers and palm free, the patient was still much too afraid to practise exercises unless he was actually stimulated into doing so. Moreover, unless he was instructed otherwise, he would most carefully protect the limb in a sling, and allow stiffness of the shoulder-joint to appear. Exercises can be secured with greatest certainty if the patient is able to follow some occupation which demands the use of his injured limb, so that his exercise is constant and subconscious. This object can be attained in Colles's fracture by using an unpadded plaster cast, which leaves the fingers and palm entirely free, allows the patient to dress normally, with the injured limb through the coat sleeve, and renders the support of a sling unnecessary.

ACTIVE MODERN METHODS

In consequence of these advances the "massage" departments of hospitals will undergo a great transformation during the next few years. A patient with a fractured limb can only be persuaded that it is safe to use it if he is taught occupational activity under supervision. In my own physiotherapy department at the Liverpool Royal Infirmary the number of patients who are actually being massaged is reduced to a negligible minimum. But every patient goes into the department as soon as x rays have shown that the fracture is perfectly reduced. Visitors to the department still see a great hall filled with patients and masseuses. But the patient is not lying on a couch being lulled to sleep; he is being galvanized into activity. The man whose Pott's fracture was reduced yesterday is to-day being taught to walk. The woman who sustained a Colles's fracture yesterday is to-day learning to make beds and peel potatoes. Men are sawing wood and hammering nails. The man whose occupation involves the climbing of ladders, climbs ladders in the department from the early days of his injury. The painter paints, the sweeper sweeps, the window-cleaner cleans windows, and as far as possible individual occupational activity is maintained throughout the period of immobilization. The patient with the fractured spine, whose trunk is encased in plaster, is taught exercises for the erector spinae, which are far more effective than faradic stimulation, even if such were possible. Every patient whose knee-joint is immobilized, whether he walks or not, is taught tonic exercises for the quadriceps, and if a patient is due for admission to hospital for a knee operation, he is first taught quadriceps drill until he is so expert in his exercise that he can practise it from the day of operation onwards.

In short, modern treatment anticipates the complications of muscle wasting, joint stiffness, and mental fear, and instead of treatment being prescribed after these conditions have developed, steps are taken to prevent them from appearing.

FRACTURES OF CARPAL SCAPHOID

Although the principle of absolute immobilization with functional activity has been of importance in accelerating recovery in all fractures, there is no injury in which its application has more profoundly influenced the prognosis than in fractures of the carpal scaphoid. The blood supply of the scaphoid is such that a fracture through the waist of the bone renders the proximal half entirely avascular. Union depends on the development between the two fragments of fine capillary vessels which will be ruptured with the slightest shearing movement. Perfect immobilization is therefore necessary, and must be maintained in recent fractures for two to four months, and in old ununited fractures for six to twelve months. But, however perfect the immobilization, without functional activity the fracture will still fail to unite, because the decalcification of disuse prevents consolidation. Böhler's¹ dorsal plaster cast, applied directly to the skin, has solved the problem. With such a cast we have had patients at work not simply as university professors and company directors, but as labourers, railway stokers, ship's riveters, painters, and decorators. Since this work was first published two and a half years ago, I have treated ninety-seven recent fractures without a single non-union, and twenty-two old ununited fractures of one to five years' duration with only a single failure (Plate, Fig. 1). The one failure was a case in which I hoped to accelerate union by using a bone graft as well. In many of the old cases it is necessary to freshen the fractured surfaces either by opening the joint or by subcutaneous drilling. I also believe that the ununited fracture unites with greater certainty if the wrist is held in radial deviation and not in ulnar deviation as advised by Böhler.

FRACTURES AND FRACTURE DISLOCATIONS OF SPINE

The method of manipulative reduction of crush fractures and fracture dislocations of the spine, which I described in 1931,² has enabled us to apply to these injuries the principles of treatment which have been enunciated—namely, reduction of the fracture, absolute immobilization, and continued functional activity. Hyperextension of the spine, which is essential for reduction, is effected in its simplest form by laying the patient prone between two tables, one higher than the other. A plaster jacket is applied, and in uncomplicated cases ambulation is commenced within a few days. Records are now available of fifty cases which have been treated in this way, and it has been found that in all cases the patient's own body weight is sufficient to reduce even the most severe crushing injuries (Fig. 2). No anaesthetic is necessary, and in the only case in which I was tempted to use general anaesthesia—a severe fracture dislocation of the dorsal spine with, so it transpired at post-mortem examination, complete transection of the cord—the patient died before recovering consciousness. In no other case has there been an operative mortality. No ill effects have been observed from early ambulation and weight bearing, and in no case has there been recurrence of deformity whilst the plaster jacket was worn—even in lumbar injuries. It has, however, been observed that a completely crushed vertebral body consolidates slowly after reduction, and in one such case, involving the fourth lumbar vertebra, though reduction was still complete after four months' ambulation in a plaster jacket, deformity recurred during the fifth month, despite the wearing of a posterior spinal support. In view of this, the simple posterior spinal support is no longer

relied on, and although four months is still regarded as an adequate immobilization period for the average crush fracture, in severely comminuted fractures the plaster jacket is retained for six months. Moreover, the fact that consolidation is so slow would suggest that the deformity is at least capable of improvement if a hyperextension plaster is applied as late as three or four months after the injury. Fracture-dislocations with paraplegia are always reduced at once, because, although recovery is impossible if the cord is transected, there can be no doubt that a recoverable lesion of the cord frequently becomes irrecoverable owing to continued compression and anaemia. Immediate relief of pressure by hyperextension is the only measure which can be of value, and has completely replaced laminectomy in the treatment of this complication. Five cases of total paraplegia have fully recovered, and in one case a fourth lumbar root paralysis completely recovered within twelve months of ambulation.

FRACTURES OF NECK OF FEMUR

The Whitman plaster remains the method of choice for fractures of the base of the neck of the femur, and bony union can always be secured here in perfect position. But the trans-cervical and subcapital fracture is an entirely different problem, and in at least 40 per cent. of such cases the Whitman method fails to secure bony union. Even when union is firm there is frequently shortening of the neck with coxa vara, and it has been necessary to retain the complete plaster for a much longer period than was originally described. Not only does a patient over 60 years of age strongly protest against three to six months' immobilization in plaster, but the knee-joint is so frequently arthritic that it becomes stiff and painful, and he rightly claims that although one disability may have been cured, another has been produced. Smith-Petersen³ has introduced a three-flanged nail, which is driven through the neck from the trochanter. The length of the nail prevents angulatory displacement and over-riding, and the three flanges prevent rotational displacement. The operation is somewhat difficult and is not to be undertaken lightly, but a surprising degree of fixation is secured when the nail is in position. No other immobilization is necessary, a bandage is applied, and the patient sits up in bed moving his knee and hip from the first day. Provided that perfect asepsis is maintained, and a stainless steel nail used, no absorptive changes occur, and the nail is still as tight in the bone four months later as it was on the first day of operation. For the first time a method of treatment of fractures of the neck of the femur has been described which fulfils the necessary criteria of perfect reduction, absolute immobilization, and continued functional activity. In my own series, bony union was secured in seven out of eight* cases, despite the fact that activity in bed was never restricted, that there was no risk of hypostatic pneumonia, no discomfort was suffered, full knee movements were practised, and normal muscle tone maintained (Fig. 3). The fact that it was not necessary several months after fracture to embark on a long course of physiotherapeutic treatment for the recovery of knee movement reduced the incapacity period by half. One patient actually walked (without permission) from the third week onwards, and suffered no ill effect. There was no mortality, although one patient was 80 and five were over 50 years of age. The shock of the operation was never a serious factor, and there can

* In the case which failed a plated nail had been used—not one of stainless steel. Four weeks after operation there was x-ray evidence of absorptive changes round it, and when explored it was found to be quite loose. There was no infection, and since in all cases where a stainless nail is used it is impossible to extract it with ordinary surgical tools even several months after operation, one must assume that the quality of the metal has accounted for local irritative effects, with resulting hyperaemia and decalcification. This difference in reaction between plated metal and stainless steel has been confirmed experimentally.

be little doubt that an old and fragile patient is better able to withstand a twenty-five-minute operation under spinal anaesthesia than three to six months' fixation in a complete plaster spica.

OVER-RIDING FRACTURES OF TIBIA AND FIBULA SHAFTS

Fractures of the shafts of both leg bones unaccompanied by over-riding offer no difficulty, whether in reduction, fixation, or early functional treatment with weight bearing. But if there is over-riding, due either to total disengagement of the fragments or to the obliquity of the fracture line, treatment is much more difficult. It is possible by the "gravity method" to reduce recent fractures, and, if an expert assistant is available, to put them up in plaster. The assistant must be prepared, however, with a very insecure grip on the patient's heel, to maintain heavy traction for ten or fifteen minutes, to prevent antero-posterior and lateral angulation, to keep the foot in the same plane of rotation as the patella, and to dorsiflex the ankle to a right angle. Moreover, assistants are not always available, they are subject to fatigue, and fractures are not always seen within a day or two of injury. For these reasons a very simple piece of apparatus has been devised,⁴ which is portable, can be clamped on to any table, allows the operator to be entirely independent of assistants, and will maintain heavy traction long enough to put the limb up in plaster (Fig. 4). An os calcis pin is used, but is removed as soon as the plaster is dry, so that there is no risk of infection along its track. The pull behind the ankle-joint automatically dorsiflexes the foot to a right angle, and at the same time the traction maintains full length and correct alignment. The limb is absolutely rigid, and the operator may apply whatever lateral pressure is necessary to impact the fragments. With practice it is easy to secure just as perfect an anatomical reduction as could be effected by an open operation. An unpadded plaster cast, extending from the toes to the mid-thigh, is applied.

Weight bearing may be commenced in three to four weeks if the fracture line is horizontal, and if the plaster was applied before the onset of swelling, so that it still grips firmly. With oblique and spiral fractures weight bearing should be deferred for six weeks. If the limb is severely swollen when the fracture is reduced, the plaster will be very slack in four weeks' time, and weight bearing must be deferred until a new unpadded plaster has been applied. When weight bearing is commenced a sorbo rubber heel is incorporated in the plaster, and a cloth boot made to cover the whole foot. The sorbo rubber adds spring to the gait and gives very considerable comfort by relieving the jar of weight bearing. A patient who has once walked with a sorbo heel plaster will always insist on the rubber being incorporated in any later plaster which may be applied. Moreover, the fact that he may dress normally, covering the plaster with stocking, boot, and trouser, gives him a mental assurance and a protection from inclement weather which will enable him to carry out the stipulated minimum of five miles walking a day.

As a rule, in eight weeks the fracture is firm enough to allow the continuance of walking with a below-knee plaster, while at the same time knee movements are recovered. No massage or physiotherapy is used. The fracture should be firmly consolidated in ten weeks, and although manipulation of the ankle for adhesions below the external malleolus is often indicated, recovery should be complete in three to four months, with a limb indistinguishable from normal.

REFERENCES

- ¹ Lorenz Böhrer: *Treatment of Fractures*, 1929, p. 98.
- ² Watson Jones: *British Medical Journal*, February 21st, 1931; *Journ. of Bone and Joint Surgery*, xiii, No. 2, 383.
- ³ Smith-Petersen: *Arch. of Surg.*, November, 1931, xxiii, 715.
- ⁴ Watson Jones: *British Medical Journal*, November 14th, 1931, 898.

R. WATSON JONES: RECENT ADVANCES IN THE TREATMENT OF FRACTURES

FIG. 1.—(a) Dislocation of carpal semilunar and fracture of the scaphoid. (b) X-ray four months after manipulative reduction. The patient had returned home, and the plaster cast had unfortunately been removed two months after injury, with the result that the scaphoid has failed to unite. (c) After six months' further immobilization, the scaphoid has firmly united by bone. This case illustrates the futility of immobilization for too short a period.

FIG. 2.—(a) Crush fracture of fourth vertebra with total paraplegia and fourth lumbar root palsy. (b) After reduction by hyperextension and plaster. The paraplegia recovered in a few weeks, and the root palsy recovered completely in twelve months.

FIG. 4.—Tibia traction apparatus. Fracture of both leg bones fully reduced, and held ready for application of plaster. No assistant is necessary. The apparatus can be applied to any table. (Made by F. H. Critchley, Liverpool.)

FIG. 3.—(a) Fracture back of femur in a patient of 78. (b) After Smith Paterson nailing operation. (c) Six months after operation. Firm bony union in perfect position, with normal movements of knee-joint and normal muscular development.