3. The asthenic type of stomach is infrequently the seat of duodenal ulceration ; but otherwise duodenal ulcers occur in the hyposthenic, sthenic, and hypersthenic types in about the same proportions as these types occur in a hundred normal individuals.

4. The average test meal curve is higher in cases of duodenal ulceration than in normal individuals.

5. The curve of gastric acidity steadily increases from being lowest in the asthenic to a maximum in the hypersthenic type.

I am much indebted to Dr. Charles Bolton for his help and suggestions in carrying out this investigation, to Dr. G. W. Goodhart for permission to use the test meal curves, to Dr. Salmond and the staff of the Radiographic Department of University College Hospital for the use of their radiograms, and to the physicians under whose care the patients were treated for permission to use the records.

REFERENCES

- ¹ Moynihan, Lord: *British Medical Journal*, 1923, i, 221.
- ² Hurst, A. F.: *Gastric and Duodenal Ulcers*.
- ³ Redding, Magnus: *X-ray Diagnosis*.
- ⁴ Mills, R. W.: *Amer. Journ. Roentgen.*, 1917, iv, 155.
- ⁵ Hurst, A. F.: *Lancet*, 1922, ii, 1369.
- ⁶ Carman, R. D.: *Roentgen Diagnosis of Diseases of the Alimentary Canal*, p. 135.
- ⁷ Hunter, Donald: *Quart. Journ. Med.*, 1923, xvi, 95.
- ⁸ Ryle, J. A.: *Lancet*, April 4th, 1925.
- ⁹ Idem: *Ibid.*, February 13th, 1932.
- ¹⁰ Bennett and Ryle: *Guy's Hospital Reports*, 1921, lxxi, No. 3, 317.

EXCRETION UROGRAPHY *

BY

REGINALD T. PAYNE, M.B., B.S., F.R.C.S.

(With Special Plate)

There are few parts of the human body where localization of the extent and nature of diseased processes can be carried out with such a high degree of accuracy as in the urinary tract. Certainly in no other system does accurate localization so constantly indicate the possibilities of surgical treatment. For this accuracy of diagnosis urology has owed much to radiology, and without its help advance would have been impossible. In this respect the year 1906 was a memorable one, for it was then that the first successful pyelography was carried out by Professor Voelker upon Alexander von Lichtenberg. Since that time there have been many modifications and improvements in the technique of carrying out instrumental urography. But all these methods have involved cystoscopy, and in the case of ureterography and pyelography have involved, in addition, the passage of ureteric catheters. The main improvements have been technical ones, and those relating to the type of material used as opaque media. With these instrumental methods, however, great advances have been made in the diagnosis of affections of the urinary tract. In the kidney, pyelography has permitted the localization of stone, and the demonstration of the configuration of the calyces and pelvis in inflammation, hydronephrosis, and new growth, to mention but a few examples. In the ureter, its course, shape, and relation to doubtful shadows in the neighbourhood have been demonstrated. In the bladder, cystography has been of great use in the demonstration of diverticula. Such radiographic methods of demonstration in the urinary tract are instrumental ones. Excretion urography, on the other hand, is the demonstration of the urinary tract by the intravenous injection of drugs which are excreted in a concentrated form in the urine—drugs which are, in addition, opaque, and in consequence of this property the outline of the urinary tract can be demonstrated during their excretion.

* Abstract of Hunterian Lecture delivered before the Royal College of Surgeons of England on February 3rd.

HISTORICAL ASPECTS

At about the same time as the introduction of instrumental pyelography Voelker and von Lichtenberg attempted the experimental production of excretion urograms by the use of the colloidal heavy metals, but found that these were too toxic. Following this, and partly due to the successful development of instrumental methods, further investigations were in abeyance for some years. In the year 1923 Osborne, Sutherland, Rowntree, and Scholl, by means of the intravenous injection of a solution of sodium iodide, were able to obtain quite satisfactory cystograms, together with a suggestion of opacities in the region of the renal pelvis. They thus proved that excretion urography was a possibility, and that success awaited the discovery of an appropriate contrast medium. Following this, a large amount of experimental work was carried out in animals in the search for a contrast medium which would be excreted in sufficient concentration to give good pictures. It was, however, Roseno who advanced the problem a step further. This worker conceived the idea of using a combination of urea and sodium iodide, known as pyelgnost. After intravenous injection of this drug radiographic shadows of the urinary tract were obtained, but frequently toxic effects were produced, and in many cases the density of the resulting pictures was insufficient for clinical purposes.

Finally, the greatest stimulus to renewed effort was the success which attended Graham's discovery of cholecystography. The search for an adequate medium for urography came in 1929 as the result of much painstaking work in chemical synthesis, biological assay, and clinical trial.

CHOICE OF DRUGS

From the chemical standpoint the credit of first synthesizing satisfactory drugs for excretion urography belongs to Professor Binz of Berlin, a pupil of Ehrlich. With the help of Professor Raeth some seventy-three drugs related to the pyridine series and containing iodine were synthesized. After animal experiments, the clinical investigations with certain of these drugs were carried out by Professor Alexander von Lichtenberg and his assistant Swick at St. Hedwig's Krankenhaus in Berlin. Since that time many other drugs have been introduced and tried clinically. The three drugs now on the market and in current use are uroselectan, abrodil, and uroselectan B.

Uroselectan.—This is a pyridine compound containing 42 per cent. of organically bound iodine ; 40 grams of the freshly prepared drug is given intravenously in 100 c.cm. of water, corresponding to 17 grams of iodine. The drug is stable, non-toxic, is eliminated through the kidneys to the extent of 85 per cent., and its presence can be determined in the blood some four hours or more after injection, the length of time depending upon the functional capacity of the kidneys.

Abrodil.—This is sodium-iodo-methane-sulphonate. Its iodine content is 52 per cent. ; it is 70 per cent. soluble in water, and is available as a sterile solution of 20 grams in 50 c.cm. of water ; 86 per cent. of the drug is eliminated in the urine.

Uroselectan B.—This also is a derivative of the iodine-pyridine series. In actual practice it has considerable advantages over the above-mentioned. Although only some 63 per cent. is eliminated through the kidneys, this occurs much more rapidly and with much less diuresis than with the other drugs, so that the concentration in the urine is 7.94 per cent. Its iodine content is 51 per cent. A great advantage is the fact that only 20 c.cm. of fluid, containing 15 grams of uroselectan B, need be injected.

It is impossible here to enter into the many technical questions which are concerned with these drugs. With all of them the success of the resulting pictures depends

upon the ultimate concentration of the drug in the urinary tract, and this in turn is mainly dependent upon the functional capacity of the kidneys. Furthermore, the rate of their elimination can be used as a functional renal test. This may be done in one of several ways. First, the amount of the drug retained in the blood at varying time intervals after injection may be estimated and compared with the normal. Secondly, the actual amount of the drugs excreted in the urine over varying time periods may be estimated. Or, finally, the specific gravity of the urine may serve as an approximate guide to the amount of the drug excreted.

MODE OF ACTION

Instrumental urography gives information as to the size, shape, and outline of the cavities of the urinary tract, but it gives no clue to the functional activity of the kidneys or to the motility or dynamics of the rest of the tract. Whilst the older method is essentially anatomical, the new one of excretion urography is strictly physiological. In excretion urography the first essential is that the kidneys shall be able to excrete the drug in sufficient concentration to give a radio-opaque picture. The time of the appearance of the dye will be a measure of the functional activity of the organ. Secondly, the outline of the calyces, the pelvis, and the ureter will be those of the normal working state, and not under any conditions of artificial distension. The work of Cumming has shown the effect of alteration of tonus in the different parts of the calyces and pelvis. From time to time separate contractions of individual calyces or of the pelvis occur, but without, apparently, any orderly sequence. Consequently the outline of an excretion pyelogram depends upon the exact functional state at the moment of the exposure. For this reason, when a series of plates is examined the calycal and pelvic outlines may vary considerably, reflecting the varying phases of filling and emptying of the component parts. In diastole of the calyces and pelvis a picture is obtained which in many respects is very similar to an instrumental pyelogram. On the other hand, a picture taken in systole may show what appear to be filling defects in one or more of the calyces, and the pelvis itself may appear much smaller. In the ureter, too, the resulting picture will be entirely different. Only too often have instrumental ureterograms suggested a comparatively passive tube, whereas in point of fact it is rarely that some point or other of the ureter is not in movement in transporting urine to the bladder. There would appear to be considerable variation in the type and frequency of the ureteric movements, but they are mainly of the nature of a segmental peristalsis, and there is little evidence to show that waves ever pass from end to end of the ureter. The excretion urogram visualizes the dynamics of these processes.

It is only on the basis of a comprehension of the method of action of excretion urography that there can be any adequate interpretation of the resulting pictures. If these are interpreted on the lines of the older instrumental methods, then not only will the urograms often be regarded as technically disappointing, but the deductions made from them will in many cases be erroneous.

TECHNIQUE OF ADMINISTRATION

Certain technical details are of importance if good urograms are to be obtained. The first of these concerns the preparation of the patient. The method adopted in the majority of the cases in the present series was as follows. Thirty-six hours before the injection an ounce of castor oil is given, and the following day the patient is kept on a light diet. Four hours before the injection a breakfast of lean ham, rusks, and a cup of tea with a minimum of milk and sugar is taken, followed by ten ounces of bovril one hour before the injection. On the

day preceding the investigation the total fluid intake should not exceed forty ounces, otherwise marked diuresis may affect the quality of the pictures. Finally, it has to be remembered that in excretion urography flatus is one of the greatest enemies of success, and it is unfortunately still true to say that many pictures are spoilt by the shadows produced by gas in the large bowel. This evil can be minimized by reducing the amount of carbohydrate in the diet on the day beforehand, and, where practicable, by keeping the patient up and about. In fact, much the best pictures are often obtained in out-patients.

The injection of the drug must be carried out with the same scrupulous care as for the dyes used in cholecystography. For this purpose the antecubital veins are by far the most accessible, though in the case of children I have had on occasion to use the external jugular. Cases have been reported in the literature of extensive sloughing after the escape of uroselectan into the tissues. Personally I have only seen one case in which a localized inflammation lasting three days followed the escape of a small amount into the tissues. As regards abrodil and uroselectan B, I have seen no complications of this type, and with the former dye they seem hardly likely, as the drug has been recommended for subcutaneous administration in the case of children. Sichel and Boeckel have followed Köhler in the administration of uroselectan and abrodil by the rectal route, but the pictures are not nearly as good as those obtained by the intravenous method.

POSSIBLE REACTIONS

Certain mild toxic effects may follow the administration of these drugs. In this respect uroselectan is the greatest offender and uroselectan B the most innocuous. In practically every case where uroselectan is used there develops during the injection a sensation of great heat all over the body, together with a marked flushing of the face, symptoms which usually disappear within ten minutes. A few of the patients develop acute thirst after the injection, and a small number nausea and vomiting. The pulse rate is invariably increased by about 40 per minute for about ten minutes. With abrodil these symptoms are very rare, and with uroselectan B they have not been observed at all. In the literature of the subject a few cases of generalized rash of urticarial type, coming on at once after the injection of uroselectan, have been reported and considered as due to the liberation of iodine. In the present series nothing of this sort has been seen, and no serious complications, either local or general, have followed the administration of any of the drugs.

TAKING OF UROGRAMS

Skiagrams of the whole urinary tract are taken at intervals after the injection. With uroselectan these have been at intervals of a quarter, three-quarters, and one and a half hours; with abrodil, a quarter, half, and three-quarters of an hour; and with uroselectan B, ten, twenty, and forty minutes. The exact times obviously are not of extreme importance. What is important is to get urograms in the individual patient during the excretion of the drug. In pathological cases the time will have to be determined after the earlier pictures have been taken, as in those cases where there is any delay in the excretion of the drug the time would obviously have to be extended. The urograms are taken with the patient supine and during full expiration. In the present series of cases no artificial means of compression of the abdomen, bladder, or ureters has been carried out. Many observers, in order to accentuate the pictures obtained by excretion urography, have compressed the lower abdomen by means of inflated bags, or have inserted balloons into the rectum or vagina to compress the ureters. I am sure such methods should in

general be condemned. If they are resorted to in order to intensify excretion pyelograms it must be remembered that the resulting pictures will be comparable to those obtained by instrumental methods. They must not be taken as showing the physiological or pathological condition of function in the urinary tract. Only as such will it be possible to interpret them, and it is questionable whether such methods, if adequately performed, are free from risks.

ANALYSIS OF CASES

During the course of the past two years 144 excretion urograms have been carried out on 138 patients at St. Bartholomew's Hospital, and these cases form the basis of the material of this lecture. Of this number 124 of the investigations were made on patients with diseases of the urinary tract, and the remaining 20 investigations were on patients with diseases of other abdominal viscera, or in cases where disease of extra-urinary origin was thought to have produced secondary changes in the urinary system. As regards the type of injection, uroselectan was used in 58, abrodil in 69, and uroselectan B in 17. The series is admittedly a small one, but I think that from experience with the method some idea of its value can be gained by a study of the cases.

Hydronephrosis.—As a means of investigation in cases of hydronephrosis excretion urography is invaluable, for it not only gives the answer to the question as to the degree of function left in the affected kidney, but in early cases it also demonstrates the site of the obstruction and the secondary changes which have occurred in the organ. It is to be expected, therefore, that a study of urography in these cases will, in the course of time, lead to the formulation of more exact knowledge as to the pathology of hydronephrosis, and consequently lead more often to earlier and more conservative treatment. At the same time, we possess in the method a ready means by which the results of plastic surgery may be checked. A case of bilateral congenital hydronephrosis is shown in Fig. 2 (on Plate), a renal type of hydronephrosis in Fig. 4, and a pelvi-renal type associated with a calculus at the uretero-pelvic junction in Fig. 5.

Horseshoe Kidney.—There were two cases of horseshoe kidney in the series. In one case the excretion pyelogram gave no indication of the condition, owing to associated severe hydronephrosis and consequently marked impairment of renal function. In the second case, shown in Fig. 3, the urograms furnished the diagnosis.

Calculi.—In the presence of functional renal tissue, excretion pyelograms will localize renal and ureteric calculi as accurately as did the older instrumental method. The excretion pyelogram has, in addition, the advantage that, owing to the relatively less radio-opaque nature of the contrast medium as compared with the sodium bromide and sodium iodide usually employed instrumentally, the outlines of calculi in the calyces or pelvis are less likely to be completely obscured by the dye. Furthermore, at the same time information is obtained as to the anatomical and functional changes which have occurred in the diseased kidney as the result of calculi, and also as to the condition of the opposite kidney (Figs. 5 and 6).

Renal Mobility.—In cases of abnormal renal mobility it is possible to demonstrate the position of the organ and at the same time to determine whether or not its descent has had any effect on the capacity of the calyces and pelvis to empty themselves, and also as to whether any dilatation of the latter has occurred. In the seventeen cases of this condition in the series it was found that in none of them was there any abnormality to be made out in the excretion urogram as regards renal function, or the size or shape of the calyces and pelvis on the affected side.

Tubercle.—As a rule the diagnosis of tuberculous disease does not present great difficulty, but in the investigation of such cases excretion urography can often be invaluable. First, in those cases of renal tubercle where a link in the diagnostic chain is missing, it provides a means of getting in early cases an excretion pyelogram of the affected organ, and also a ureterogram and cystogram, while in advanced cases a func-

tionless kidney may be demonstrated; secondly, the excretion urogram provides information as to the functional capacity of the opposite kidney. Of the six cases of proved renal tubercle in the present series, four showed no secretion on the side of the lesion owing either to the extent of renal destruction or on account of ureteric obstruction. In the earliest case hitherto encountered the excretion pyelogram (Fig. 9) shows elongation and deformity of the upper calyx, together with a persistent dilatation of the ureter. Fig. 8 shows the appearance of a kidney, the site of moderately advanced cavernous tubercle, in which a fair amount of functional renal tissue was present; the dye was delayed in appearance, and gradually outlined the cavernous spaces of the organ.

Renal and Bladder Tumours.—In the early diagnosis of renal tumour the one essential is extremely accurate anatomical information, and in view of this, and in the light of the knowledge of the mode of action of excretion urography, it is very doubtful if the latter can ever become an entirely reliable substitute for the instrumental method. On the other hand, in the absence of haematuria the excretion method may prove a useful preliminary in indicating the side of impaired renal function. At times also the aid of excretion urography may be invoked in the investigation of bladder tumours by visualizing the effects which the latter have produced in the upper urinary tract. This is particularly the case where either the extent of the growth or haematuria prevents cystoscopy.

The Ureter.—Perhaps the most important contribution which excretion urography brings in connexion with the ureter relates to the vexed question of stricture. On this subject there has often been great difficulty in the interpretation of instrumental ureterograms. With the excretion method, if, in a series of pictures, pathological changes suggesting angulation, fixation, or absence of motility are detected, then stricture and peri-ureteritis may be diagnosed. In Fig. 7 are shown the changes in a double ureter following an operation some time previously for the removal of stone from the pelvic course. Fig. 10 shows a condition of hydronephrosis and hydroureter in a patient with multiple urinary fistulae to the rectum, perineum, and suprapubic region. In a case in the series (not illustrated) the tremendous lateral displacement of the ureter associated with a huge retroperitoneal ganglion neuroma was well shown. Similarly the relationship of both the kidney and the ureter to external shadows, such as tuberculous glands, was frequently illustrated.

VALUE IN TESTING RENAL EFFICIENCY

A discussion as to how far excretion urography can be regarded as a test of renal efficiency involves a consideration of all the factors which determine the density of the resulting shadows. Now as regards the pyelogram these factors may be summarized as follows: (1) the degree of function of the kidney; (2) the effect of diuresis; (3) the contour of the calyces and pelvis; (4) the degree of obstruction in the urinary tract; (5) the presence of infection; and, in the interpretation of the urograms, the questions as to the time of the appearance of the dye in the renal pelvis and the bladder, and the density of the pictures, are the relevant ones.

Hughes and Péterfi have shown that drugs of the uroselectan group are eliminated principally through the glomeruli. Thus, in any form of nephritis with glomerular lesions there will be delayed excretion. Now in surgical diseases of the urinary tract, especially those where any obstructive lesion is present, it is principally the tubules which are affected. Consequently in such cases glomerular activity may let a considerable amount of dye through, and the effects of the obstructive lesion will be to heighten the resulting shadow. Unless one realizes this, there will be in such cases a marked over-estimation of the real renal capacity. In any excretion urograms where there is a suggestion of any obstructive lesion, it is essential to avoid such a pitfall.

As an example of the above may be quoted the case of a woman who had a uretero-vaginal fistula following pan-hysterectomy some months before. The right kidney never

secreted at all, but on the left side a moderately good renal shadow, with a slightly dilated ureter showing dye in its whole length, was made out. At the time the patient had a blood urea of 50 mg. per cent.; the urea concentration test never reached more than 0.9 per cent. in the urine, and, after indigo-carmin, the dye appeared very faintly at twenty-five minutes on the left side, and not at all on the right. In this case the effects of a moderate degree of ureteric obstruction enhanced the shadow from an extremely defective kidney. The patient died in uraemia about a month later.

As a test of functional capacity the method is important in those cases where for some reason or another it is inadvisable or impossible to obtain the information by instrumental methods. Acute infections of the urinary tract lead to a delayed excretion of the dye, and the improvement in function of the kidneys can be demonstrated with the clearing up of the infection. Similarly in the case of stone in the ureter it is possible to tell whether or no there is complete obstruction (Fig. 12), and, after the removal of the obstruction, to watch the return of renal function.

COMPARISON WITH INSTRUMENTAL METHODS

In their inherent nature instrumental urograms must be regarded as artefacts produced by the cystoscopist for diagnostic purposes, yet capable of giving invaluable information. In this connexion I shall confine my remarks to a comparison of instrumental and excretion pyelograms. In carrying out an instrumental pyelogram the essential condition is that it shall be possible to introduce the opaque media directly into the renal pelvis, and this implies an absence of obstruction to cystoscopy and to the passage of a ureteric catheter. On the other hand, in the case of an excretion pyelogram, the first consideration is that there shall be a sufficient amount of renal tissue to excrete the dye in a concentration necessary to produce a radio-opaque shadow in the renal pelvis. An instrumental pyelogram records the anatomical outline of the calyces and pelvis. The excretion pyelogram records, first, the functional capacity of the kidney; secondly, the anatomical outline of the calyces and pelvis; and, thirdly, the dynamic function of these parts. So that in comparing the two methods we can only compare the instrumental pyelogram with the anatomical results of the excretion pyelogram.

Amongst the present series were thirty-four cases where the excretion pyelogram was checked by an instrumental one, but space does not permit a detailed consideration of these.

INDICATIONS FOR USE

The indications for the employment of excretion urography in diagnosis fall in the main into two groups. In the first group are those cases where, for some reason or other, instrumental methods are impossible or inadvisable. These are as follows: (1) in children, especially males, and in intolerant adults; (2) in urethral obstructions, such as strictures or enlarged prostate, or in the presence of urethritis; (3) in acute infections or tuberculosis of the bladder, or in the presence of haemorrhage, growth, or fistula in that organ; (4) in ureteric lesions of the nature of stricture or fistula, or where malformations are suspected; (5) in suspected calculus anuria. In the second group the indications are relative ones, and under this heading come all those pathological conditions of the urinary tract where it is desired to gain information as to the function of the kidneys on the two sides, or to determine the outline and motility of the ureters.

As regards contraindications to the employment of the method, I find it very difficult to formulate any. The available evidence does not suggest that excretion urography is contraindicated in the presence of even severe renal disease, though with severe associated hepatic

disease it has been regarded as inadvisable. Personally, the only condition in the urinary tract in which I have deliberately refrained from this investigation has been in the presence of very acute infection associated with a high degree of fever.

CONCLUSION

It has been possible in the course of this lecture to touch upon only a few aspects of excretion urography, but I trust I have given you some idea of the scope and possibilities of the method. For the extravagant claims that have already been made by some, that the method is one which will replace the older methods of visualization of the urinary tract, and even in many cases do away with the need for cystoscopic examination, I hold no brief. But experience has shown that excretion urography is safe; that it is, in addition, applicable to a wider range of cases than is the instrumental method; and that it answers questions in terms of anatomy and function. On the other hand, it must be admitted that the interpretation of the urograms is not always easy, and that for the uninitiated the method may provide a large number of pitfalls.

With time, improvements in technique are bound to occur, and it may not be too great a hope that in the near future oral administration of the drugs may become a possibility. Meanwhile the extended use of the method will lead to a fuller knowledge of the physiology and pathology of the urinary tract, especially in regard to the effect of disease of one part upon the function and dynamics of another. In this way the method justly deserves to be incorporated with the older and more tried ones, with which its results must be checked and its advantages and limitations defined. If this is done, then many difficulties of urological diagnosis will be simplified, surgery will gain in outlook and technique from a renewed contact with physiology, and, most important of all, the slender barque of our patient's life is further safeguarded from shipwreck on those dangerous twin rocks of inaccurate diagnosis and ill-advised surgery.

A FATAL CASE OF ACUTE NECROSIS OF THE LIVER ASSOCIATED WITH EPIDEMIC CATARRHAL JAUNDICE

BY

G. M. FINDLAY, M.D., Sc.D.

AND

J. L. DUNLOP, M.B., Ch.B.

ASSISTANT M.O.H. FOR SURREY

(Wellcome Bureau of Scientific Research, London)

(With Special Plate)

During the past few years considerable interest has been aroused, both in this country and abroad, by the occurrence of extensive outbreaks of epidemic catarrhal jaundice. In association with certain of these outbreaks, notably in Sweden, there have been reported a considerable number of deaths from either acute or subacute necrosis of the liver. In this country the disease has been comparatively mild, and relatively few deaths have been recorded. Nevertheless, Morgan and Brown (1927) and Okell (1931) have described fatal cases during epidemics in Northamptonshire and Surrey. The present case, however, is the first in this country in which it has been possible to carry out full pathological and bacteriological examinations.

The patient, a girl aged 3 years and 2 months, was resident in a town in Hampshire immediately adjoining the Surrey boundary. During the past two years, as

R. T. PAYNE: EXCRETION UROGRAPHY

FIG. 1.—Normal. On the right side there is a bifid pelvis, and on both sides there is well-marked cupping of the calyces.

FIG. 2.—Bilateral congenital hydronephrosis. From a youth of 22, whose only symptom was headache.

FIG. 3.—Horseshoe kidney with bilateral hydronephrosis, and calculi in the lower calyx on left side.

FIG. 4.—Hydronephrosis on the right side, and on the left a persistent kink at the pelvi-ureteral junction with early hydronephrosis.

FIG. 5.—Calculus at the uretero-pelvic junction on the right side, showing pelvi-renal type of hydronephrosis.

FIG. 6.—Calculus in the lower calyx of right kidney.

R. T. PAYNE: EXCRETION UROGRAPHY

FIG. 7.—Double ureters on right side, showing marked dilatation as the result of periureteritis following removal of a calculus from the course of the pelvic ureter.

FIG. 8.—Right renal tubercle. The secretion on the affected side was delayed in appearance, and it shows a cavernous type of lesion.

FIG. 9.—Very early right renal tubercle. Showing elongation and deformity of the upper calyx, and persistent dilatation of the ureter.

FIG. 10.—Hydronephrosis and hydroureter. From a patient with urinary fistulas to the rectum, perineum, and suprapubic region. Many years before, left nephrectomy for tubercle, also lithotripsy, etc.

FIG. 11.—From same patient as Fig. 10. Both pictures show the extreme degree of dilatation and kinking of the ureter. This also shows a diverticulum of the bladder.

FIG. 12.—Showing calculus in middle calyx of right kidney, and one in the course of ureter. At no stage any secretion on this side; history suggested that ureteric calculus had been in position about nine weeks.