mittent signs of dry bronchitis, but the prominent symptom was dyspnoea, which had been increasing in severity. This x-ray illustrates the parenchymatous type of growth. Re-examination with lipiodol showed no evidence of bronchial obstruction, the opacity being surrounded by bronchial tubes of normal size and appearance into which the oil had entered without difficulty, though none was visible in the mass itself.

Treatment consisted in the introduction of radon seeds through the thoracotomy opening by one of my surgical colleagues. Considerable improvement resulted, and for about a year the patient lost his symptoms and was able to get about in comparative comfort. Later he developed other symptoms, which suggested the presence of metastatic deposits in the brain, and from the last account I received from his own doctor it was evident that he was going downhill.

Fig. 4 illustrates the typical appearances in a case of "blocked bronchus," demonstrated by injection of lipiodol into the trachea below the cricoid cartilage. The patient

Diagrammatic sketch explanatory of Fig. 4 (showing blocked bronchus).

from whom this skiagram was taken was a middle-aged man, who came under my observation on account of haemoptysis, the cause of which had not been determined. From the history, the presence of bronchial obstruction, and the occurrence of a small pleural effusion, the diagnosis of bronchial carcinoma appeared inevitable. Bronchoscopic examination showed a greyish-white glistening tumour projecting from the mucous membrane of the bronchus into the lumen. Two or three days after this examination the patient had a sudden fit of coughing, and expectorated about an ounce of blood, in which a small portion of solid tissue was discovered. This, on section, proved to be a carcinoma.

REFERENCES

- ¹ Barnard, W. G.: *Journ. Pathol. and Bacteriol.*, 1926, xxix, 241.
- ² Davidson, M.: *Cancer of the Lung and other Intrathoracic Tumours*, Bristol, 1930, p. 75.
- ³ Idem: *Ibid.*, p. 92.
- ⁴ Davidson, M., and Ledlie, R. C. B.: *Brit. Journ. Surg.*, 1928, xvi, 198.
- ⁵ Hanf, D.: *Virch. Arch. f. Path. Anat.*, 1927, cclxiv, 367.
- ⁶ Huguenin, R.: *Cancer primitif du Poumon*, Paris, 1928.
- ⁷ Jackson, C.: *Amer. Journ. Med. Sci.*, 1917, cliii, 371.
- ⁸ Kikuth, W.: *Virch. Arch. f. Path. Anat.*, 1925, cclv, 107.
- ⁹ Levy-Simpson, S.: *Quart. Journ. Med.*, 1928-9, xxii, 413.
- ¹⁰ Orton, H. B.: *Surg. Clin. N. Amer.*, 1926, vi, 1529.

ANTE-NATAL RADIOLOGY*

BY

R. E. ROBERTS, M.D., B.Sc., D.P.H., D.M.R.E.

LECTURER IN RADIOLOGY, UNIVERSITY OF LIVERPOOL; HONORARY RADIOLOGIST, LIVERPOOL ROYAL INFIRMARY, LIVERPOOL MATERNITY HOSPITAL, AND VICTORIA CENTRAL HOSPITAL, WALLASEY

(With Special Plate)

This paper, based on the author's experience in the investigation by means of x rays of six hundred selected cases of pregnancy, is put forward with the object of briefly indicating what information may be expected from this type of examination. The cases are designated "selected" in that the x-ray examination was not of the nature of a routine procedure, but was in each case undertaken at the request of an obstetrician, either to confirm some clinical suspicion of abnormality or to solve some problem which could not be clearly elucidated by the ordinary methods of examination. Among the reasons

* Read in the Section of Obstetrics and Gynaecology at the Centenary Meeting of the British Medical Association, London, 1932.

which resulted in the help of the radiologist being solicited the following may be regarded as the more important:

1. To make a differential diagnosis between pregnancy and tumour in the case of an obscure abdominal swelling.
2. To ascertain the age of the foetus when this was in doubt clinically.
3. To determine the position and presentation of the foetus in doubtful cases.
4. To assess, by radiological pelvimetry, the exact measurements of the pelvic brim or outlet in a case of suspected contracted pelvis.
5. To investigate any suspected disproportion between the foetal head and the maternal pelvis.
6. To decide whether more than one foetus was present.
7. To determine the presence or otherwise of extra-uterine pregnancy.
8. To investigate the cause of hydramnios, with special reference to the presence and nature of foetal abnormalities.
9. To help in the diagnosis of suspected intrauterine death.

Differential Diagnosis of Pregnancy from Hydatidiform Mole and Pelvic Tumours

Since the advent of the Zondek-Aschheim test the radiological diagnosis of early pregnancy is of less importance than previously. The positive diagnosis by direct radiography is sometimes possible at the twelfth or thirteenth week. As a rule, however, it is not safe to exclude pregnancy by this method till after the sixteenth week. Even then the absence of foetal shadows should not be taken as conclusive evidence on one film only. At least three radiographs should be taken to exclude the possibility of non-visualization of the foetal parts through movement.

The Age of the Foetus

Where, owing to disparity between the period of amenorrhoea and the size of the uterus, an element of doubt exists as to the period of gestation, useful information can usually be given by x rays. In addition to the radiographic size of the foetus and the stage of ossification of its bones, one must take into consideration its position relative to the maternal parts, the presence of hydramnios or multiple pregnancy, or any other factor which may bear on its relative distance from the film and the consequent enlargement of its shadow on the radiograph. In order to build up a standard of comparison of sizes of foetuses at different ages it is important that a standard distance between the x-ray tube and the film should be maintained by the radiologist. A useful working distance is 30 inches.

Position and Presentation

For a complete orientation of the foetus three radiographs are, as a rule, necessary—namely, postero-anterior, antero-posterior, and lateral. The radiographs will give not only a true picture of the presentation, but also clear information as to the relation of the foetal parts to each other and to the maternal pelvis and spine. This is of special importance with vertex presentations in assessing the degree of flexion or otherwise of the head, or with breech cases in demonstrating the degree of flexion or extension of the legs and the position of the hands in relation to the head (Fig. 1). Out of 100 breech cases occurring in this series seventy-six showed extension of the legs.

Maternal Pelvic Deformities

By placing the patient in such a position that the pelvic brim is parallel to the x-ray film and the tube directly over the centre of the brim, it is possible to obtain a symmetrically enlarged picture of the superior strait. By measuring the distance of the tube from the upper border of the symphysis pubis and from the film it is quite a simple matter to estimate the exact dimensions of the pelvic brim in all diameters.^{1 2 3} With the

patient in a "squatting" position on the film it is also easy to estimate accurately the transverse diameter of the pelvic outlet and the pubic angle.⁴

Disproportion

While exact pelvimetry is of importance, the relative size of the foetal head to the maternal pelvis is naturally of greater importance (Fig. 2). Information on this point is as a rule readily available if radiographs are taken in the three standard positions mentioned. By means of three such radiographs one has a good idea as to what allowance to make for relative enlargement of the skull and pelvis due to their different distances from the films.

Multiple Pregnancy

The x -ray examination will settle the question of multiple pregnancy beyond doubt. Moreover, the radiograph shows clearly the relative position of the foetuses to each other and to the maternal parts. In this series twins were demonstrated by x rays in eighty-four cases and triplets once (Fig. 3). In many cases twins were demonstrated radiologically where they were, at the time of examination, unsuspected clinically; while in many cases of hydramnios only one foetus was present where a tentative suggestion of multiple pregnancy had been made clinically.

Extrauterine Pregnancy

From the size and abnormal relation of the foetal parts to the uterine shadow it is at times possible to make a definite radiological diagnosis of extrauterine pregnancy. But unless the uterine shadow can be clearly seen in the radiograph an abnormal position of the foetus alone is insufficient for a positive diagnosis. In such cases most careful collaboration between the radiologist and the obstetrician is of paramount importance. Possibly the introduction of a rubber tube, or of lipiodol, into the uterine cavity might give the required information as to the relation of foetus to the uterus, but the justifiability of such a procedure is one which the obstetrician alone can decide. The tube method was resorted to in one of the cases in this series. Three cases of extrauterine pregnancy occurred; two were diagnosed radiologically, the third ought to have been.

Foetal Abnormalities

Foetal abnormalities are often suspected from the presence of hydramnios, but usually not definitely diagnosed clinically. They are very readily demonstrated by means of x rays. In this series the following foetal abnormalities were found at the ante-natal x -ray examination and confirmed at birth.

	Cases
Anencephalus (Fig. 4)	17
Hydrocephalus (Fig. 5)	6
Iniencephalus (Fig. 6)	3
Meningocele (skull abnormality diagnosed) ...	1
Rudimentary upper limbs	1

In the radiological diagnosis of hydrocephalus it is important to take into consideration the degree of ossification of the cranial bones in addition to the size of the foetal head, otherwise a post-mature foetal head might be mistaken for a mild degree of hydrocephalus. In iniencephaly the radiographic demonstration of the deformity and retroflexion of the cervical spine gives the clue to the diagnosis. Radiographs in more than one plane are necessary to avoid mistaking a normal foetus with a hyperextended head for an iniencephalic foetus.

Intrauterine Death

The only reliable radiological sign of intrauterine death is that first described by Spalding. Owing to shrinkage of the brain there is, within five to seven days, a falling in of the vault producing an overriding or crinkling of the cranial bones (Figs. 7 and 8). This must not of

course be confused with the overriding or moulding which normally takes place during labour. In this series there were sixteen cases of intrauterine death. In nine of these Spalding's sign was positive at the first x -ray examination. In cases of doubt the examination should be repeated after an interval of seven to fourteen days, when, in the case of intrauterine death there usually will be seen to be absence of further growth, increased overlapping of cranial bones, and often an increased curvature of the foetal spine.

RADIOGRAPHIC TECHNIQUE ADOPTED

Apparatus: Snook transformer, hot cathode tube, Potter-Bucky couch with compressor band.

Kilovoltage: 80 to 100 kV peak.

Exposure: 70 to 350 milliamperes-seconds.

Film target distance: 30 inches.

Position of patient: prone; supine; lateral; occasionally oblique. All in full expiration, maintained when necessary by closure of lips and pinching of nostrils by the patient.

CONCLUSIONS

Ante-natal radiology gives valuable information in the following respects:

1. Diagnosis of pregnancy after the sixteenth week, differential diagnosis of pregnancy from hydatidiform mole and pelvic tumours.
2. Estimation of age of foetus.
3. Position and presentation of foetus.
4. Estimation of pelvic measurements.
5. Investigation of possible disproportion between foetus and maternal pelvis.
6. Investigation of cause of hydramnios, with special reference to multiple pregnancy and foetal abnormalities.
7. Diagnosis of extrauterine pregnancy.
8. Diagnosis of intrauterine death.

Neither in the author's experience nor that of others is there any evidence that diagnostic ante-natal exposure to x rays is in any way harmful to the foetus or mother.⁵

The author wishes to express his grateful appreciation of the facilities afforded and help given by the obstetrical staff of the Liverpool Maternity Hospital and of the Liverpool Royal Infirmary in the radiological investigation of their cases.

REFERENCES

- ¹ Thoms: *Amer. Journ. Obstet. and Gynec.*, October, 1926.
- ² Roberts: *Brit. Journ. Radiol.*, January, 1927.
- ³ Rowden: *Ibid.*, September, 1931.
- ⁴ Chassard and Lapiné: *Journ. de Radiol. et de l'Electrol.*, 1923, p. 113.
- ⁵ Goldstein and Murphy: *Amer. Journ. Roentgenol.*, 1929.

A CASE OF ADDISON'S DISEASE TREATED WITH CORTICAL SUPRARENAL EXTRACT

BY

J. G. McCRIE, M.B., F.R.C.P.Ed.

LECTURER IN MEDICINE, UNIVERSITY OF EDINBURGH; TUTOR IN
CLINICAL MEDICINE, ROYAL INFIRMARY, EDINBURGH

I. M. MEARS, M.B., Ch.B.

ASSISTANT MEDICAL SUPERINTENDENT, THE ASTLEY AINSLIE
INSTITUTION, EDINBURGH

AND

W. G. MILLAR, M.B., Ch.B.

LECTURER IN PATHOLOGY, UNIVERSITY OF EDINBURGH; PATHOLOGIST,
ROYAL INFIRMARY, EDINBURGH

(With Special Plate)

Examples of Addison's disease in which an opportunity for full pathological examination is presented are not commonly encountered, Davidson² stating that, up to the present, there have been published full reports of only eighty-two cases. The fact that the practical utility of treatment with cortical suprarenal extract is at present under consideration is an additional justification for the publication of a single case report.

R. E. ROBERTS: ANTE-NATAL RADIOLOGY

FIG. 1.—Breech with extended legs—hands in front of brow.

FIG. 2.—Disproportion between full-term foetus and generally contracted pelvis.

FIG. 3.—Triplets.

FIG. 4.—Anencephalic foetus—post-mature. (Note well-ossified femoral and tibial epiphysis.)

R. E. ROBERTS: ANTE-NATAL RADIOLOGY

FIG. 5.—Hydrocephalic foetus.

FIG. 6.—Iniencephalic foetus. (Note lesion of cervical spine causing hyperextension of head and brow presentation.)

FIGS. 7 and 8.—Intrauterine death (Spalding's sign positive).