- Bernhardt, H., and Levy Simpson, S.: To be published.
 Bulloch, W., and Sequeira, J.: *Trans. Path. Soc. London*, 1905, lxvi, 189.
 Cannon, W. B.: *Lancet*, 1930, i, 1109.
 Corey, E. L. and Britton, S. W.: *Amer. Journ. Physiol.*, 1931, xciv, 33.
 Idem: *Ibid.*, 1931, xcix, 33.
 Elliot, T. R.: *Journ. Physiol.*, 1914, xlix, 38.
 Fordyce, A. D., and Evans, W. H.: *Quart. Journ. Med.*, 1929, xxii, 557.
 Glynn, E.: *Journ. Obstet. and Gynaec.*, 1921, xxviii, 23.
 Greene, C. H., Rowntree, L. G., Swingle, W. W., and Piffner, J. J.: *Amer. Journ. Med. Sci.*, 1932, clxxxiii, 1.
 Hartmann, F. A., MacArthur, C. G., and Hartmann, W. E.: *Proc. Soc. Exper. Biol. and Med.*, 1927, xxv, 69.
 Hartmann, F. A., Thom, G. W., Lockie, L. M., Greene, C. W., and Bowen, B. D.: *Journ. Amer. Med. Assoc.*, 1932, xcvi, 788.
 Hutchison, R.: *Quart. Journ. Med.*, 1907-8, i, 33.
 Labbé, M., Tinel, J., and Doumer: *Bull. et Mém. Soc.*, 1922, xli, 982.
 Levy Simpson, S.: *Proc. Roy. Soc. Med.*, 1930-1, xxiv, 497.
 Idem: *Journ. Physiol.*, 1931, lxxii, 4.
 Idem: *Quart. Journ. Med.*, 1932, i, 99.
 Marshall, E. K., and Davis, D. M.: *Journ. Pharm. and Exper. Therap.*, 1916, viii, 525.
 Meirowsky, E.: *Munch. med. Woch.*, 1911, lviii, 1005.
 Neusser: Quoted by Orth, J.
 Oppenheimer, B. S., and Fishberg, A. M.: *Arch. Int. Med.*, 1924, xxxiv, 631.
 Orth, J.: *Sitzung. d. k. Preussischen Akad. d. Wissensch.*, 1914, ii, 34.
 Rogoff, J. M., and Stewart, G. N.: *Science*, 1927, lxvi, 327.
 Rolleston, H. D.: *West London Med. Journ.*, 1925, xxx, 105.
 Rowlands, R. A.: Personal communication.
 Rowlands, R. A., and Levy Simpson, S.: *Proc. Roy. Soc. Med. (Section of Medicine)*, 1929, xxiii, 16.
 Rowntree, L. G., Greene, C. H., Ball, R. G., Swingle, W. W., and Piffner, J. J.: *Trans. Assoc. Amer. Phys.*, 1931, xli, 123.
 Rowntree, L. G., Greene, C. H., Swingle, W. W., and Piffner, J. J.: *Ibid.*, 1931, xcvi, 231.
 Rowntree, L. G., and Snell, A. M.: Mayo Clinic Monographs, 1931.
 Stewart, G. N.: *Arch. Int. Med.*, 1929, xliii, 733.
 Swingle, W. W., and Piffner, J. J.: *Science*, New York, 1930, N.S., lxxi, 321.
 Wadi, W.: *Klin. Woch.*, 1927, vii, 2107.
 Wiesel: Quoted by Oppenheimer, B. S., and Fishberg, A. M.

CARE OF THE LUNGS IN ANAESTHESIA*

BY

H. W. FEATHERSTONE, M.D.

LECTURER ON ANAESTHETICS, UNIVERSITY OF BIRMINGHAM

This subject may be described with truth as a vast one. In the course of a surgical operation under general anaesthesia the widespread effects of the drugs employed, the altered state of the entire physiology of the body, and the immediate and remote consequences of the actual operation are problems which must receive the constant attention of the surgeon and of the anaesthetist during their work. Since the lungs are affected in greater or lesser degree by these bodily disturbances, the anaesthetist must regard his responsibilities as being wide in their scope and profound in their detail.

Formerly, post-operative lung disorders were summed up by many as "ether bronchitis" or "ether pneumonia"; more recently research has been devoted to the subject of "post-operative massive collapse of the lungs," and during the last year or two attention has been drawn to the therapeutic value, and the prophylactic wisdom, of insufflations of carbon dioxide gas when mixed with air or with oxygen.

In this paper the difficulties which an anaesthetist may look for are summarized, but it is hoped that subsequent papers from physicians and physiologists may open our eyes still further to certain risks and wise precautions concerning which these workers are more qualified to show the way. Perhaps I may group the problems which have to be considered under the following general headings: (a) the anaesthetic agent; (b) the method of administration; (c) the operation itself; (d) the patient's general condition; (e) the patient's respiratory system; (f) the process of infection.

ANAESTHETIC AGENT

Of the anaesthetic agents which are available for inhalation, nitrous oxide, ethylene, ether, and chloroform are the drugs most commonly employed. They provide examples of different chemical and physical states—namely, gaseous oxide of nitrogen, gaseous hydrocarbon, liquid oxide of a hydrocarbon, and liquid halogen compound of a hydrocarbon. Ether vapour has an irritant effect on the mucous membrane of the respiratory passages, but sound technique renders this ill effect negligible. Impurities such as ketones and peroxides are harmful,¹ and should be avoided by employing only the purest ether. It is essential that the ether should be well vaporized, for a cold fog of ether droplets is particularly irritating. Sir Francis Shipway was the pioneer who drew attention to the importance of administering warmed ether gas, and his apparatus is one of the most important items in an anaesthetist's armamentarium. The proportion of ether vapour in the inhaled mixture of gases should be as low as is practicable, for the higher concentrations are very pungent; indeed, experiments on dogs² have shown that massive doses of ether produce petechial haemorrhages and oedema of the lungs. In lesser concentrations, narcotic drugs paralyse the vasomotor control of the blood vessels of the lungs, and physiological friends have suggested to me that this factor may be of importance in determining a considerable degree of damage to the lung tissue.

Salivation and profuse secretion from the upper air passages and the lungs are controlled by pre-operative administration of atropine, but the local injury to the epithelial cells, and the paralysis of the ciliated epithelium, should be prevented as far as possible by using only pure ether, warmed, thoroughly vaporized, and well diluted with air or oxygen. When these precautions have been taken, it is a common experience to find that the lungs of patients who have suffered from subacute bronchitis for a week or more prior to operation clear up rapidly after the administration of ether.

It may be of interest to remark that, in addition to the local effects of ether on the lungs, Graham³ showed that while ether checked phagocytosis in the body, olive oil administered by the rectum, or lecithin injected subcutaneously, restored the opsonic index to normal. Cutaneous vaso-dilatation and sweating should be dealt with by covering the body adequately with blankets and by the use of atropine, lest these indirect effects of ether predispose to lung disease. In long operations it has been realized that the dangers attaching to the prolonged administration of ether should be avoided by the employment of morphine or basal narcosis, together with nitrous oxide or ethylene with oxygen, and as little ether as possible.

Chloroform, in the weak dilution which is necessary for safety in inhalation anaesthesia, is not pungent to the nasal and oral mucous membrane, but experimentally it has been shown that strong chloroform vapour is destructive in its action on animal cells.⁴ Although the more serious risks of chloroform anaesthesia are due to the damage to the cardiac nerves, the heart muscle, and the liver, the frequency of post-operative pneumonia following chloroform anaesthesia is noteworthy, and perhaps the vasomotor disturbances in the lung tissues are more considerable than has been generally recognized.

Nitrous oxide and ethylene are gases which are comparatively inert in their local effects on the respiratory epithelium. The absence of poisonous effects is emphasized by the excellent results which have been obtained by Magill and Rowbotham⁵ in anaesthesia for the prolonged multiple operations which are entailed in plastic surgery of the face. Furthermore, they are the anaesthetic agents of choice in chest surgery.

* Read in opening a discussion in the Section of Anaesthetics at the Centenary Meeting of the British Medical Association, London, 1932.

Spinal analgesia when first introduced was hailed as the certain method for the protection of patients against post-operative lung sequelae, but experience has shown that lung disease is by no means avoided by the use of this technique. The local effect of paralysing some of the intercostal muscles perhaps conduces to immobility of the lower lobes and pulmonary congestion. Marked lowering of blood pressure may reduce the body's resistance to infection, and on several occasions in my experience,⁶ within a few hours of the intrathecal injection of a local anaesthetic a slight bronchitis has developed into definite consolidation of the lungs. It has been shown by Meakins and others⁷ that in spinal paraplegia, involving the lower intercostal muscles, carbon dioxide accumulates in the blood, and at the same time oxygenation is deficient. Furthermore, lesions of the dorsal region of the spinal cord are usually complicated by purulent bronchitis or bronchopneumonia. Therefore splanchnic analgesia, by preventing reflex inhibition of the muscles of respiration, should be a valuable safeguard, and many workers are satisfied that that is true.

METHOD OF ADMINISTRATION

In the selection of the method or technique of administration of an anaesthetic many considerations must be reviewed, but attention may here be drawn to certain precautions. The closed method of Clover, Hewitt, and others has had a wide vogue. Rebreathing produces excess of carbon dioxide in the bag, and the respirations become deeper and more rapid. From the apparatus the patient may then inhale considerable volumes of cold and concentrated ether vapour. Breaths of air and careful manipulation of the ether control will avoid this danger. Open or "per-halation" ether nowadays is a routine method. Perhaps it may be stated with fairness that beginners soak the gauze too freely with ether, which tends to a frozen mask and a cold ether fog. More gauze and a steady drip from the ether bottle uses less ether and gives a less pungent odour. Intratracheal methods require gentle introduction of the catheter between fully abducted vocal cords (Magill's rubber tubes provide a wider margin of safety than the gum-elastic variety). During the operation the concentration of ether should be low. The cough reflex should return before the catheter is withdrawn, lest collapse of the trachea or of a bronchus should supervene.⁶

EFFECT OF THE OPERATION

So far as the effect of the operation on the lungs is concerned there are several points to be considered. The posture of the patient, whether supine, prone, sitting, or the Trendelenburg position, is not of much importance. Perhaps the sitting posture increases the risk of aspiration from the upper air passages, while the Trendelenburg position reduces this risk. Compression of the ribs, and limitation of chest movement in the prone position, may conduce to stagnation of the respiration and of the circulation in the lungs, but two long parallel sandbags, supporting the sides of the chest, allow the sternum and the anterior portion of the chest to move more freely. The site of operation may exert a powerful influence on the condition of the lungs. While operations on the pelvic organs or on the lower abdomen are rarely complicated by lung disorders, surgery of the upper abdomen is often marred by post-operative lung collapse or consolidation.⁶ The disturbance to the muscles and nerves which is occasioned by upper abdominal operations will be discussed at a later stage of this paper, but here it may be pointed out that in diseases of the stomach, starvation or indigestion or delayed absorption of food-stuffs may predispose to lung disease. The pandemic of influenza in the later stages of the war and the high

morbidity rate for post-operative pneumonia just after the war⁶ serve to remind us of the effects of malnutrition. Purgation and pre-operative starvation should be avoided, and rectal feeds of glucose, water, and other substances are important.

Operative shock is a potent cause of lung disease. Analysis of hospital post-mortem records⁶ shows that close approach to death, whether due to accidental injuries, to surgical trauma, or to disease, involves the onset of pneumonia in 30 to 50 per cent. of fatal cases. Care should be taken to avoid needless damage to the tissues, loss of blood, and chilling of the patient's body, before, during, and after the operation. Loss of heat from the skin is increased by exposing the skin to draughts, and by general cutaneous vaso-dilatation. Where dry cold air is inhaled, the consequent warming of the air and evaporation in the alveoli to saturate it with water vapour abstracts much heat from the body. Especial care should be taken with elderly persons, and perhaps men are more prone to lung disorders than women. It is possible that the abdominal or diaphragmatic type of respiration in men predisposes to lung trouble. Women breathe without straining the abdominal wound, and perhaps it is fair to say that coughing disturbs abdominal wounds less in women than in men.

GENERAL CONDITION AND RESPIRATORY SYSTEM

Disturbance of the metabolism and of the organs of defence occurs during anaesthesia. If the patient is already suffering from poor general nutrition, from cachexia due to disease such as cancer, from drug addiction such as chronic alcoholism, or from circulatory failure, special care must be taken to spare the lungs from direct irritation and the body from the effects of the more toxic anaesthetics. There is strong evidence that patients who have suffered from pneumonia during the year or two prior to a major operation are especially liable to post-operative pneumonia. Care should be taken to ensure that these patients may be in the best possible state of health when an operation is to be undertaken under general anaesthesia.

When considering the effect of anaesthesia upon respiratory function, the importance of avoiding anoxaemia must be borne in mind. Additional oxygen must readily be given to those whose respiratory exchange is embarrassed by any of the following conditions: deformities of the chest; paralysis of the muscles of respiration; laryngeal obstruction; loss of function in part of the lungs due to injury; emphysema, or previous disease (such as permanent collapse from pneumonia); frothy secretion in ether anaesthesia without pre-medication with atropine; shallow breathing due to enfeeblement of the respiratory centre as the result of shock or anaesthetic overdose. In this connexion the work of Haldane,⁸ Meakins, and many others is gratefully acknowledged by anaesthetists. In the surgery of toxic goitre extra oxygen is a most valuable safeguard, both during the operation and during the recovery stage. When conscious these patients recognize the relief given by oxygen, and the risk of acute hyperthyroidism is reduced.

The clinical entity first described by Pasteur⁹ and called by him massive collapse of the lungs has received wide attention and study. Knowledge of the mechanism of massive collapse is of the first importance to anaesthetists in order that all possible precautions may be taken to avoid the condition. Many workers believe that occlusion of bronchioles or of a bronchus by inflammation, spasm, or secretion¹⁰ is followed by absorption of the imprisoned air and collapse of the affected area. Woodburn Morison,¹¹ from his extensive radiological experience, believes that "the primary cause of active collapse is a reflex disturbance of the sympathetic and vagus innerva-

tion of the lungs." Others attribute the condition to inactivity of the muscles of respiration resulting from (1) paralysis (as in post-diphtheritic cases); (2) inhibition (the result of trauma or of post-operative disturbance of the action of the abdominal muscles, of the diaphragm, or of the intercostal muscles). Sir Charlton Briscoe¹² believes that the crural portion of the diaphragm is the only part of that muscle which contracts in normal respiration in the supine position, but after an operation, or in cases of debility or toxæmia, the contraction of the crural portion diminishes, and deflation of the lower lobe follows. Pneumonia or pleurisy may start this process and increase it.

Some years ago I brought forward arguments⁶ which suggested that post-operative massive collapse is the result of several factors acting in conjunction. Thus, rigidity of the abdominal wall from pain and trauma leads to reflex inhibition of the diaphragm; congestion and oedema occur in the quiescent lower lobes; the consequent swelling and spasm of the bronchioles is increased by irritation from the anaesthetic agent, and infective material is introduced by aspiration and possibly by lymphatic infection; closure of the bronchioles and absorption of air from the bronchi then ensues.

In January of this year Band and Simson Hall of Edinburgh published some important work on the subject.¹³ After describing four clinical cases of post-operative massive collapse, they give experimental evidence which indicates that certain factors, acting in combination, may lead to collapse. These are, first, a viscid intrabronchial secretion; secondly, abolition of the cough reflex; thirdly, limitation of the movement of respiration. In dogs, gum acacia solution of 75 per cent. viscosity was instilled into certain bronchi, while the cough reflex was abolished by means of morphine, and the movements of the ribs were limited by strapping. In each experiment massive collapse of the lung was demonstrated by radiography. Valuable as this work undoubtedly is, it should be noted that the conditions of the experiment were more harsh than the circumstances of an upper abdominal operation, and the application of a strong solution of gum acacia to the respiratory epithelium surely was directly irritant, and in these experiments perhaps a certain degree of oedema and even of inflammation should be added to the three causal factors which the authors suggest.

In the treatment of massive collapse the therapeutic value of inhalations of carbon dioxide is emphasized by Band and Simson Hall. Carbon dioxide, by its action in deepening respiration and increasing the frequency of breathing, is a most valuable friend to the anaesthetist. Induction is expedited, the deepening of anaesthesia in morphinized patients becomes practicable, rapid recovery of the cough reflex may be achieved, and post-operative collapse is guarded against by stimulating strong expansion of the lungs at the end of operations on the upper abdomen. Many prefer to use a short administration of concentrated carbon dioxide, which is arrested when deep breathing has commenced. Others consider that the administration of 7 per cent. carbon dioxide with oxygen for several minutes is safer, and equally efficacious in promoting maximum expansion of the lungs.

LUNG INFECTION

Infection of the lungs, whether they be prone to massive collapse or not, is to some extent the province of the anaesthetist. The aspiration of infected material from the mouth, nose, and pharynx is to be reduced by pre-operative cleansing of the mouth and nose, by dental treatment, by a free airway, and by encouraging the early return of the cough reflex. The existence of small epidemics of coughs, colds, bronchitis, and influenza

renders the subjects of upper abdominal operations specially prone to post-operative lung inflammation. Whenever possible it is reasonable to suggest that the more serious operations in a ward where colds are rife should be postponed for a time. Infection of the lung from other parts of the body by direct spread, by lymph-borne and by blood-borne infection, is usually unavoidable. In the post-mortem room, pneumonia secondary to peritonitis, middle-ear suppuration, septicaemia, and the like are shown to be pathological sequences of events, which are often overwhelmingly powerful.

Post-operative pulmonary embolism remains serious and unexplained. Sepsis, vascular stasis, excess of blood platelets, or dearth of hepatic antithrombin are causes beyond the ken of the anaesthetist, but the suggestion has been made that the temporary concentration of ether in the blood stream actually predisposes to intravascular clotting. Embolism after brief operations under gas-oxygen for the ligation of haemorrhoids, for herniotomy, and for prostatectomy, is by no means unknown, so that the administration of ether can hardly be counted of primary responsibility in these catastrophes.

I have dealt in a somewhat sketchy fashion with a number of the points which must engage the attention of the careful anaesthetist in his endeavour to protect the lungs of his patient against harm, but I trust enough has been said to prove that the care of the lungs in anaesthesia involves wide knowledge and meticulous care.

REFERENCES

- ¹ Storm van Leeuwen: *Proc. Roy. Soc. Med.*, March, 1924.
- ² Poppert: *Deut. Zeit. f. Chir.*, Band 67.
- ³ Graham: *Journ. Amer. Med. Assoc.*, 1911.
- ⁴ Flemming: *British Medical Journal*, 1922, ii, 22.
- ⁵ Rowbotham and Magill: *Proc. Roy. Soc. Med.*, February, 1921 (and elsewhere).
- ⁶ Featherstone: *Brit. Journ. of Surg.*, January, 1925.
- ⁷ Meakins: *British Medical Journal*, 1924, i (and other papers).
- ⁸ Haldane: *Proc. Roy. Soc. Med.*, January, 1932.
- ⁹ Pasteur: *Lancet*, November 7th, 1908, and May 9th, 1914; *Brit. Journ. of Surg.*, i, 1914.
- ¹⁰ Elliott and Dingley: *Lancet*, 1914, i.
- ¹¹ Woodburn Morison: *British Medical Journal*, 1930, ii, 237.
- ¹² Briscoe: *Quart. Journ. of Med.*, 1919, 332.
- ¹³ Band and Simson Hall: *Brit. Journ. of Surg.*, January, 1932, 387.

Memoranda

MEDICAL, SURGICAL, OBSTETRICAL

A CASE OF ICHTHYOSIS HYSTRIX

(With Special Plate)

Linear naevus, itself a rare disease, is usually met with in the form of a band of varying length following the line of a nerve in a limb or one side of the body. The pathologically identical giant form is still more rare, and the photographs reproduced in the Plate illustrate an example of ichthyosis hystrix, the most extensive, I believe, on record.

The growth in this case is distributed over the whole trunk, neck, axillae, and groins. It is most dense in the flexural sites, where it presents sodden masses half an inch thick. The growths on the front and back of the neck are like small yellow warts. Similar patches (not shown in the photographs) lie about the sides of the forehead and angles of the mouth, but elsewhere the arrangement in lines is peculiarly distinct. When first seen the linear growth was of a dark-brown colour, actually black in parts, but following the application of salicylic acid ointment is now a dark fawn. The patient is 16 years of age, and the mother assures me that the eruption was so pale that she paid no attention to it until it darkened rapidly three years ago. It may interest those who believe in maternal impressions to learn that