The main argument in favour of immediate operation in all cases is the difficulty of diagnosis. It may not be possible to be certain that the peritonitis is not secondary to appendicitis. Many cases have been reported where an erroneous diagnosis of pneumococcal peritonitis has been made and a diseased appendix found at operation. In the two cases I have already mentioned which were admitted to hospital within an hour, the withholding of operation in the first case would have meant certain death from appendicitis and peritonitis; in the second case, operation made no difference to the inevitable fatal termination.

My rules of conduct would be these:

If the diagnosis is certain and if the infection as shown by the physical signs is diffuse, then operation is contraindicated.

If the diagnosis is certain and if there is reason to believe that the infection is most intense in one area—for example, the pelvis in girls—then a simple drainage operation is performed. If the diagnosis is uncertain, especially in males, then a quick laparotomy under gas and oxygen, to determine the condition of the appendix and to institute pelvic drainage, is the safer treatment.

The intra-abdominal manipulations should be reduced to a minimum. It is usually possible to realize the nature of the infection at once on opening the peritoneum. The straw-coloured odourless fluid, the bright red roughened peritoneum, the fibrinous patches, some filmy adhesions between the slightly distended coils, form a picture that cannot be mistaken.

In the absence of much exudate it might be justifiable to close the peritoneum, although I have not had the opportunity of testing this procedure.

Whether operation is undertaken or not, immediate steps must be adopted to combat the intense septicaemia. The organism should be rapidly typed in the manner suggested by Armstrong, and the appropriate serum administered. I will leave to others in this discussion, more qualified to speak, the question of the appropriate amount and frequency. Blood transfusion is considered of great value, but I must confess, in my own cases, I have found it a little disappointing. Perhaps one expected too much. Fluids must be administered in large quantities, and glucose solution intravenously is of great value. Mercuriochrome and mercury perchloride intravenously have seemed to do good in some cases.

STREPTOCOCCAL PERITONITIS

I must give a separate word to that very fatal condition streptococcal peritonitis. I have alluded to it more than once in discussing pneumococcal peritonitis, and it has many points of resemblance. The clinical features and pathological appearance are very similar, and differ only in degree: 71 per cent. of the cases were in females. Its origin corresponds very closely. I believe the vaginal route of infection is very common, and infection of the peritoneum by the blood stream accounts for all the other cases. Its septicaemic complications are very severe. In my experience, recovery with or without subacute encysted intra-peritoneal collections never occurs. Operation has no effect. In fifteen years there have been twenty-two cases at Great Ormond Street: twelve were operated upon, ten were not, but all terminated fatally.

One can only hope that, with the passage of time, the streptococcus will become less virulent, or the human immunity greater, and that the disease will become more benign. Hopefully one may recall the vast reduction in the number of cases of fulminating osteomyelitis due to the staphylococcus which have occurred in our own memory. In the severe streptococcal peritonitis, operation is not demanded, save for the exclusion of an appendix peritonitis. Serum, blood transfusion, and the administration of large quantities of fluids will be the better line of treatment.

THE EFFECTS OF ANAESTHETICS ON THE LIVER *

BY

WESLEY BOURNE, M.D., C.M., M.Sc., F.R.C.P.(C.)

MONTREAL, CANADA

Attention was first directed to "chronic chloroform poisoning"¹ shortly after the discovery of anaesthetics. Since that time interest in the action of this drug on the liver has been more or less constant. Stiles and M'Donald² have reviewed the earlier literature and given their findings, and Whipple and Sperry³ have contributed most invaluable data; yet I shall not recount extensively the inceptive investigations, but rather the more recent ones concerning the effects of several chemical substances which are used in anaesthesia.

Latterly, considerable information has emanated from the Mayo Foundation on the physiology of the liver (^{4 5 6 7 et al}), and the value of functional tests of this organ.^{4 8} What with partial and total removal of the liver, heart-lung-liver and heart-lung-leg perfusions, denervation, studies on detoxication, and so forth, our knowledge of the several known vital functions of this great organ has been made much more clear and a rational evaluation given to the many tests, of which it may be said with authority that Rosenthal's⁹ is the most dependable in a relative and comparative sense.

It has been shown¹⁰ that brief periods of chloroform anaesthesia are sufficient to produce immediate and delayed toxic effects on the liver in the dog; that half an hour of chloroform causes injury which requires eight days for functional recovery, while it takes as long as six weeks for return to normal after two hours of chloroform anaesthesia; that disturbances of function could be demonstrated with the bromsulphalein test long after pigment metabolism had returned to normal; that ether produces a definite but transitory impairment of function, recovery being complete usually within twenty-four hours; and that nitrous oxide and ethylene do not produce any change in this dye test for hepatic function or any disturbance of pigment metabolism, but when there is a reduction in the oxygen supply during the administration of these gases, liver function is so deranged that it takes about six days to come back to normal. At this juncture a word may be said on the topic of cyanosis. Nothing but harm comes to aerobiotic cells through deprivation of oxygen; take, for example, the finding of Koehler, Brunquist, and Loevenhart,¹¹ that anoxaemia produces a more marked acidosis than has been seen in any other condition correlative with depression to all functional activity. This is the point at which one may put it down most axiomatically that any measure of functional perturbation in any one part of an organism is in direct proportion to the damage going on in all of the rest of that being, and, as the liver is such an important structure, it may be looked upon as a good index.

Large doses of morphine cause considerable, but transitory, impairment of function of the liver in dogs,¹⁰ but in the human being,^{10 12} with whom the therapeutic dose is from thirty to fifty times smaller in size, it is believed that no such damage occurs. Amytal, a representative of the malonyl-urea derivatives, has very little action upon liver function, comparing very favourably with ether in this respect.¹³ Tribromethanol^{14 15} produces only a mild parenchymatous degeneration of the liver after repeated administration. With this drug, fatty changes in the liver occur occasionally. When they do occur they are very slight in the normal dog.

* Paper read in the Section of Anaesthetics at the Centenary Meeting of the British Medical Association, London, 1932.

Hepatic function is impaired so little that there is good comparison with ether and the barbiturates. Even when a liver has been purposely damaged by chloroform, avertin causes very little additional impediment.

More may be said about the liver in its activity. During anaesthesia phosphoric acid appears to leave the muscles and to sojourn in that gland until the resumption of kidney function after the recovery of an animal, when it is redistributed and partly excreted,¹⁶ and Mann¹⁷ has given it out that ether produces demonstrable effects on hepatic activity; it causes a hyperglycaemia dependent wholly on the liver, and it also depresses hepatic function in regard to the secretion of bile.

These informative data could easily be amplified: however, they may be sufficient for the present, and this fact remains, that we should not decry any one drug too much. I am convinced that even chloroform has a definite place in anaesthesia.

REFERENCES

- ¹ Casper: *Woch. f. d. Ges. Heilkunde*, 1850, xvi.
- ² Stiles, H. J., and McDonald, S.: *Scottish Med. and Surg. Journ.*, 1904, xv, 97.
- ³ Whipple, G. H., and Sperry, J. A.: *Bull. Johns Hopkins Hosp.*, 1909, xx, 278.
- ⁴ Greene, C. H., Snell, A. M., and Walters, W.: *Arch. Int. Med.*, 1925, xxxvi, 248.
- ⁵ Snell, A. M., Greene, C. H., and Rowntree, L. G.: *Ibid.*, 273; Greene, C. H., McVicar, C. S., Rowntree, L. G., and Walters, W.: *Ibid.*, 418 and 542.
- ⁶ Mann, F. C., and Boothby, W. M.: *Amer. Journ. Physiol.*, 1928, lxxxvii, No. 2, 486.
- ⁷ Bollman, J. L., and Mann, F. C.: *Ann. Int. Med.*, 1931, v, No. 6, 699.
- ⁸ Mann, F. C., and Bollman, J. L.: *Arch. Path. and Lab. Med.*, 1926, i, 681.
- ⁹ Rosenthal, S. M., and White, E. C.: *Journ. Amer. Med. Assoc.*, 1925, lxxxiv, 1112.
- ¹⁰ Rosenthal, S. M., and Bourne, Wesley: *Ibid.*, 1928, xc, 377.
- ¹¹ Koehler, A. E., Brunquist, E. H., and Loevenhart, A. S.: *Journ. Biol. Chem.*, 1925, lxxiv, 313.
- ¹² Bourne, Wesley: *Journ. Can. Med. Assoc.*, 1927, xvii, 1030.
- ¹³ Bourne, Wesley, Bruger, Maurice, and Dreyer, N. B.: *Surg., Gynecol. and Obstet.*, 1930, li, 356.
- ¹⁴ Bruger, Maurice, Bourne, Wesley, and Dreyer, N. B.: *Amer. Journ. of Surg.*, 1930, ix, 82.
- ¹⁵ Bourne, Wesley, and Raginsky, B. B.: *Ibid.*, 1931, xiv, 653.
- ¹⁶ Stehle, R. L., and Bourne, Wesley: *Journ. Biol. Chem.*, 1924, lx, No. 1, 17.
- ¹⁷ Mann, F. C.: *Anesthesia and Analgesia*, April, 1925.

A NOTE ON HEPATIC INSUFFICIENCY WITH SPECIAL REFERENCE TO EFFECTS OF DRUGS USED IN TREATMENT OF ARTHRITIS, GOUT, AND RHEUMATISM

BY

H. A. HARRIS, D.Sc., M.B., B.S.

PROFESSOR OF CLINICAL ANATOMY, UNIVERSITY COLLEGE, AND
MEDICAL UNIT, UNIVERSITY COLLEGE HOSPITAL

The urgent attention of the medical profession and of the manufacturers of reputable drugs should be drawn to the series of papers relating to jaundice, cirrhosis, and even acute yellow atrophy of the liver after the assimilation of various drugs of the cinchophen group administered in the treatment of arthritis, gout, and rheumatism. The widespread use of derivatives of cinchophen both in the frank legitimate form and in numerous patent remedies has become a serious danger. Already Weis¹ has recorded ninety-two cases of such poisoning, of which forty terminated fatally.

One is tempted to say that our knowledge of the metabolic processes in an organ is inversely proportional to its size. We now have much accurate scientific information concerning the exact function of the constituent parts of so small a gland as the pituitary, both as regards the pars anterior and the pars posterior. Anatomy, physiology, and pharmacology have kept in step during the dramatic progress of the last quarter of a century. In the case of the suprarenal gland progress has not been so

rapid, but from the days of Oliver and Sharpey on the one hand and Addison on the other we have lived to see the spectacle of the victims of Addison's disease returning from the death-bed to routine work as a result of the administration of the suprarenal extract of Pfiffner and Swingle.

In the case of so large an organ as the liver our lack of definite knowledge is astounding. Mann and Bollman² state that the known functions of the liver, named in the order in which they appear to be most easily impaired, are: (1) the ability to destroy uric acid; (2) the excretion of bile pigment; (3) the ability to prevent reaction from the administration of certain toxic substances. Mann and Bollman have rightly emphasized the dangers of talking of a functional test for the liver when the function of the liver is unknown.

Tests of hepatic function have grouped themselves around the problem of jaundice, sugar tolerance, protein metabolism, hepatic excretion of dyes, and detoxicating action of the liver on certain ingested drugs. The numerous tests depending on the excretion of bile pigment have of late been overshadowed by the van den Bergh reaction. The van den Bergh test is not purely and simply a test of hepatic function. There is more bile pigment made in the spleen and marrow than in the liver. The rate at which bilirubin is formed and the site of maximum formation are variable. The subtle differences in the pigment which determine the recurrence of a direct or indirect van den Bergh reaction are not yet fully understood. In some animals a direct reaction is obtained from the gall-bladder, and only an indirect reaction with the bile obtained from the hepatic ducts. The bile from some animals which do not possess a gall-bladder, such as the rat, never gives a direct van den Bergh reaction. The van den Bergh reaction is as inexplicably fortuitous as the Wassermann reaction—but less reliable.

The liver is concerned with the maintenance of the blood sugar at a particular level in much the same way as the kidney tubule is concerned with maintaining the salt content of the tissue fluids, the parathyroid with the calcium content, or the respiratory centre with the CO₂ content of the blood. There is no test of hepatic sufficiency in relation to carbohydrate metabolism which is reliable. Such tests involve not only the glycogen of the liver but also the glycogen of the muscles and the peculiar factors imposed upon carbohydrate metabolism by the islet tissue of the pancreas and the anterior lobe of the pituitary. Neither blood sugar level, glycogen mobilization, glucose tolerance, nor laevulose tolerance are fundamental tests for hepatic efficiency. When the function of the pancreas is impaired, as in diabetes, the liver actively maintains a higher blood sugar level, so that any further addition of glucose to the blood stream does not furnish evidence of hepatic deficiency. Conversely, after enforced starvation the rate at which injected glucose returns to pre-injection level is greatly retarded, and the degree of retardation is greater in the normal liver than in the diseased liver. The laevulose test is somewhat vitiated by the variation in the time taken by laevulose to replace glucose.

The ammonia and urea of the blood and urine, although closely associated with metabolic processes occurring in the liver in relation to protein metabolism, have failed to give reliable information concerning hepatic function as distinct from renal function. The destruction of uric acid in the normal animal seems to be entirely dependent on the presence of the liver, since no uric acid is destroyed in the absence of the liver. The complete removal of the liver of dogs produces a great increase in the uric acid content of the blood and tissues and causes the appearance of large amounts of uric acid in the urine, yet Mann says that the amount of uric acid in the urine