than his vagotonic brother? Is the peptic ulcer more common in persons of the sympatheticotonic type? Does the excessive use of an organ lead to structural changes in that organ? These are some of the problems which occur to one. The observation of J. Beattie, already referred to, is here of interest. There would appear to be a fruitful field for investigation on these lines.

MORBID ANATOMY OF THE AUTONOMIC SYSTEM

Apart from the new growths which present distinctive histological features little is known of the morbid anatomy of the autonomic system. A variety of appearances have, it is true, been described in the ganglion cells; but, as Nissl indicated many years ago, the histological picture presented by a pathological nerve cell is not distinctive of the process which produces it. Nor have attempts to study the morbid anatomy of disorders, the clinical features of which suggest their dependence upon autonomic disease, so far afforded much information. As Hurst¹⁶ points out, achalasia of the oesophagus—some eighteen cases of which have been recorded in which there was inflammation or degeneration of Auerbach's plexus in the cardiac sphincter—is the only malady definitely known to be due to disease of the sympathetic nervous system. But there must be many others so produced. Those distressing cases of thin blue caecum with a paper-like colon, of which I have seen several instances in otherwise healthy and robust young women, must, one is almost forced to believe, have a similar aetiology. Then again, the muscular wasting and trophic skin changes occurring in some cases of arthritis and the localized atrophy of the adjacent muscles of the thorax seen in some cases of phthisis are difficult to account for unless by structural changes in the autonomic nerve supply. Is it not reasonable to suppose that the autonomic system is liable to be affected by the same morbid processes which attack the voluntary nervous system and other tissues? In 1925 I advanced the view that the muscular dystrophies may be due to an abiotrophy of the sympathetic system.¹⁷ Ken Kuré of Tokio¹⁸ has since brought forward additional evidence in support of this suggestion, for he has shown not only that atrophy of muscle may be produced by sympathetic denervation, but that the histological appearances are similar to those met with in the dystrophies. I suggested further that cases of muscular dystrophy (reported in the literature by recognized authorities) in which recovery had occurred might perhaps be explained by a polyneuritis of the sympathetic, with a consequent distribution of the atrophy similar to that observed in the dystrophies.

CONCLUDING REMARKS

The purpose of this introductory paper has been to indicate some of the more prominent anatomical features and general functions of the autonomic system. I shall have attained my object if I have succeeded in arousing the interest of those of you who profess no special knowledge in a subject of great clinical importance. From the therapeutic standpoint I have emphasized the fact that sympathetic-parasympathetic imbalance accounts for certain attributes in the "make-up" of the individual, and have indicated certain factors which demand consideration in the province of individual preventive medicine. To the many diseases and disorders which have been attributed on clinical grounds to autonomic disturbance—and they are legion—I have made no reference. Among others I may mention Raynaud's disease, the syndrome of intermittent claudication, thrombo-angiitis obliterans, migraine, angina pectoris, vasovagal attacks, spasmodic asthma, excessive sweating, erythromelalgia, angio-neurotic oedema, "trophic" oedemas, scleroderma, unilateral hemiatrophies and hemihypertrophies, lipo-

dystrophy, achalasia of the oesophagus, "congenital" hypertrophy of the pylorus, spastic colon, megalocolon, certain anomalous types of neuralgia, striate rigidity, the hypothalamic syndromes, etc. These are but some of the conditions which at the moment occur to me. Each must have its particular aetiology. What a field for exploration! So long as a disorder is dependent upon functional or recoverable disturbance it comes within the province of the physician, but when it is due to disabling structural changes the possibilities of correction by surgical procedures may have to be considered. The surgeon has already achieved notable successes in the treatment of several of the disorders above referred to (Raynaud's disease, thrombo-angiitis obliterans, congenital megalocolon, etc.), and he will no doubt continue to extend his field of operation. But I would add a final note of warning. The surgical expert, before he interferes—and this applies especially to conditions of sympathetic-parasympathetic imbalance—must be very sure of his diagnostic ground. He must see to it that he attaches due weight to the mental factor in the production of symptoms, and that he is not carried away by his enthusiasm for the niceties of technical procedure.

REFERENCES

- ¹ Bell: *An Idea of a New Anatomy*, London, 1811.
- ² Waller: *Müller's Archiv*, 1852, p. 392.
- ³ Broca: *Bull. Soc. Anat. de Paris*, 1861.
- ⁴ Marshall Hall: *Diseases and Derangements of the Nervous System*, London, 1841, p. 123.
- ⁵ Eppinger and Hess: *Zeit. f. klin. Med.*, 1909, lxxvi, 345, lxxviii, 205.
- ⁶ Crile: *The Origin and Nature of the Emotions*, Philadelphia and London, 1915.
- ⁷ Cannon: *Bodily Changes in Pain, Hunger, Fear, and Rage*, New York and London, 1915.
- ⁸ Langdon Brown: *Lancet*, 1919, i, 827.
- ⁹ Hunter: *British Medical Journal*, 1925, i, 197.
- ¹⁰ Beattie, Brow, and Long: *Proc. Roy. Soc.*, 1930, cvi, 253.
- ¹¹ Cushing: *Proc. Nat. Acad. of Sciences, U.S.A.*, 1931, xvii, 253.
- ¹² Beattie: *Can. Med. Assoc. Journ.*, 1932, xxvi, 278.
- ¹³ Cannon: Abstract in *Bassoe's Nervous and Mental Diseases*, 1929, p. 234.
- ¹⁴ Quoted by McDowall in *Clinical Physiology*, 1927, p. 359.
- ¹⁵ Quoted by Carver in *Journ. of Neurol. and Psychopathol.*, 1921-2, ii, 39.
- ¹⁶ Hurst: *Lancet*, 1932, i, 673.
- ¹⁷ Bramwell: *Ibid.*, 1925, ii, 1103.
- ¹⁸ Ken Kuré: *Ibid.*, 1928, i, 441.

RADIOGRAPHIC INVESTIGATION OF LUMBAR AND SCIATIC PAIN*

BY

JAMES F. BRAILSFORD, M.D., M.R.C.S.

HONORARY ASSISTANT RADIOLOGIST, THE QUEEN'S HOSPITAL, BIRMINGHAM

(With Special Plate)

No symptom or group of symptoms as those known to the lay public under the terms "lumbago" or "sciatica" may be due to such a diversity of causes. So diverse are the lesions that it is impossible in this discussion, with the short time at our disposal, to cover the whole. The radiographic investigation should always include radiographs of the lumbar and sacral spine, pelvis, and hip-joints, and these should be supplemented by radiographs of other organs or structures according to the indications of the clinician. It is advisable to include lateral radiographs of the lumbar and sacral spine because these may show lesions which would not be suspected from the antero-posterior radiographs. I have recorded the details of a case of aneurysm of the abdominal aorta, the diagnosis of which was made only after a lateral radiograph had been taken. Antero-posterior radiographs of this patient had been made on several occasions because of the persistent

* Read in opening a discussion in the Section of Radiology at the Centenary Meeting of the British Medical Association, London, 1932.

pain in the back, but the essential lesion had not been suspected or recognized from the antero-posterior radiographs.

By means of radiography it is possible to determine the cause of the symptoms in a large number of cases, and though in many others radiography may only serve to exclude suspected lesions, the help given is unique, and no wise clinician will neglect its possibilities.

The lesion of the skeletal tissues or viscera which gives rise to the symptoms may be due to (a) congenital or developmental abnormalities; (b) trauma; (c) toxæmia; (d) acute or chronic inflammatory conditions; (e) tumours.

CONGENITAL OR DEVELOPMENTAL ABNORMALITIES

These occur in a large percentage of apparently normal individuals. In the bones of the lumbo-sacral region of the spine I found abnormalities present in about 25 per cent. of cases. Other observers have given much higher figures. These abnormalities may be present and the patient have no symptoms, consequently their demonstration should not be interpreted as the cause of the patient's symptoms without very good reasons, after all other possible lesions have been excluded. In some patients these abnormalities are the only lesions found, and operative treatment results in alleviation. Some authorities place more importance upon these abnormalities than others. Thus Willis, who has made a long study of the anatomy of the lower spine, in a recent article, after complaining of the lack of interest of the student in the anatomy of the lumbo-sacral region of the spine, says:

"This may explain the difficulties of the average physician when confronted with the problem of backache, as well as the tendency of the surgeon to turn his patient over and remove some of the dispensable parts from the abdomen or female pelvis, concerning the anatomy of which his curiosity has been more active."

The abnormalities known as spina bifida occulta, sacralization of the fifth lumbar vertebra, hemivertebrae, and spondylolisthesis are sometimes associated with intractable pain in the lower back, which is only to be relieved by operative measures.

The gynaecologist naturally considers that the majority of cases of low back pain in women are due to retroversion and retroflexion of the uterus, and to such other conditions as chronic cervicitis and prolapse, while less frequently he proves that the pain is due to pressure from large fibroid tumours, incarcerated cysts, salpingitis, and carcinoma. At a recent discussion on backache in women it was suggested by Brooke that in 80 per cent. of the cases the pain was gynaecological in origin and the remaining 20 per cent. orthopaedic. Kark is of the opinion that while these malpositions of the female pelvic organs occur in cases of low back pain, they are merely the accompaniment of structural and postural changes in the back. Unfortunately we are familiar with many other causes, and the importance of a preliminary radiographic examination is illustrated by the details of the following case.

The patient, a woman of 45, had had no fewer than four gynaecological operations on account of low back pain. With a change of medical practitioner necessitated by removal to another district, an x-ray examination was advised, and this revealed a collection of stones in one of the ureters, all too large to pass into the bladder. In another patient, whose backache had been attributed to "dropping" of the kidney, the kidney had been fixed without affording any relief from the pain: a large stone was demonstrated on a radiograph in the pelvis of the kidney, and her symptoms disappeared after its removal.

TRAUMA

The part which trauma plays in lumbar and sciatic pain is difficult to assess. Young men and women frequently sustain severe injuries to the back from which they readily

recover and no sign of injury can be detected on a radiograph. Injuries to the back in middle life would appear to be far more significant. The complicated structures and movements of the spine render it difficult without apparatus completely to rest the injured spine. Pain of a severity suggesting fracture may occur without any radiographic evidence of fracture, but subsequent radiographs taken after an interval of one or two years may show partial or complete ossification of the ligaments at the site of the painful vertebrae. Lesser degrees of trauma produce more pain in the "arthritic" spine and the spine showing ossification of the ligaments than in the normal, and the pain is more persistent. Many attacks of "lumbago" appear to originate in sudden involuntary movements of the spine. It is also known that in individuals harbouring a focus of chronic sepsis, progress towards recovery from trauma is much slower than in the healthy individual. That it is difficult to assess the importance of trauma and pain is illustrated by the histories of the following two patients.

The first, a woman, had been thrown from a horse while hunting, and had sustained a fracture of the transverse processes of four lumbar vertebrae, but after a few days resumed her normal life and habits. The second, a man, while pulling on a rope, had fallen backwards because of the breaking of the rope, and sustained a fracture of the transverse process of one lumbar vertebra; he had not worked since—two years after his accident—and he was seeking compensation for total disablement on account of pain in his back.

Where there is the possibility of a claim for compensation it is essential that a radiograph should be taken as soon as possible after any injury, and the radiograph very carefully examined for evidence of recent bone injury. That it is possible to give some idea of the age of a bone injury from the radiograph, and the importance of this, was illustrated in a case which came before the courts for compensation.

A man had received an injury to his back. He was taken to hospital the following day, when it was seen that he had a bruise over the site of the injury. A radiograph was taken and a fracture of the last dorsal and first lumbar vertebrae was diagnosed. The patient was put into a plaster jacket, kept on his back for some months, and then allowed to get up, wearing first a plaster jacket and later a metal back support. Two years after, his case came before the court on the plea of total disablement owing to pain in the back, but the case was referred for further examination. Radiographs taken one and a half years after the accident were submitted to me for an opinion. These showed fusion of the damaged vertebrae, but the appearance suggested a much older lesion. I asked for the patient to be sent to me for further radiographic examination. These radiographs showed almost identical appearances, and when I obtained, by request, the radiographs taken the day following his injury, I saw that these also showed similar appearances. Further, on inquiring into the nature of the injury, the patient said that he was loading coke and that a piece of coke fell off the railway truck and struck him on the back as he was bending down. Having weighed some of the largest pieces of coke obtainable I found that the heaviest only weighed 1 lb., and I came to the conclusion that such a blow could not possibly have produced so severe an injury. I advised the company to inquire into the man's history, because I was convinced that five to ten years previously he must have had a very severe injury. They were unable to obtain any evidence about this and the case was brought into court.

Before the hearing the man's barrister was given to understand that he had a poor case, and after a long consultation with the patient, he decided to accept a relatively small sum of money as compensation compared with his original claim. After the case had been agreed and settled in court, the man's counsel asked me what evidence I was prepared to give, and when I told him, he said that at the last consultation the man had admitted to him that he had had a fracture of his back seven to eight years before, for which he had received

a large sum of money as compensation. Apparently after receiving this first compensation money, he obtained work with another company, from which he had almost succeeded in obtaining a further large sum for the same injury.

I have previously recorded that radiographs taken immediately after an injury to a vertebral body may show little evidence of deformity, but that after an interval of a week or more, the radiographic appearance of a crushed body may be obtained, and I suggested that when there was a possibility of such an injury the vertebral bodies should be relieved of all pressure and strain by extension.

Radiographs following injury may reveal an unsuspected bone disease such as osteitis deformans (Fig. 1 on Plate).

TOXAEMIA

Absorption from any focus of chronic sepsis may, by setting up a neuritis, fibrositis, or arthritis, render the patient liable to attacks of lumbar or sciatic pain, and this is probably the most common cause of such symptoms. In a large proportion of these cases radiographs of the lumbar and sacral spine, pelvis, and hip-joints may only show pointing of the articular margins and none of the gross changes usually illustrated as examples of chronic arthritis. On the other hand, patients whose spines show marked lipping of the vertebral bodies, though they may, and usually do, give histories of past attacks of acute lumbago or sciatica, do not appear to suffer attacks of the same severity now that bone changes are well marked. It is these acute conditions which form the majority of the cases, and to which the terms "lumbago" and "sciatica" are usually understood by the lay public to apply, that help in treatment is obtained by radiographic investigation for foci of sepsis.

Men who have paid particular attention to these cases are influenced according to the incidence of the sites of the septic foci and the improvement of the patients after these foci have been eradicated. The colon, gall-bladder, appendix, teeth, sinuses, and genito-urinary sepsis receive varying degrees of attention with different authorities. The fact is that in particular patients relief is obtained by the discovery and eradication of any one or several of these septic foci; but because removal of colon sepsis, septic gall-bladder, septic appendix, septic teeth, or genital sepsis has resulted in relief of symptoms, any one of these should not be allowed to control our judgement in every case. Such obsessions have been responsible in the past for the extraction of many healthy teeth and other operative treatment. Definite lesions which have been demonstrated should receive attention before apparently normal organs or structures are removed; it is advisable to have as complete an investigation as possible before jumping to conclusions as to the cause of the pain. This is illustrated by the history of a patient who showed well-marked evidence of colitis, clinically and radiologically, to which was attributed the cause of lumbago and sciatica, but at a later examination a large fibroid was discovered, which was causing marked pressure on the structures in the pelvis. In another patient the source of the toxin causing severe lumbago and sciatica was attributed to the gall-bladder, and a cholecystectomy was advised; the radiograms, however, showed destruction of the second lumbar vertebra by secondary carcinoma. The primary, a carcinoma of the prostate, was not discovered for six months after the lesion in the spine had been demonstrated on the radiograph.

Severe pains in the back and down the course of the sciatic nerve are not infrequent in diverticulitis. Spriggs and Marxer have pointed out that

"in 100 recent consecutive cases of diverticulosis examined with this point of view, spondylitis was present in 72, whereas in 100 non-diverticulosis patients of the same average age as

the diverticulosis ones—that is, 58 years—spondylitis was present in 20."

These figures suggest that in a fair proportion of cases of lumbar and sciatic pain the toxic absorption from the colon may be responsible. The prevalence of arthritis of the lower spine and hip-joints in women is probably due largely to sepsis of the genito-urinary system.

The value of radiography in investigating the alimentary and urinary tracts and the accessory sinuses in these cases is dealt with in other papers. I shall confine myself here to dental sepsis and its radiographic interpretation in these cases.

DENTAL SEPSIS

No more characteristic lesion is detected by radiography than the abscess or focus of sepsis at the apices of infected teeth, and no more clearly related cases of cause and effect can be seen than in those instances of acute lumbago or sciatica in which the radiologist demonstrates one or more apical abscesses, and in which the painful symptoms disappear after the extraction of the infected teeth, though no response had previously been seen to any form of treatment. Unfortunately septic processes may exist at the apices of teeth from which the patient is absorbing toxins or having his blood supplied with virulent bacteria without the patient being conscious of any local discomfort, or without the teeth showing any evidence of sepsis even to the trained observer. It is most important for the patient, whose health is our first and chief care, that the doctor attending the patient is acquainted with the presence of any such focus of danger to health. The dental surgeon may know of it; he may have knowingly root-filled a septic tooth, believing that it could have no influence on the health of the patient or because he considered it of importance in holding a denture, or because the patient expressed a desire to keep it and the dental surgeon has been persuaded against his better judgement. But when the acute lumbago or sciatica or more grievous malady resulting from it comes along, this evidence is not in the hands of the doctor, and the patient, having no local symptoms, does not think of the teeth as a cause. Further, if the dental surgeon has taken radiographs, the patient and doctor are apt to conclude that these were satisfactory and interpreted correctly, consequently any septic focus has been effectively dealt with. Unfortunately only too often the radiologist discovers that this is not the case. Of the last 1,000 patients submitted to me by doctors searching for a focus of sepsis I found definite evidence of hidden sepsis in no fewer than 682.

The radiograph always shows evidence of chronic septic absorption, and it is often the only evidence of the presence of this type of dental sepsis. Further, I am convinced that if the teeth give the patient no discomfort and present no clinical or radiographic signs of sepsis, they are not the cause of chronic septic absorption and should not be extracted on account of this. Acute dental sepsis will cause the patient to seek treatment because of the local pain; it does not even require skilled training to detect it, but of the apical abscess the patient is generally unconscious and clinical examination will not detect it. It is natural, therefore, that the doctor should refer his patient to a radiologist for an independent opinion as to the possibility of hidden dental sepsis. Most dental surgeons welcome the help which a radiologist can give.

The following remarks made by a surgeon of world-wide repute who has probably seen more radiographs than any surgeon, and recorded in the *British Medical Journal* (April 23rd, 1932), is known to apply to dental radiography with even greater emphasis.

"Radiography was an essential diagnostic agent, and not only should the surgeon insist on procuring a film, but should welcome the radiologist's help in reading it. Some surgeons said

'Give me the film and I can read it for myself,' which was stupid and arrogant, and never to the patient's advantage."

Apart from the fact that the radiologist has far more experience than the practising dental surgeon can possibly obtain, he, unlike the dental surgeon, is a medical man, familiar with the dangers of dental sepsis to the general health, and, perhaps most important, his opinion is unbiased and independent. This opinion of radiography by dental surgeons is most strongly supported by those dental surgeons who have devoted much time to its study.

L. M. Ennis, D.D.S., in his textbook, *Dental Radiography*, says:

"The trouble is that many dentists consider themselves radiologists merely because they possess the apparatus. Putting it straightforwardly, no dentist can qualify as a radiologist merely from sporadic experiments after a desultory course of reading and a superficial line of instruction at the hands of the equipment salesman, and yet there are many dentists who presume to do so. There lies a danger that needs no lengthy explanation. The making of radiographs demands more than the average unthinking dentist will comprehend unless he is thoroughly versed in the requirements of both the technique of making them and the ability to interpret them. Either one of these two realms of radiography can be mastered only after a deep study of the fundamentals that demands time and patience, and the casual operator in the field cannot break into his regular practice to allow sufficient time, or patience for that matter, to become a master."

The importance of the independent opinion is still further emphasized by the fact that dental sepsis or other foci of sepsis may be present, yet may be of little or no importance in causing the grave symptoms exhibited by the patient, and the eradication of these septic foci may subject the patient to unnecessary suffering. The following is an instance. On January 1st, 1932, I was asked to radiograph the teeth of a patient who had been confined to bed for seven weeks with acute sciatica. I took the opportunity also to radiograph the lower spine, pelvis, and hip-joints. The dental radiographs showed two large apical abscesses, to which, had these been the only radiographs taken, might have been attributed the cause of the pain, and the patient subjected to the pain and discomfort of extraction. Unfortunately the radiographs of the lumbar spine, pelvis, and hip-joints showed extensive carcinoma metastases (Fig. 2), though no primary had ever been suspected. This case also emphasizes the importance of radiographing the site of the pain as well as the possible source of toxin. In another patient suffering pains in the back due to erosion of the vertebral bodies by aneurysm, the teeth were extracted because it was thought that these, being septic, were the cause of the pain.

Many instances have occurred in which radiographically normal teeth have been extracted because of lumbar or sciatic pain when unquestionable genital or colonic sepsis had been given secondary consideration or entirely neglected. Only harm can be caused by the removal of any normal structure. This fact is continually brought to the notice of radiologists when investigating cases of gastric disturbance following dental extractions which have failed to relieve the symptoms for which they were extracted. The radiographic appearances of dental sepsis and their significance I have described and illustrated in the *British Medical Journal* (June 16th, 1928).

ACUTE AND CHRONIC INFLAMMATORY CONDITIONS AND TUMOURS

The radiographic appearances of acute and chronic inflammatory diseases and tumours of the bones of the lumbar and sacral spine, pelvis, and hip-joints and their significance will be dealt with in other papers. I have previously illustrated and described a series of cases in

which the radiographs showed definite evidence of such conditions as Paget's disease of the bone (osteitis deformans), secondary malignant deposits in bone, and evidence of syphilis and tubercle which had previously not been suspected as the cause for lumbar or sciatic pain.

SUMMARY

1. In the investigation of the cases of lumbar and sciatic pain, antero-posterior and lateral radiographs of the lumbar and sacral spine, pelvis, and hip-joints should always be made. These may give an immediate clue to the cause of the symptoms.

2. Congenital or developmental irregularities occur in a large percentage of cases, and should not be ascribed as the cause of the symptoms until all other possible lesions have been excluded.

3. In cases of injury to the back or pelvis, radiographs should be taken as soon as possible after the injury, and note made of any old as well as recent injury and the films kept for reference. Changes indicative of injury may be shown on radiographs taken subsequently which were not present on the first radiographs.

4. Acute lumbar and sciatic pain is most frequently associated with toxæmia from some focus of septic absorption such as teeth, colon, genito-urinary system, appendix, gall-bladder, sinuses. Radiographs will often help in the discovery of such a focus.

5. Radiographs of teeth will show evidence of chronic septic absorption though local symptoms and clinical signs may be absent. The essential feature of the radiologist's report on teeth is that it is independent and unbiased.

6. Paget's disease of bone and secondary deposits of malignant disease may be discovered in the preliminary radiographic examination before there is any clinical evidence that such conditions exist.

REFERENCES

- ¹ Brailsford, James F.: Observations on the Value of Radiography in the Diagnosis of Obscure Dental Sepsis, *British Medical Journal*, 1928, i, 1013.
- ² Idem: Aneurysm of the Abdominal Aorta, *Brit. Journ. of Surg.*, 1926, No. 54.
- ³ Idem: Deformities of the Lumbo-sacral Region of the Spine, *Ibid.*, 1929, No. 64.
- ⁴ Idem: Diseases of the Hip-joint, *Lancet*, 1932, i, 721.

THE INFECTIVE ECZEMAS OF THE SCALP*

BY

G. B. DOWLING, M.D., M.R.C.P.

PHYSICIAN IN CHARGE OF THE DEPARTMENT FOR SKIN DISEASES,
ST. THOMAS'S HOSPITAL; PHYSICIAN TO ST. JOHN'S HOSPITAL
FOR SKIN DISEASES

The infective eczemas of the scalp are of greater importance than is generally supposed. They are seen in numbers in every large skin clinic, and in some cases present the physician with a task which under the present conditions he is quite unable to accomplish. But before attempting to discuss the rational therapy of these conditions it is essential to divide them into definite aetiological groups.

It is particularly desirable to avoid the common error of grouping them all loosely under the term "seborrhoeic eczema." Though this name is unsatisfactory, even for the disease to which it belongs, it is in fact frequently applied to a number of other conditions which are totally unrelated to seborrhoeic eczema, and especially to the eczemas of the scalp, whatever their cause may be.

It will be agreed that the term "eczema" may, however, be correctly applied to all of them; for eczema is

* Read in the Section of Dermatology at the Centenary Meeting of the British Medical Association, London, 1932.

J. F. BRAILSFORD: RADIOGRAPHIC INVESTIGATION OF LUMBAR AND SCIATIC PAIN

FIG. 1.—Radiograph showing general coarseness of bone trabeculae of left side of pelvis and upper half of femur, and increase in density. Bones of pelvis show pressure deformities; pelvis is flattened and walls of acetabulum are being pushed into pelvis. The transverse fracture of upper end of femur is characteristic of Paget's disease of bone.

FIG. 2.—Radiograph of left side of pelvis of woman aged 52, showing marked increase in density of acetabular region with almost complete absorption of lime salts from symphysis and ramus of pubis and ischium. No definite striation of involved bone of ilium indicating trabeculae of cancellous tissue can be seen, but normal contour of pelvis is preserved. Secondary carcinoma.

D. F. HEGARTY: CHRONIC DUODENAL ILEUS

Showing duodenal ileus with distended duodenum.

K. HUNTER: METAL RING ROUND TONGUE

Lateral view showing ring in situ.