2. The large numbers of polymorphs in the cerebro-spinal fluid, and the fact that in subacute cases meningococci may be very scarce.

3. Cases of meningococcal infection with an onset resembling nephritis have been described.

4. A prolonged course with relapses is one recognized form of meningococcal septicaemia.

CONGENITAL ABSENCE OF HEAD OF FEMUR

BY

ALEX BROWNLEE, M.D., F.R.C.S.Ed.

MEDICAL SUPERINTENDENT, TUBERCULOSIS HOSPITAL, GLAN ELY,
FAIRWATER, CARDIFF

(With Special Plate)

The following cases of this anomaly seem of sufficient interest to warrant record. Briefly, they are examples of congenital absence of the head of the femur on both sides in two members of the same family, with a condition resembling double Perthes's disease in a third member. The particulars are as follows:

CASE RECORD

M. M. W., a female child of 5 years, was sent to me on March 12th, 1931. She complained of slight pain in the left hip, and the mother stated that she was limping. On clinical examination nothing definite could be made out. There was certainly no sign of limping when the child was made to walk, and her general condition was good. There was no muscular wasting, and movements of the joint were practically normal. There was no tenderness anywhere, and no pain on percussion of the heel or of the great trochanter. The patient was *x*-rayed, and an extraordinary condition was seen. The head of the femur on both sides was completely absent, and a space of about 1 inch existed between the neck and the acetabulum (Fig. 1). A report was sent to the tuberculosis officer concerned, giving particulars of what was found, and stating that the condition was not tuberculous, and that she should be recommended to an orthopaedic hospital for any treatment required.

At a later date Dr. Folliott (the tuberculosis officer who had sent the case) reported that he had reason to believe that two other members of the same family were affected by some hip condition, and they were requested to attend for examination. On January 8th, 1932, the first patient's mother and brother attended for examination. Neither of them complained of any symptoms whatsoever. The mother displayed rather a "waddling" gait, and her appearance was somewhat suggestive of a congenital specific condition. On *x*-ray examination of the mother, aged 44, it was found that the head of the femur on both sides was absent, and both the necks were considerably thickened (Fig. 2).

The boy, aged 7, showed nothing abnormal on clinical examination, but the *x* rays revealed a condition resembling double Perthes's disease (Fig. 3).

COMMENTARY AND REVIEW OF LITERATURE

In the *Annals of Surgery*, 1930, this condition is reported on by Dr. Franklin E. Walton of St. Louis. He describes congenital absence of the neck of the femur as a comparatively rare anomaly and gives the history as one of a painless, sinking limp, first noticeable when the child begins to walk, suggestive of the likelihood of congenital dislocation of the hip. He further states that the usual confirmatory clinical features are—shortening of the limb, displacement of the upper trochanter, absence of the head from its normal position (below the centre of the inguinal ligament), and mild limitation of abduction, with the other movements comparatively free. None of these symptoms or signs were present in any of the cases recorded above. Walton states that the cases on record are few, but he

refers to eleven. Of these six were bilateral cases (two of which later developed coxa vara), two unilateral (one of which showed absence of the upper epiphysis and part of the shaft), one showing absence of the lower femoral epiphysis, and one extensive arrest of development of the femur.

Treves, in his *Surgical Anatomy*, states that there are three epiphyses in the upper end of the femur, one for the head, which unites at between 18 and 19 years; one for the lesser trochanter, which unites at about 17; and one for the great trochanter, which unites at about 18. The neck is formed by extension of ossification from the shaft and grows as a part of the shaft. Slovitch, in a thesis for the doctorate of medicine of the University of Paris, wrote on these conditions. He stated that these malformations were unknown until forty years ago, the first observation of them being published in 1883 by John H. Morgan. In this case there was congenital absence of the right femur. Some authors, however, doubt the occurrence of total congenital absence. Slovitch refers to the work of Reiner, who has divided these malformations into five groups. In the first are placed all the cases in which the femur is shortened and thinned—in these there is an extreme degree of congenital coxa vara. In the second the femur does not exist as a complete unit, but is divided into several fragments—below, the inferior extremity of the femoral shaft with its epiphyses; above, the head and the trochanter. In the third group the same disposition exists as in the second, but the lower extremity of the femoral shaft is fixed like a cork on to the tibia, without the formation of a knee-joint. In the fourth group certain parts of the femur form a complete unit, but the upper part of the femoral shaft is arrested in its development and retains its embryo form. The head is situated in the prolongation of the shaft of the femur; the neck does not exist. Above, the axis of the femur is bent towards the pelvis. In the last group the malformation consists of extreme shortening of the shaft. The epiphysis, hip-joints, and knee-joints are normally developed and present their normal dimensions. It is pointed out by this author that there is one characteristic found in common—namely, that the proximal part of the femur is that which shows the greatest malformation.

Blencke, who has adopted Reiner's classification, has examined sixty-three cases, adding three new ones of his own—total sixty-six. Of these, thirty were males and nineteen females; in the remaining seventeen the sex was not stated. From this he assumed that the malformation was commoner in males than in females, but this point is not yet certain. Eleven times the deformity affected the femur alone; twelve times the femur and the fibula; eleven times the femur, the fibula, and the patella; and three times the femur, the foot, and the fibula. In twelve cases the deformity affected all the bones of the lower extremity without exception, and thirty-one times the femur, fibula, patella, and the foot. Detailed observations are wanting on nine of the cases. In the great majority the patella was absent. Naturally, the extent of the deformity differed in these cases. In some, absence of the femur was complete, these being very rare, and denied by some authors. In others the fragments, more or less large, were present as vestiges, and sometimes the malformation consisted in a true shortening of the hypoplasia, the shape being preserved. These statistics show that the deformity can be either unilateral or bilateral, and can exist alone or in association with other deformities of the lower limbs, having a predilection for certain of them—in particular for certain parts of the same bone—but not absolutely sparing any part of the lower extremity.

In discussing the pathology of this condition various authors have suggested alcoholism of the parents, emotion,

infections, and injuries. Other less likely causes have been quoted as being capable of producing the deformity, some of these being associated with pre-natal obstetrical abnormalities. According to Reiner, it is logical that any external cause, whether pathological or mechanical, which localizes its action on a special vascular area of the femur in the embryo, can only interfere with the development of the particular vessel interfered with. Slovitch, in his thesis, comments on the fact that the congenital absence of the head of the femur was also associated with the absence of the fibula, and absence of several toes in one case.

Memoranda MEDICAL, SURGICAL, OBSTETRICAL

CHRONIC DUODENAL ILEUS

(With Special Plate)

I wish to report the following case, which came under my care recently, first, because the condition is a rare one; secondly, because of the excellent radiographs taken by Dr. J. Fielding, which made a confident pre-operative diagnosis possible; and finally, because operative treatment in this condition is so successful.

CASE RECORD

The patient was a female, aged 24, on whom I had operated for acute appendicitis three years previously. She stated that the present complaint had persisted for nine months. She first noticed a sensation of fullness and pain in the epigastric region, coming on a short time after meals. Then she began to vomit. This usually occurred from half to one hour after meals. Food was returned unaltered, but was usually bile-stained. The vomiting became gradually more frequent, she lost weight, and was very constipated.

On admission to hospital she was very emaciated, weighing 6 st. 2 lb. Her general condition was very poor. Cardiac, respiratory, renal, and nervous systems were normal. I was not able to make a provisional diagnosis from the clinical history. She was kept in bed for a week and put on a milk diet. The vomiting diminished. She was then x-rayed, the report being as follows:

"Stomach normal in size, tone, and mobility. No evidence of ulceration. Duodenal cap filled, but no evidence of deformity. Early on during the screen examination a lag was noticed in the third portion of the duodenum. As the passage of the meal progressed, the third and fourth portions of the duodenum were observed to be very much dilated, and little or no peristalsis was evident. The x-ray evidence is of chronic duodenal ileus."

Assisted by Dr. Fennell I explored the abdomen by a right paramedian incision. There was no ulcer of the stomach or duodenum to be seen, or any evidence of gall-bladder disease. On drawing out the transverse colon and turning it upwards the third part of the duodenum, as seen through the transverse mesocolon, was dilated. This dilatation terminated abruptly where the duodenum was crossed by the superior mesenteric vessels. The vessels were tightly stretched across the intestine at this point, greatly diminishing the lumen. I decided to perform a duodeno-jejunostomy as recommended by Moynihan. On incising the mesocolon over the selected portion of the duodenum, and on mobilizing the latter, I noticed that the coats of the intestine were much thickened. For the anastomosis I used a loop of jejunum close to the flexure. Owing to the dilatation of the duodenum the operation was comparatively simple.

Convalescence was smooth and she has not vomited since (over two months), and is gaining weight steadily.

North Infirmary, Cork.

D. F. HEGARTY, F.R.C.S.Ed.

BIBLIOGRAPHY

British Medical Journal, 1921, ii, 795.
Moynihan: *Abdominal Operations*, i, 445.

TWO UNUSUAL TONGUE CASES

(With Special Plate)

Case 1: Metal Ring Round Tongue.—This case is of interest in view of its duration and the slight inconvenience which the foreign body caused.

A little girl 6 years old came to the surgical out-patient department of this hospital complaining of a sore throat and a lump under her tongue. Her mother thought that she had enlarged tonsils. On examination a hard mass was found under the tongue, the surface of which was covered with tartar. Closer examination revealed it to be a metal ring, which, when followed round the base of the tongue, completely surrounded it. The posterior part could be felt with the finger in the pharynx. When questioned the child said that a little friend of hers had put the metal frame of the front of a watch round her tongue about one year before. Under general anaesthesia the ring was removed after dividing it in one place.

The x-ray photograph reproduced on the plate shows the ring *in situ*.

Case 2: Bilateral Congenital Tumours of the Tongue.—As congenital tumours of the tongue are rare, the publication of a short note on this case seems justifiable.

A female child 9 days old was admitted to the hospital with two swellings in its mouth which had been noticed by the maternity nurse. On examination these were seen to be two pedunculated tumours arising from the dorsum of the tongue on each side, just in front of the junction of its anterior two-thirds and posterior third. The one on the right side was the larger, being about the size of a gooseberry, and accommodated itself in a cleft in the soft palate; the left one was a little smaller. The surface of the tumours had the same appearance as the mucous membrane of the tongue. The lower gum presented a serrated appearance due to segmentation of the labiogingival lamina, a condition sometimes seen in the newborn. The tumours were ligatured and removed. On microscopical examination they presented the characteristics of accessory salivary glands.

I have to thank Mr. Ian Fraser for the permission to publish these cases.

K. HUNTER, M.B.,

House-Surgeon, Belfast Hospital for
Sick Children.

OSSIFICATION OF BOTH ACHILLES TENDONS WITH TRAUMATIC FRACTURE OF ONE

(With Special Plate)

The condition below described appears to be unique—at any rate in the extent of involvement of the tendons in the bony formation and in the results of injury.

The patient, a man 56 years of age, was born with a deformity of both feet which, so far as can be ascertained, was a certain degree of bilateral talipes equinus. At the age of 8, owing to a severe walking disability, he had had the operation of tenotomy performed. Both tendons were divided by the subcutaneous method. The result was a great improvement in his gait, though the patient had always been described as an "awkward walker." On reaching adult life he took up a vocation which necessitated a great deal of walking, but one which he has followed up to the present. About a year ago he began to experience aching pains in both heels particularly and in the feet generally. His ankles used also to swell slightly. These symptoms greatly increased in March and April, 1932, and made the carrying out of his work very painful and difficult. The left foot was especially affected. On May 2nd, whilst walking, he suddenly felt a sensation as of a heavy blow on his left heel, followed by a "fiery pain" up the back of his leg. After this he was just able to crawl home, very slowly and painfully.

The patient was first seen the next day. Examination revealed a tall man, thin, under-nourished, and of rather poor physique. Heart, lungs, and other systems were normal, but indigestion was complained of, and there was some dental caries. The patient had been living on a diet consisting

ALEX BROWNLEE: CONGENITAL ABSENCE OF HEAD OF FEMUR

FIG. 1.—Girl, aged 5. Radiograph showing absence of head of femur.

FIG. 2.—Mother, aged 44. Radiograph showing absence of head of femur and thickening of necks.

ALEX BROWNLEE: CONGENITAL ABSENCE OF
HEAD OF FEMUR

FIG. 3.—Boy, aged 7. Radiograph showing condition resembling double Perthes's disease.

F. B. MALLINSON: OSSIFICATION OF BOTH
ACHILLES TENDONS

Radiograph showing wired bony fragment of left tendo Achillis.