the vagina cannot prolapse, the uterus above it cannot prolapse, and there is therefore seldom any need to treat the uterus at all in dealing operatively with uncomplicated prolapse. It may be advisable where the cervix is lacerated, infected, elongated, or thickened to amputate it in the course of carrying out an operation of colporrhaphies—I do this in the bulk of cases. But never will I believe that an abdominal suspension operation is necessary. I am a regenerate, for, until 1922, I was well satisfied with a combination of plastic operation with ventro-suspension. My results now, by vaginal operation alone, are not a whit inferior to those of earlier years. At the same time I regret that those who agree with me in the operative treatment of parous prolapse that I have outlined have not always clearly indicated their reasons for supplementing this operation, in an occasional case, by some abdominal procedure. If they will only be persuaded to study their case-history afresh I feel sure they will realize that abdominal operation was indicated, not for prolapse alone, but for a complication of it—nearly always a subinvolution of the uterus with retroflexion. A curettage plus some suspension operation is, to my mind, an ideal procedure for this complication, but I never regard it as part and parcel of my treatment of prolapse, and on occasion I do it in cases of subinvolution where no question of prolapse is involved.

It has been argued that colporrhaphies are contra-indicated within the child-bearing period. I do not subscribe to any such view, but I admit that the carrying out of an operation of colporrhaphies to produce the desired result, without causing dyspareunia, and without leading to the likelihood of trouble at a future pregnancy, is a matter of expertness. The age of the patient must always be considered. It is very easy to cure prolapse by narrowing the vagina, but if this should lead to dyspareunia some patients no doubt would prefer their prolapse. Where there is a probability of further pregnancies, it is best to do the least that, as one judges from experience of similar cases, will cure the tendency to prolapse. But never let it be recorded that operations of this type should be postponed for either of the reasons I have mentioned. Sound obstetrical treatment will be just as necessary at any subsequent confinement as at the original one, but, with an expert operator, a reasonable interlude, and a good obstetrician, there need be no great fear of recurrence.

THE AETIOLOGY OF DENTAL CARIES*

BY

E. W. FISH, M.D.

ASSISTANT DENTAL SURGEON, ROYAL DENTAL HOSPITAL, AND
ST. MARY'S HOSPITAL

In this introduction to the discussion on the cause of dental caries some reasons are offered for considering that a peculiarity of the surface of the teeth, involving the presence of a crevice permeable to mouth fluids, is a necessary factor in producing the disease. The pathological reactions of the tooth to faults of the enamel and early enamel caries are briefly referred to. It is suggested that caries is a saprophytic phenomenon occurring either in morphological crevices or in developmental or acquired faults in the tooth surface, and that the tooth is powerless to prevent the onset of caries, though the progress of the disease may be retarded by vital reactions in the dentine. The aetiology of enamel faults and types of immunity to caries are discussed.

* Read in opening a discussion in the Section of Comparative Medicine at the Centenary Meeting of the British Medical Association, London, 1932.

PERMEABLE SURFACE DEFECTS AS PRECURSORS OF CARIES

There are four main clinical types of caries.

1. Fissure caries.
2. Interstitial caries.
3. Cervical caries.
4. Recurrent caries (after fillings).

Each of these can be shown to start in either a permeable crevice or a faulty area of the tooth surface.

Fissure Caries.—This starts in the depths of a natural fissure in the surface of the enamel. These fissures vary in their depth, and their susceptibility to caries increases with it. When unerupted teeth with deep fissures are immersed in dye, the dye is found on section to have penetrated to the bottom of the fissure (Fig. 1). The

FIG. 1.—Drawing of an unerupted human premolar exhibiting a deep permeable fissure (P.F.) which is distended in its deepest part into a natural cavity, filled with epithelial debris. The fissure was shown to be permeable by immersing the tooth in methyl blue solution for a few days.

crevice, which is doomed almost inevitably to become carious because it forms a natural incubator for germs, is present before the tooth erupts.

Interstitial Caries.—Although the contact points of two adjacent teeth constitute an area of relative stagnation, they do not offer a cul-de-sac for harbouring germs, and are not so certain to become carious as are deep occlusal fissures. Nevertheless caries is much more common at the contact point than on any other lateral surface of the crown, so that stagnation is a factor. But as all contact points do not become carious in a given mouth despite the uniformity of their environment, we may assume another factor presumably of a structural nature. We can indeed narrow the issue, for in almost every case when a tooth does become carious at the contact point its neighbour also suffers in the same way, though it is a matter of experience that the two cavities are seldom equal in size. Often one is very large and the other only just starting in the enamel. It is significant, too, that the tooth with the larger cavity is as likely as not to be the more recently erupted of the two.

Now these two teeth, so far as their contact points are concerned, have exactly the same environment. The same drop of saliva hangs between them with the same lysozymes, the same pH value, and the same carbohydrate content. The stagnation is the same and the tissue fluid in the dentine is the same, yet one decays before—or more quickly than—the other, and it is by no means always the one that was there first which decays first. It follows that the factor which accounts for the earlier onset or quicker growth of caries in one of them must be a difference of structure in the two teeth, since the environment is the same; and since a difference in structure in these circumstances implies an imperfection in the structure of at least one of the teeth, it would appear that caries starts in a fault in the enamel. These faults occur in the form of permeable lamellae and hypoplastic areas distributed at random on the surface of many teeth.

The bacteria multiply in them undisturbed, and gradually invade the surrounding enamel. It is reasonable to suppose that in interstitial caries the destructive process can spread to the enamel of the neighbouring tooth even though there is no pre-existing fault in it; but if the second tooth has a permeable fault also it will succumb all the sooner and—depending on its permeability—tend to have as large a cavity as the former. This does quite often occur, although the cavities are much more frequently dissimilar in size.

Cervical Caries.—This obviously occurs in a permeable fault in the tooth surface. The fault in this case is acquired and not developmental, for the gum has receded and exposed the cementum, which is permeable by virtue of the Sharpey fibres of the periodontal membrane embedded in it.

Recurrent Caries.—This only happens when a filling leaks, or when the enamel edge of the cavity is structurally so faulty that it becomes carious *de novo* and so falls into one of the other groups.

PATHOLOGY OF PERMEABLE ENAMEL FAULTS

It is necessary to discuss first whether enamel is capable of displaying any vital phenomena.

The enamel of a young dog is freely permeable to fluids both from the dentine and from the mouth. As the dog grows older the enamel becomes impermeable in its outer layers, probably by a process of keratinization of its organic content. Human enamel, on the other hand, is never freely permeable to fluids diffusing into it either from the dentine or from the mouth, though it sometimes contains traces of tissue fluid (Fish, 1932). Laidlaw and Jefferies, by using cataphoresis, have driven dye through the human enamel, but agree that it is far less permeable than dog's enamel, and that old dogs' enamel is far less permeable than young dogs. There can therefore be no reasonable doubt that while a fluid path exists across the enamel in human beings, it is not at all a free path, and that molecules the size of methyl blue or acid fuchsin can only pass along it in the majority of cases if they are impressed by a cataphoretic current of the order of 100 volts.

When a permeable fault exists in the surface of the enamel, it becomes filled immediately upon eruption of the tooth with mouth fluids containing food in solution and even solid food debris and micro-organisms. It is inconceivable that in the absence of any cells or blood vessels the traces of tissue fluid which sometimes enter the enamel for a short distance could, even if they reached the fault at the surface of the enamel, wash away the contained debris, seal up the fault temporarily, and then repair it. Such a service, when a permeable breach in the skin occurs, demands the presence of leucocytes, new capillaries, profuse extravasation of serum, granulation tissue, epithelialization, and fibrous tissue production.

But a more serious bar exists to the possibility of any vital reaction occurring in the enamel. Whenever mouth fluids enter a fault in enamel to a sufficient depth to irritate the dentine beneath, either the dentine hypercalcifies, producing a translucent zone which effectively isolates the lesion from the pulp, or the contents of the affected tubules die and are shut off, together with the enamel beyond, by secondary dentine, thus forming a dead tract in the hard tissues of the tooth, which is cut off by a barrier and lies completely outside the pale of nutrition. This observation (Fish, 1929)* has been corroborated by Boedecker (1930) and by Jefferies (1932). The latter shows that the barriers thus formed are still impermeable to dyes, even when they are subjected to cataphoresis.

It therefore follows that a permeable developmental fault in enamel becoming contaminated upon eruption of the tooth is not only beyond the possibility of repair, but the whole area very quickly becomes shut off from tissue fluids, together with the underlying dentine (Fig. 2).

FIG. 2.—Drawing of adult human molar. Dye was placed in the pulp chamber and allowed to diffuse into the dentine before the section was cut. The dye entered the normal dentine everywhere, but was shut off from reaching the lesions in each case by a calcific barrier. The section exhibits a permeable enamel lamella at L1 stained by mouth fluids, but non-carious, and shut off from tissue fluids by a translucent zone (TZ1). L2 is a similar enamel lamella which is beginning to become carious and is isolated by a similar translucent zone (TZ2). C1 is a later stage, and exhibits enamel caries which is invading the dentine. The original translucent zone is becoming decalcified again at C1 and the tubules are infected, but the translucent zone (TZ3) is spreading and still completely isolates the caries from the tissue fluids of the pulp. C2 is either a permeable hypoplastic fault of enamel or early enamel caries. Not being in the occlusal surface it shows a dead tract instead of a translucent zone type of reaction. The invaded tubules, being devoid of dye throughout their whole length, must be sealed off at the pulp margin by secondary dentine, and so lie beyond the reach of tissue fluid from the pulp.

Such faults therefore constitute areas of necrotic tissue in which saprophytic phenomena may occur, and the only possibility of vital reaction checking the "disease" is by means of the calcific barriers surrounding these necrotic areas in the dentine. These barriers continue to be laid down in advance of the carious process, and it is evident that if they can be built of more resistant material they will delay the ravages of caries. No doubt the enormous differences observed in the rate of spread of caries in different teeth are due in part to this cause, though obviously the suitability of the environment to the saprophytes will also have a bearing on the rate of growth of the cavity.

It is, however, clear that once a tooth has been developed with a permeable fault in the enamel surface, no vital reaction can occur in the tooth which would prevent the incidence of caries. This can only be done by rendering the mouth immune to the saprophytes responsible for the putrefactive processes which, occurring in the fault, constitute caries.

AETIOLOGY OF FAULTS IN ENAMEL

It has been shown by M. Mellanby that an absence of vitamin D from the diet during the development of the teeth in dogs always causes the enamel to be faulty, and it may be that the converse is true and that faults in the enamel are always due to vitamin D deficiency. It is, however, difficult to reconcile this with the clinical findings.

In view of the wide distribution of vitamin D in milk, eggs, butter, cheese, and animal fats generally, it seems surprising that almost the whole population, not only of this country, but of sunny countries like Australia, should be suffering from a deficiency serious enough to produce extensive caries. Yet a small percentage of the population is immune, not by any means confined to the wealthier classes, who are better housed and fed. Again, the faults in teeth which become carious are not anything like the hypoplasia of rickets, but consist of enamel lamellae and small patches of discoloured and chalky enamel on the undistorted surface of the tooth. A fault due to rickets is a distortion or pitting of the

* A more complete account is in the press, to be published in the *British Dental Journal*.

enamel, which is not necessarily "chalky" in character. The rachitic fault, moreover, occurs in certain definite parts of the tooth corresponding to a given period of growth. Caries does not. Amongst the teeth most affected by vitamin shortage in rickets are the lower incisors; yet these teeth are almost immune to enamel caries. Surely if vitamin shortage in early life predisposed the teeth to caries, the lower incisors would be amongst the most susceptible, instead of being amazingly immune.

Mackay (1931), in a report on the association of vitamin D deficiency and dental caries, showed that there was only slightly greater incidence of caries in a group of rachitic children, 22 per cent. of whom showed hypoplasia of the enamel, than in a control group of normal children in which only 2.5 per cent. showed hypoplasia.

In any case the permeable point in the tooth surface which becomes carious is quite often not a developmental fault at all, and therefore has no relation to avitaminosis. The depth of fissures is a morphological characteristic. Recession of the gums which gives rise to cervical caries is a fault of the parodontal tissues rather than of the tooth, and recurrent caries is a matter of technique. Thus at most vitamin deficiency can only account for one of the four types of caries observed clinically—namely, the interstitial group. Even so, the deficiency would have to extend over a period of about twelve years to affect all the teeth in the way that caries does, and should logically be most intense in the twelfth year, when the wisdom tooth is developing. It seems more likely that the poor calcification of the wisdom tooth is a racial retrogression.

Interstitial and fissure caries appear to be definitely racial and hereditary in character. Tamil and Chinese coolies working together show an extraordinary difference in the incidence of caries. J. W. Field (1929) recorded that 100 Tamils had seven carious teeth while 100 Chinese had 163. Their diet was similar and so was their exposure to sunlight, though the chewing by the Tamils of betel nut with its caustic content might have had an antiseptic effect on the saprophytes.

Up to the Third Dynasty Elliot Smith reports that the Egyptians were free from caries. At this time an invasion of a foreign race occurred, and caries became as prevalent as it is in England to-day.* Even on Tristan da Cunha (Sampson, 1932), where hereditary and not racial factors predominate, of the 156 people examined 26 had an average of eight teeth each, either carious or missing, while 130 were caries-free, and these 26 were largely the "foreign" element of the population, in contradistinction from the 130, who were chiefly descended from the original stock.

In clinical practice in England it is a matter of common experience that good teeth run in families, so that whatever the factors may be which produce the faults and crevices in which caries develops, they appear to be hereditary.

IMMUNITY TO CARIES

There are certain cases of this which are significant.

The human subject presents two types of immunity to caries. In one we find a perfect set of teeth with shallow fissures and faultless enamel. In the other type there are deep fissures and numerous discoloured patches of enamel which show traces of early caries, but the caries has now become completely arrested. The former type appears to have no flaws in which the organisms can lodge, the latter appears to have a mouth which has become inimical to the growth of the saprophytes concerned. If, as he gets older, the man with perfect enamel gets recession of the gums, the exposed necks of the teeth often decay, yet the enamel remains immune, showing definitely that his immunity had been due to absence of faults in the enamel.

* See page 760.

On the other hand the presence of faults alone is not sufficient to produce caries, as M. Mellanby has shown in her experiments on dogs where faulty enamel was constantly produced, but caries never supervened. Indeed, the dog is immune to caries, and yet it has both deep fissures and permeable lamellae with dead tracts of dentine under them (Fish, 1929). The same observation may be made in the human subject, for as we have seen the hypoplastic teeth of rickets are not especially prone to caries.

There is clearly a factor of susceptibility to an organism or a group of organisms without which caries cannot supervene. The apparently spontaneous arrest of caries is due to acquired immunity and not to a vital reaction in the tooth, since caries often becomes arrested in old broken-down dead roots as well as in deep fissures in vital teeth.

Normally, a change to relative immunity takes place as children grow up. No doubt this is to some extent due to the fact that most permeable faults have by this time been treated, or the tooth has been lost, but it may result in part from a change of diet from milk and sticky carbohydrates to savouries and the use of tea, alcohol, and tobacco. In the same way the use of betel nut may account in some degree for the relative immunity of the Tamil.

CONCLUSION

Thus all the evidence seems to point to caries being a saprophytic phenomenon occurring in morphological fissures, developmental enamel faults, or in the permeable necks of teeth. In no case can its occurrence be prevented by vital reaction on the part of the formed tooth, though its extension may be delayed. The prevention of decay appears to be possible only by rendering the mouth unsuitable as an environment for the saprophytes concerned, or by breeding a race free from morphological crevices and developmental enamel faults, whose gums will never recede.

REFERENCES

- Boedecker, C. F., and Applebaum, E.: *Dental Cosmos*, 1930, xxii, 1001.
Field, J. W.: *British Medical Journal*, 1929, i, p. 707.
Fish, E. W.: *Dental Record*, 1929, xlix, 151.
Idem: *British Dental Journal*, 1932, liii, 563.
Jefferies: *British Dental Journal*, 1932 (in the press).
Mackay, H. M. M.: *Lancet*, 1931, ii, 1230.
Sampson, W. E. A.: *British Dental Journal* 1932, p. 397.
Smith, G. Elliot: *Archaeological Survey of Nubia*, 1907-8, ii, "Report on Human Remains at Cairo."

THE AETIOLOGY OF DENTAL CARIES*

BY

MAY MELLANBY

My only claim to address you is that of one who has devoted many years to physiological and biochemical research, chiefly in relation to teeth and their associated tissues. My approach to the subject of dental disease is therefore somewhat different from that of the clinician, and though we must ultimately arrive at the same destination, we travel by different roads. As none of us can claim to have reached the goal, it is perhaps only natural that we should each tend to emphasize the points elucidated in the course of our own work. No doubt the very fact that Dr. Fish and I have approached the subject from different angles and hold such opposing views prompted the committee to invite us to open this discussion.

* Read in opening a discussion in the Section of Comparative Medicine at the Centenary Meeting of the British Medical Association, London, 1932.