vascularized." And again, "Of 7 that remained insensitive, 5 were clear, 1 was opaque, and 1 was partially opaque, after an average time of thirteen months."

The margins of the graft become sensitive before the centre, which is in accord with my findings and with what would be expected.

NEURO-PARALYTIC KERATITIS

Leslie Paton, with characteristic industry, discusses seven possible explanations of the development of neuro-paralytic keratitis,¹⁶ and comes to the conclusion that no complete and satisfactory explanation is forthcoming. I do not propose to discuss an eighth possible explanation.

The chief controversy ranges round the question of the existence of trophic nerves, the destruction of which is responsible for the ulceration. Sherrington says, "Every nerve and nerve centre that can influence the proper and specific activity of a tissue possesses in so far a trophic influence on that tissue. If by trophic nerve be understood no more than this, all nerves are trophic nerves. That, on the other hand, any nerve directly modifies the nutrition of a tissue independently of inducing vascular changes in it, or influencing those particular metamorphoses of material for which the tissue is the machine specifically adapted, there is no proof." Duke-Elder suggests, in his recent excellent book, that the formation of histamine in the tissues is under the control of impulses passing antidromically down the fifth nerve; when therefore this control is eliminated the changes in the blood vessels and capillaries of the tissues, the so-called triple response of Lewis, no longer take place, and the so-called trophic changes in the tissues develop as a consequence.

Whatever may be the truth in this regard, there are certain facts that are clear.

Anaesthesia of the cornea alone will not produce ulceration, for, if the cornea be completely protected by a tarsorrhaphy, it does not develop. Further, after it has developed it is surprising how quickly healing takes place if tarsorrhaphy be done. Again, although, as has been said, a corneal section causes anaesthesia, I have not seen an ulcer arise on this account, though it is well known that exposure without anaesthesia readily produces ulceration. The patient who lies in a semi-conscious condition with the lids only partly closed, so that the lower part of the cornea remains exposed, is liable to ulceration of the exposed area. The patient with prominent eyes from Graves's disease or an animal with prominent eyes, such as the Pekinese dog, is predisposed to corneal ulceration from exposure. No doubt the insensitive cornea leads to a reduced reflex secretion of tears and frequency of blinking, and so provides an additional factor. My belief is that, when the ganglion is destroyed, the lids should at the same time be sewn together. I do not forget that, with care, no serious involvement will take place in a good many instances, and where it does occur, in spite of bandaging, I believe it is either because the patient's cornea comes into contact with the dressing, which, not being felt, is not resented, or that the patient rubs his eye. Nevertheless, tarsorrhaphy is so easily done—if done neatly it will not show the slightest deformity of the lids after separation—and is so certain in its results that I believe it should be done as a routine. My practice is to make raw the juxtaposed edges of the lids for about a quarter of an inch and a little to the outer side of the middle, avoiding the lash-bearing margin, and to join them by two silk stitches. I believe the bridge should be maintained for six months, but even after this period ulceration may take place on separation, and compel a resuturing.

I should now like to refer shortly to a special type of ulceration in which I believe a neuropathic element exists. Cases of it have been reported by Messrs. Neame, Balfour Barrow, Herbert Fisher, de Courcy, and myself.¹⁷ During the war we were all familiar with the curious milky-white

coagulated appearance of the conjunctiva which resulted from "mustard gas" poisoning. Now, many years after exposure to this gas, one still sees a condition which counterfeits very closely the appearance that was produced by the gas at the time of exposure. There is the same milky-white appearance of the conjunctiva, over which vessels run, producing a marbled effect. These vessels are in great measure characteristic, in that they are of a very irregular calibre, being now narrow, and now becoming widened out into broader, irregular channels, only again to narrow down. In some cases no further lesion occurs, but in some the cornea becomes ulcerated, and this may occur for the first time ten or twelve years after exposure to the gas. Ulcers extend deeply into the interstitial tissue, separating its layers, destroying much tissue, sometimes producing hypopyon, showing little tendency to heal, and leading finally to extensive and dense nebulae. A striking feature is the mild degree of the symptoms produced; pain, lachrymation, and photophobia being conspicuous by their almost complete absence. That these ulcers are due to the gassing no one who has seen them will doubt, and it seems to me that the clinical course they run is suggestive of a neuropathic cause for them—a suggestion which gains support from the fact that tarsorrhaphy helps markedly in their healing.

REFERENCES

- ¹ *Brit. Journ. Derm. and Syph.*, October, 1917.
- ² *Royal London Ophth. Hosp. Reports*, 1866, v, 191; vi, 181, 263.
- ³ *British Medical Journal*, 1866, i, 470.
- ⁴ *Brain*, 1900, xxiii, 353.
- ⁵ *Trans. Oph. Soc.*, 1923, xliii, 659.
- ⁶ *Annals of Ophthalmology*, 1915, xxiv, 693.
- ⁷ *Trans. Oph. Soc.*, 1928, xlviii, 179.
- ⁸ *Royal London Ophth. Hosp. Reports*, 1869, vi, 183.
- ⁹ *Ohio State Med. Journ.*, December, 1913.
- ¹⁰ *Journ. Amer. Med. Assoc.*, 1909, liii, 1456.
- ¹¹ *Royal London Ophth. Hosp. Reports*, 1869, vi, 3.
- ¹² *Amer. Ophth. Soc.*, June, 1919.
- ¹³ Mayou: *Trans. Ophth. Soc.*, 1917, xxxvii, 226.
- ¹⁴ *Ibid.*, 1918, xxxviii, 153.
- ¹⁵ Penman, G. G.: *Brit. Journ. Ophth.*, 1931, xv, 585.
- ¹⁶ *Ibid.*, 1926, x, 305.
- ¹⁷ *British Medical Journal*, 1929, i, 497.

OCULAR MANIFESTATIONS OF LESIONS OF THE FIFTH CRANIAL NERVE *

BY

C. M. HINDS HOWELL, M.D., F.R.C.P.

PHYSICIAN TO ST. BARTHOLOMEW'S HOSPITAL AND TO THE NATIONAL
HOSPITAL FOR THE PARALYSED AND EPILEPTIC

The eye receives its sensory innervation by the first or ophthalmic division of the fifth cranial nerve. Afferent impulses pass this route to the Gasserian ganglion, thence by the central root to the pons. Here some of the fibres end in the chief sensory nucleus, but a mass of fibres descends through the pons and medulla to the upper cervical segments of the spinal cord. These form the so-called ascending or spinal root of the fifth nerve, and make connexions with a nucleus—the substantia gelatinosa of Rolando, which extends in a corresponding manner from the pons to the upper cervical cord. From the sensory nuclei a second afferent neuron arises which decussates to form the trigeminal fillet, passing on the lateral aspect of the mesial fillet, and so to the optic thalamus, whence a third afferent neuron passes via the internal capsule and corona radiata to the cortex.

The fifth nerve fibres are thus exposed to injury over a very wide area; they may be damaged in their supra-nuclear path, at the level of the nuclei, and in their peripheral distribution. A constant result of damage to

* Read in opening a discussion in the Section of Ophthalmology at the Centenary Meeting of the British Medical Association, London, 1932.

the fibres which carry afferent impulses from the eye is desensitization of the globus oculi, with immediate loss of corneal reflex; there must also be loss of function in the afferent fibres from the extrinsic muscles of the eyeball. Whether this results in any incoordination of muscular movements I cannot say; ordinary clinical observation would not suggest that it does. I have never observed any abnormality in ocular movements in cases in which there has been complete loss of sensation in the area supplied by the first division of the fifth nerve. The chief danger which results from desensitization of the globus oculi is corneal ulceration. The questions which I would submit for your consideration are:

1. Is corneal ulceration more likely to occur according to the part of the afferent path which is affected?

2. What may be done to prevent such ulceration from occurring, and if it occurs what methods should be applied to deal with it?

3. What are the ultimate results of ulceration likely to be?

With regard to the first question, I think we must agree that the part of the afferent path which is affected is of great importance in determining whether a consecutive effect on the eye will develop. It is true that in whatever part of their path the ocular fibres are destroyed, desensitization of the eye with loss of corneal reflex will result. In such a case there is an obvious danger from foreign bodies entering the eye unperceived, and thereby setting up inflammation, which is likely to result in corneal ulcer—a state of affairs which might be designated as traumatic ulcer.

I have never seen what for purposes of convenience I will call "spontaneous corneal ulcer" develop from a supranuclear or intramedullary lesion, whatever its nature—whether slow in destructive quality, such as a pontine tumour, or acute, as in a vascular lesion. It is when a lesion affects the first afferent neuron in its extramedullary portion that spontaneous corneal ulcer is likely to occur. Further, I think that acute injury—such as trauma from the surgeon's hands; acute inflammation, as, for example, in herpes affecting the Gasserian ganglion; or damage caused by the injection of alcohol—is more likely to produce a corneal ulcer than when the peripheral nervous tissues are subjected to a slower destructive process, as, for example, by pressure from tumours.

I have treated a considerable number of cases of tic douloureux by alcohol injection. For the purposes of this discussion I have looked up my cases in the last ten years; I have to thank Dr. Fordham, my late house-physician at St. Bartholomew's, for collecting my cases there. I have records of sixty which I have injected with alcohol in that period.

Tic douloureux rarely affects the first division of the fifth nerve primarily, and I am always anxious to avoid desensitizing the eye. Nevertheless, as it is of the utmost importance, if recurrence of the tic is to be avoided, to destroy thoroughly the cells of the Gasserian ganglion supplying the second and third divisions, it frequently happens that the cells of the first division are damaged as well. This is not the place and I do not now intend to discuss the relative merits of operative treatment as compared with alcohol injections of the ganglion in cases of tic douloureux, but modern surgical methods certainly have the advantage of more easily avoiding the fibres of the first division, and in consequence corneal ulcer should occur less frequently after surgical operation than after the use of alcohol.

Of my sixty cases, I find that thirty-six had complete anaesthesia in the first division of the fifth nerves as the result of the alcohol injection; sixteen of these developed corneal ulcer. As a result of my observation in these cases, I find that the period of greatest danger of corneal ulcer supervening is in the first fortnight following injec-

tion. After that period the risk steadily diminishes, and after a month or so the risk is quite small. So far as I have been able to ascertain, only two patients with anaesthesia of the first division out of twenty who escaped ulcer immediately following injection have developed corneal ulcers or any other ocular trouble at a later date. One of these cases was caused by a "splash of lime" getting into the eye when a room was being whitewashed. I do not know what occurred in the second case.

The question naturally arises as to why there should be this tendency to inflammation and corneal ulceration in the early days following injury to the peripheral fifth neurons. The course of events suggests that it may be due to irritation of fibres—perhaps the antidromic fibres described by Bayliss in posterior roots. If such is the explanation the irritation must be due to degenerative changes in the nerve fibres. It might be argued that alcohol produces destruction of some ganglion cells and irritation of others, and that the ulcer develops as the result of irritation of nerve cells by the alcohol. This, however, cannot be an adequate explanation of the cause of ulceration, because removal of the ganglion, including the cells of the first division, is likely to produce the same risk of corneal ulceration. For that reason Horsley was accustomed to stitch the lids over the cornea before removing the ganglion, or twisting the central root out of the pons. If the ulceration of the cornea is due to peripheral nerve irritation it should occur whether the lids are stitched or not. I think, however, it would seem probable that careful protection of the eye does minimize the risk of ulcer developing, and that there is an ocular factor in the shape of an external trauma, which may be very slight, but which will aggravate the tendency to inflammation or ulceration in the early days after operation.

I have never seen any affections of the deeper structures of the eye following injury of the ganglion by alcohol. Herpetic eruptions occur occasionally, but I do not recollect having seen any in the distribution of the ophthalmic division. In herpes ophthalmicus another example is afforded of damage to ganglion cells, whether by haemorrhage or inflammation, resulting in conspicuous cutaneous changes in the peripheral distribution of the first division of the fifth nerve. So far as the eye is concerned, corneal ulceration takes the place of the cutaneous vesicular eruption. I believe that the iris is sometimes involved and an iritis produced, but of this I cannot speak from personal observation; as Mr. Foster Moore has dealt fully with this aspect of the question I do not propose to say more about it myself.

A point to which I wish to refer briefly is the effect of intensive x-ray therapy on a desensitized eye. This question is one of some importance in cases of malignant disease—usually sarcoma of the base of the skull—in which the eye has become insensitive either by the direct action of the tumour on the Gasserian ganglion or from the destruction of the ganglion by alcohol injected for the relief of pain. In any case, the reaction to x rays in the desensitized eye is much more acute than in the eye with a normal nerve supply. Dr. Levett, who has charge of our deep x-ray therapy department at St. Bartholomew's, tells me that a dose of x rays which will under ordinary conditions of innervation produce only slight injection of the conjunctiva, will produce an acute chemosis in a desensitized eye. From my limited experience, it would seem that malignant ulcers progress more rapidly in a desensitized area. I remember a case of rodent ulcer involving the orbit which progressed with devastating rapidity after I had produced an anaesthesia in the area of the first division of the fifth nerve.

What may be done to prevent ulceration from occurring in a desensitized eye? As I have already mentioned,

Horsley used to stitch the lids together before removal of the ganglion. I do not adopt this procedure as a routine because more than half my patients with anaesthesia have escaped without any ulceration. Mr. Foster Moore has frequently rebuked me for not stitching the eyelids—in fact, he does so whenever I have to invoke his aid. Perhaps he is right, but I am anxious to hear the views of others as the result of their experience. I used to irrigate the eye daily, keeping it carefully bandaged for the first fortnight or so, but I no longer do this. A plan that seems to work very well is to keep the upper lid carefully strapped down for two or three weeks, then to allow the lids to open, but to put an eye shield on, which is worn at first both indoors and out, then only outdoors, and finally removed altogether.

The first sign of commencing trouble is injection of the conjunctival vessels, and if this persists I always get the lids stitched at once. If an ulcer has already formed—and they form very quickly—it will be a very superficial affair at this stage, and will heal as a rule without leaving any corneal opacity. I usually leave the lids stitched together for a period of four to six months in those cases in which an ulcer is developing.

Let me repeat that in this series of cases corneal ulceration has occurred within a short interval in 45 per cent. of those individuals in whom a complete anaesthesia in the distribution of the first division of the fifth nerve has been produced. I do not know whether my experience is similar to that of others, and the figures are not large enough by themselves to establish any theory, but the suggestion which arises from a consideration of these figures is that the tendency to ulceration in the cornea desensitized by injury to the ganglion is not due to trauma alone. If there is a nervous factor involved it is difficult to define the manner in which it operates. I do not think that there is any evidence which would support the theory of a trophic nerve influence, but such cases as these we are considering, perforating ulcers in tabes dorsalis, and bed-sores arising with startling rapidity in acute myelitis, prove that interference with the sensory supply of a part renders the tissues peculiarly susceptible to degenerative or destructive processes.

INTRAVENOUS UROGRAPHY IN CHILDREN*

BY

CECIL G. TEALL, M.B., CH.B.

RADIOLOGIST, CHILDREN'S HOSPITAL, BIRMINGHAM

The radiological investigation of the urinary tract in children has in the past been far from satisfactory. The introduction of retrograde pyelography brought about an enormous improvement in the value of the investigation of the supravescical portion of the urinary tract in adults, and in them it has become a routine method of examination, but in children, for a number of reasons, it is by no means possible to use it as a routine method.

DIFFICULTIES IN RETROGRADE PYELOGRAPHY

While it is true that in the hands of a skilled urologist the technique of the examination does not present any insuperable difficulty, many surgeons who find no trouble in the procedure required in adult patients find a considerable amount in the case of children—it is, in fact, a procedure calling for special skill and training. This, in the first place, is a very definite reason why retrograde pyelography cannot be used as a routine method of investigation in children. When the technical difficulties have been overcome, however, there still remain other objections to this method of examination. Even the slight

trauma resulting from the skilled manipulations of the urologist is liable to be followed in children by much more severe general disturbance than is the case with adults, while as a result of catheterization of the ureters children also seem to be much more prone to develop ascending infections of the urinary tract. There is, further, the fact that in children the procedure calls for general anaesthesia. This not only makes the production of satisfactory radiographs free from respiratory movement much more difficult for the radiologist, but from the surgeon's point of view it is also more difficult to know when the kidney pelvis is adequately filled without being over-distended. Lastly, there is the fact that even when this method of investigation is advised permission to proceed with the examination is frequently refused by the parents.

VALUE OF INTRAVENOUS METHOD

The introduction of a method of visualizing the urinary tract without having to resort to instrumentation was therefore a matter of considerable importance to paediatricians. Interest was at once aroused when in 1929 Zwick, working in von Lichtenberg's clinic, used uroselectan for this purpose, and with it was able to show that intravenous urography was not only possible, but was a practical clinical method of investigation. Although for a time opinions were in doubt as to the diagnostic value of intravenous urography as compared with retrograde pyelography in adults, it was at once obvious that, whatever its limitations, it was of considerable value in children, for it would at least give some information which had in many cases been previously unobtainable. This was in the first few weeks after the introduction of uroselectan: further experience has shown that its value is by no means small, and that in intravenous urography we have a method of investigation of the greatest possible value.

That the method of examination has been of practical clinical value is shown by the fact that while in the x-ray department at the Children's Hospital, Birmingham, in the two years from 1928 to 1930, only six retrograde pyelographic examinations were made, during the last two years fifty patients have been investigated by the intravenous method. The technique is quite simple to apply in the case of a child, especially since the introduction of uroselectan B, which has been in every way an improvement on the other preparations, uroselectan and abrodil, which had been in use previously. This substance causes no general disturbance, and with no more upset to the child than is occasioned by an intravenous injection, urograms of great practical value can be obtained—in very marked contrast to the procedure necessary to produce pyelograms by the retrograde method. I do not suggest that the intravenous method has replaced the retrograde method, for the limitations of intravenous urography are the same in children as in adults. As in adults, the two methods are complementary and not mutually exclusive, but in a large number of cases the intravenous method will give all the information that is required, and since it can so easily be used without upsetting the patient, it is often employed where one would hesitate to make a pyelogram by the retrograde method.

TECHNIQUE

The technique of the examination is extremely simple. It consists of the intravenous injection of uroselectan B. The dose used varies with the age of the child. For a child of 5 to 7 years we give a third of the adult dose, with a reduction for younger children, but in no case have we injected less than a quarter of the adult dose—namely, 10 grams of uroselectan. No case has shown any untoward reaction, nor has any patient been in any way upset to a greater extent than is the case with any intravenous injection. The first film is made almost as soon as the injection is finished, certainly within the first ten minutes, and we find it best to inject the uroselectan in the x-ray department with the child on

* Read in the Section of Radiology at the Centenary Meeting of the British Medical Association, London, 1932.