on the right side and the median opening on the left. No urine was seen coming from the median opening on the right side or from the outer orifice on the left. Catheters were passed without difficulty up all four ureters.

DIFFERENTIAL URINE ANALYSIS

Catheter Specimens from Right Kidney

Right Outer Ureter.—Urine somewhat blood-stained; specific gravity, 1012; acid; urea, 1.16 per cent. White blood cells +, red blood cells ++. Gram-negative bacilli and a few streptococci present. Culture yielded a heavy growth of *B. coli*.

Right Median Ureter.—Urine slightly turbid; acid; urea, 1.63 per cent. White blood cells +, red cells ++. Gram-negative bacilli and a few streptococci present. Culture yielded heavy growth of *B. coli*.

Catheter Specimens from Left Kidney

Left Outer Ureter.—Urine turbid and very blood-stained; faintly acid; specific gravity, 1010; urea, 0.70 per cent. Very few white blood cells, and many red cells. No organisms seen. On culture there was a heavy growth of *B. coli*.

Left Median Ureter.—Urine turbid; amphoteric; specific gravity, 1006; urea, 0.43 per cent. Many white cells but very few red cells present. There were some Gram-negative bacilli. On culture there was a heavy growth of *B. coli*.

The urine from all four catheters correspond closely, all being heavily infected and showing a low urea content, the right side being a little the better.

RESULTS OF TREATMENT

As the patient did not appear to benefit under the usual urinary antiseptics, a trial was made of the ketogenic treatment. At first the organisms per cubic centimetre of urine diminished from being uncountable to about 1,000 per c.cm. at the end of a week, but at the end of a fortnight a sudden increase took place, and the daily count varied from 15,300 per c.cm. to approximately 100,000,000,000 per c.cm. After about four weeks, therefore, the diet was discontinued, as it was clearly doing no good.

During the ketogenic treatment one or two evening rises of temperature occurred, associated with some headache, otherwise the general condition of the patient seemed unaffected. She was subsequently discharged to the country branch of the hospital at Tadworth, where her general health improved considerably, though the condition of the urine remained unchanged.

MONILETHRIX*

A GROUP OF TWENTY-TWO CASES

BY

J. GOODWIN TOMKINSON, M.D.

McCALL ANDERSON MEMORIAL LECTURER IN DERMATOLOGY,
UNIVERSITY OF GLASGOW; VISITING PHYSICIAN (SKIN
WARDS), WESTERN INFIRMARY, GLASGOW

(With Special Plate)

In emphasizing the infrequent incidence of monilethrix O'Donovan¹ cites the experience of J. Schutz, who, during seventeen years' hospital practice in Frankfort, saw only one case of this affection. Up to the present time I am unaware of ever having encountered the condition save in the group, including five generations, which forms the subject of this communication.

THE SON

Early in May of this year a boy, aged 5½ years, was sent to me as a possible sufferer from tinea tonsurans. His hair, however, had not been examined microscopically. At first sight the scalp posteriorly certainly suggested ringworm, but the furfuraceous scaling which so often accompanies that

disease was not in evidence. About the occipital protuberance, which was sparsely covered, and suboccipital region stumps of hairs—some of which were twisted—and short hairs were seen, while keratosis pilaris was well marked on the nape (Fig. 1 on Plate) and very slightly present on arms and forearms. His father, who accompanied him, strenuously asserted it was not ringworm, and said that he himself was similarly affected, as was his small daughter, aged 7½ years, who was with them. Microscopical examination of hairs from the boy's scalp revealed the pathognomonic features of monilethrix. The boy has to have his hair cut in front only. The macroscopical appearance—stumps of hairs, some of which were twisted, and short hairs, distributed in a patchy manner with keratosis pilaris roughly simulating a mycelial sheath—might quite well have led a busy physician to a hasty diagnosis of tinea tonsurans.

HIS FATHER

Some weeks later, after several unsuccessful attempts owing to the father's occupation as a miner, I saw the three subjects for further investigation. Microscopical examination of the father's scalp hairs showed profusion of nodes in the occipital ones. Frontally, where the hair was longer, they appeared to be more elongated. Axillary hair showed some irregular variation in diameter. Hair from the arm exhibited nodes, and in one hair from that region the medulla was broken up into distinct islets, while at one area between two islets some narrowing of the capillary calibre was suggested, raising the question of whether the malformation might originate in the medulla. Hair from the pubis showed definite nodes. No nodes were detected in lanugo hairs from the trunk. The affection in this subject was situated, in so far as short hairs are concerned, mainly posteriorly and on the vertex. He stated that the hair anteriorly, although much longer (two inches at least) tended to break, that he had only once had it cut—while recuperating after a war wound. At the time of examination infiltrated erythematous elevations were noticed on the nape, which he stated were the result of midge bites. As pediculosis capitis was detected, however, it is quite possible that they were due to the latter condition, and were urticarial in nature. No keratosis pilaris was present except on the thighs. With some hesitation a biopsy was suggested, to which he most readily and generously agreed.

THE DAUGHTER

On his daughter's scalp short hairs occurred mainly in the suboccipital and extreme frontal regions; some of the hairs were twisted and were very easily broken. The pigmentation in some parts appeared to concentrate half nodally and half internodally. Keratosis pilaris was profuse on the nape and occiput, and also present on the thighs and legs, and very slightly on the arms and forearms. Pediculosis capitis was noticed, as was abortive impetigo contagiosa on the face. This was also evident in the son.

FAMILY HISTORY

The father told me he had other relatives similarly affected. Their family physician kindly assembled them, and, accompanied by my house-physician, I visited them, removing hairs from the scalps of fourteen other persons, varying in age from 1½ to 77 years. Microscopical examination of the scalp showed pathognomonic hairs in every case. The septuagenarian—the grandfather of the boy sent for diagnosis—said that both his father and grandfather had been similarly affected. His alopecia was almost absolute. He wore a wig. He was the only one of seventeen children to be affected. His father, who had the disease, had never spoken of a sister or brother, and was probably an only child. His grandfather also had the disease; here again there is a doubt as to whether he had brothers or sisters. He himself has had fifteen children—nine boys and six girls. At birth they all had good heads of hair. Three of the daughters died in infancy, aged respectively 12 months, 18 months, and 3 years—all with good heads of hair. Five out of the nine boys developed the condition, and also one out of the

* Read in the Section of Dermatology at the Centenary Meeting of the British Medical Association, London, 1932.

three surviving girls. All his affected children eventually married. Of these:

A son, aged 55, has four children, two boys and two girls; all are affected.

A son who was killed at the age of 47 left four children, two boys and two girls, of whom two girls are affected. Both wear wigs.

A son, aged about 46, has five children, four boys and one girl, of whom one boy and one girl are affected.

A son, aged 44, has three children, all boys, of whom one boy is affected.

A son, aged 37, has five children, three boys and two girls, of whom one boy—the suspected tinea tonsurans case—and one girl are affected.

A daughter, aged 35, has had three children—two boys, and one girl who died. The two boys are affected.

early as the sixth month and in others at dates varying from this up to the fourth and fifth years. One parent states that the children (two) were born with hair which faded away at the age of 6 months, and this was followed by keratosis pilaris. This is of interest, as it has been advanced that keratosis pilaris determines the development of nodes. Certainly it is a very frequent accompaniment, for of the eleven children of the fifth generation seen by me ten had it on the scalp. I must admit that the solitary exception, a boy of 1½ years, was very refractory under examination. Of the six adults examined only three had the disease on the scalp, and one of these only slightly. From its almost invariable presence in the children one might deduce that the keratosis pilaris of the scalp tends to disappear with advancing years. Hairs

FAMILY TREE.—Showing Members Affected with the Disease

Of this series of thirteen children seven are boys and six are girls. In the whole group of twenty-two subjects occurring in five generations I have seen seventeen, and microscopically examined affected hairs from all their scalps. I have seen portraits of four of the remaining five subjects. From these data it would appear that in the later generations there had been increase in the frequency of incidence of the affection. In the third generation the percentage was 5.88; in the fourth 40; and in the fifth 54.16. I might not be allowed, in the event of further inheritors of this condition, to note their numbers in the next generation, but it would be interesting to observe whether the percentage of sufferers continued to increase.

All the adult males from the most remote have followed the occupation of coal-miner, and the affected daughter of the fourth generation married a miner. The extent and intensity of the disease in the scalp varies from almost complete baldness—as seen in the septuagenarian—to a scalp well covered, except suboccipitally and in the extreme frontal region observed in the sister of the boy whose scalp suggested ringworm. Wigs have been or are worn by five of the affected—two men and three women.

It was noticed by his mother that the hair of one youth, now 15, whom I saw, showed at the age of 14 years signs of improvement of growth. This would appear to be coincident with the advent of puberty, a period associated with increased glandular activity, which obviously introduces the question of an aetiological role sustained by the sebaceous glands, which are so intimately involved with the root sheath in the development of the embryo.⁶ The strong hereditary and familial features of the disease are well brought out in this group of successive cases in five generations with six out of fifteen children affected in the fourth generation, and thirteen out of twenty-four in the subsequent one.

DISCUSSION

Solitary cases of monilethrix have been recorded. Stelwagon and Gaskill² mention that Gilchrist Morrow and Ruggles describe cases limited to the legs and also cases which appear in adolescent or adult life. In this matter of date of onset Sabouraud³ states that the affection begins by a fall of hair about the sixth week. I have found some difficulty in getting exact information as to onset in the fourth and fifth generations, but roughly it would appear to have been in some cases as

from the scalps of the first three subjects were sown on Sabouraud's medium and colonies of *Sarcinae* obtained in each of the three tubes, in one of which a culture of *Staphylococcus albus* also appeared. Attempts at culture were made in four other cases, but at a considerable interval after removal of the scalp hairs. A yeast and *Staphylococcus albus* resulted in one case, and in another a growth of *Penicillium* type. One of the male adults, aged 46 years, developed at the age of 10 years right cervical tuberculous adenitis, and on starting to work in the coal pit wore a band on the neck to which his lamp was attached. Lupus vulgaris appeared on the right side of the neck and extended to the nape. One youth aged 16 suffers from right internal strabismus.

I have seen it recorded that in monilethrix an affected hair is involved from root to extremity. I do not think

Photomicrograph of monilethrix hair, showing nodes, with concentration of pigment in one-half of a node and continuous internode.

that this invariably occurs, as I have examined a pubic hair which appeared to be affected in only a part of its length. While all observers state that fracture of the hair occurs at an internode, there is some variance of description in the matter of pigmentation. Most describe a concentration of pigment in the nodes with corresponding depigmentation of the internodes. Max Joseph⁴ states, however, that the nodes appear lighter than the internodes in hair of slight pigmentation owing to the presence of air in the medullary canal of the nodes. This optical effect, he states, disappears in deep black hair, where the appearances are reversed. Microscopical examination, however, of grey hairs from the grandfather in this group showed pigmentation of the nodes. It has been stated that the pigmentation contrast is due

to the difference in diameter of the nodes and internodes. This would appear to be an inadequate explanation owing in some instances to the extreme degree of internodal depigmentation, and I have more than once seen concentration of pigment in areas composed of contiguous halves of nodes and internodes (see Figure in text).

An arresting feature seen in an arm hair was the separation of medulla into distinct islets, a possible abortive attempt at node formation, for at one point narrowing of calibre was suggested. In some hairs twisting of the internodes was detected, but it was decided that this was sometimes only apparent and due to an unequal distribution of pigment in the internode.

Aetiology of the Disease

This still remains very debatable. Darier⁵ quotes a theory, for which Golay appears to be responsible, that the formative papilla undergoes alternate dilatation and atrophy comparable to a pulsation of which the rhythm is daily. Other observers, however, state that the nodes represent two days' growth. Certainly the process arises intrafollicularly, as the photomicrograph of a section of skin from the scalp of the father of the boy who was suspected of having tinea tonsurans, containing three nodes within the follicle, clearly demonstrates (Fig. 2 on Plate). The almost invariable association of monilethrix and

keratosis pilaris is arresting, but would tend to point to a common aetiology rather than that the latter was the cause of the former, for monilethrix may precede keratosis pilaris. Again, keratosis pilaris arises from hypertrophy, whereas the abnormality in monilethrix lies, not in the node, which is of normal diameter, but in the atrophied internode. It will be recalled that attention was drawn to the occurrence of islets of medulla, and that this might be an early stage in the development of nodes. It should also not be forgotten that improvement occurred in one case at puberty, with its increased glandular activity, which might include the sebaceous or other glands in the aetiological picture.

These various features open up a large field of possibilities into which I do not at present intend to venture further. Investigations are not by any means completed, and I hope to resume them in the near future.

My best thanks are due to Dr. Donald Douglas, Mr. Charles Bennett, Professor E. M. Dunlop, Dr. A. B. Kerr, and my house-physician, Dr. Brownlie, for much generous assistance.

REFERENCES

- ¹ O'Donovan: *The Hair*, 1930, p. 20.
- ² Stelwagon and Gaskill: *Diseases of the Skin*, 1921, p. 1025.
- ³ Sabouraud: *Diagnostic et Traitement des Affections du Cuir Chevelu*, 1932, p. 13.
- ⁴ Joseph, Max: *Hautkrankheiten*, 1905, p. 237.
- ⁵ Darier: *Précis de Dermatologie*, 1923, p. 512.
- ⁶ Jordan and Kindred: *Embryology*, 1926, p. 495.

Memoranda

MEDICAL, SURGICAL, OBSTETRICAL

AN UNUSUAL CASE OF HYDRO-PNEUMOTHORAX

The following case presents features of unusual interest, and illustrates a group of conditions which would appear to be extremely rare.

History.—The patient, a married woman, aged 47, was admitted to Heathfield Sanatorium on September 2nd, 1932, with the history that six days previously she had consulted her doctor for a severe pain on the left side of the chest, which radiated to the left shoulder. Two days later she carried out her ordinary duties as charwoman in a cinema, but the same evening became violently sick, and vomited. The vomitus consisted of glairy fluid only, and anything she drank was returned immediately. Retching continued almost up to the time of admission to hospital, and was accompanied by pain of a pleuritic character on the left side. Two days after the appearance of these acute symptoms dullness was demonstrated at the left base, and the following day—the day preceding her removal to hospital—paracentesis was carried out and fluid drawn off. At 9 a.m. on the day of admission she had sudden acute pain in the chest, accompanied by severe dyspnoea. A diagnosis of pneumothorax was made, confirmed by an escape of air on the introduction of a needle into the chest, and the patient was sent into hospital with that diagnosis.

On admission she was extremely ill, presenting all the symptoms of acute heart failure and shock, was very dyspnoeic, and cyanosis was extreme, the hands being almost purple. Immediate examination was deemed inadvisable, treatment for shock being instituted at once. Physical examination without disturbing the patient revealed diffuse pulsation on the right side of the chest due to gross displacement of the heart, and a tympanitic note to percussion all down the left side of the thorax. "Coin" percussion failed to elicit the "bell" sound. The patient's condition gradually became worse, and she died a few hours after admission.

The following additional history was obtained from the patient's husband after her death. She had enjoyed good

health until the last few years of her life. She had had three normal full-time pregnancies, and two abdominal operations for prolapse of the uterus. Several years previously she became troubled with a slight cough, which was accompanied by a frothy spit, and for the past two years she had definitely lost weight. Frequently, over the same period, she would rise from the table, vomit, and then return to resume her meal; she also complained of flatulence, and occasional pain in the abdomen. Two or three months before she died she first complained of a pain, which recurred at frequent intervals up to the time of her death. This pain was located over the left side of the chest behind, radiating to the left shoulder, to the left side of the neck, and was sufficiently severe at times to cause her to lie down to obtain relief; sickness did not accompany this pain. An additional point of interest is that the patient remarked to her husband just prior to her removal to hospital "that she did not think anything was getting into the stomach."

Post-mortem Findings.—When the chest was opened the collapsed left lung was found to occupy the upper part of the thoracic cavity on the left side. The heart and mediastinum were markedly displaced to the right, and the remainder of the left thoracic cavity was occupied by the stomach, spleen, and some free fluid, which gave an acid reaction to litmus and was similar in appearance to the fluid found in the stomach. The stomach was slightly ballooned, and did not present the appearance of a strangulated viscus. When it was compressed more gastric contents found their way into the thoracic cavity from a perforated ulcer, situated on the posterior part of the stomach wall, near the gastro-oesophageal junction. The spleen was found lying behind the stomach close to the posterior margin of a hernial opening in the diaphragm, and was pale and shrunken, its surface having a crenated appearance; its diameters measured $3\frac{1}{4}$ by $2\frac{1}{2}$ inches. The hernial opening was ovoid in shape, occupied a position in the centre of the left leaflet of the diaphragmatic tendon, and measured $2\frac{1}{2}$ inches and $1\frac{1}{2}$ inches in its long and short axis respectively. Portions of the great omentum and gastro-splenic ligament adhered to its margin. At the hernial opening the pylorus came into contact with the cardiac end of the stomach, and by traction on the former the stomach could be partially returned to the abdominal cavity. The capacity of the stomach was found to be approximately 1,000 c.cm. To the left of the hernial opening and in close relation to the spleen, but from which it was separated by an invaginated portion of the diaphragm, lay the

OSWALD ADDISON: A PATIENT WITH FOUR URETERS

FIG. 1.—Pyelogram after intravenous injection of uroselectan. On right side two pelves and two ureters show clearly; the lower pelvis is dilated. On left side the two ureters can only be traced for part of their course.

FIG. 2.—Opaque catheters in position, showing four separate ureters.

OSWALD ADDISON: A PATIENT WITH FOUR URETERS

FIG. 3.—Catheters passed up median ureters, and 5 c.cm. sodium iodide injected in each side. The upper pelves only are filled, and do not show any communication with the lower pelves.

J. GOODWIN TOMKINSON: MONILETHRIX

FIG. 1.—Monilethrix, in son, aged 5½ years, sent for microscopical examination; suspected of having contracted tinea tonsurans. Keratosis pilaris is also seen.

FIG. 2.—Photomicrograph: monilethrix. Section of skin from scalp of father of boy aged 5½ years, showing three nodes within the hair follicle.