Published in final edited form as:

J Med Chem. 2007 April 19; 50(8): 1731–1734. doi:10.1021/jm070141c.

Novel Tricyclic Compounds Having Acetylene Groups at C-8a and Cyano Enones in Rings A and C: Highly Potent Anti-inflammatory and Cytoprotective Agents

Tadashi Honda^{*,†}, Chitra Sundararajan[†], Hidenori Yoshizawa^{†,§}, Xiaobo Su[†], Yukiko Honda[†], Karen T. Liby[‡], Michael B. Sporn[‡], and Gordon W. Gribble^{*,†}

Department of Chemistry, Dartmouth College, 6128 Burke Laboratory, Hanover, New Hampshire, 03755, and Department of Pharmacology and Toxicology, Dartmouth Medical School, Hanover, New Hampshire, 03755

Abstract

Novel C-8a functionalized tricyclic compounds having cyano enones in rings A and C have been synthesized and biologically evaluated. Amongst them, compounds with acetylene groups at C-8a show extremely high potency in in vitro and in vivo bioassays for anti-inflammatory and cytoprotective activities. Both in vitro and in vivo potencies are markedly higher than CDDO, which is being evaluated as an anti-cancer drug in phase I clinical trials.

Introduction

We have been engaged in the improvement of anti-inflammatory and anti-proliferative activity of oleanolic acid, a naturally occurring triterpenoid. This led to the discovery of CDDO and related compounds (e.g., CDDO-Me, CDDO-Im, CDDO-CN, and others).^{1–5} CDDO and its methyl ester (CDDO-Me) are presently being evaluated in phase I clinical trials for the treatment of leukemia and solid tumors.

During the course of these investigations, we have found that three-ringed compounds with similar enone functionalities in rings A and C to those of CDDO are also a novel class of potent anti-inflammatory, cytoprotective, growth suppressive, and pro-apoptotic compounds. Particularly, compound **1** is nearly equivalent in potency to CDDO for inhibition of nitric oxide

*To whom correspondence should be addressed. For T. H.: (phone) 603-646-1591; (fax) 603-646-3946; (e-mail) th9@dartmouth.edu. For G. W. G.: (phone) 603-646-3118; (fax) 603-646-3946; (e-mail) ggribble@dartmouth.edu.

[†]Dartmouth College

[‡]Dartmouth Medical School

[§]Visiting scholar from Shionogi & Co., Ltd.

(NO) production in primary mouse macrophages stimulated with interferon- γ , and is orally active in a preliminary in vivo inflammation model.^{6,7} These results encouraged us to synthesize additional new analogues. Thus, we initially designed C-8a functionalized analogues because insertion of functionalities at this position would be expected to improve the potency and pharmacokinetics due to the favorable balance shift between hydrophilicity and hydrophobicity. Also, insertion of polar and hydrophilic functionalities (e.g., carboxyl group, amine group, etc.) would give water-soluble compounds.

We have synthesized various C-8a functionalized analogues using new tricycles **2a–2c** (Scheme 1) as starting materials, whose efficient synthesis we have previously established for our projected synthesis,⁸ and evaluated their potency for inhibition of NO production in RAW 264.7 cells stimulated with interferon- γ (RAW cell assay). These analogues include typical electron-withdrawing, electron-releasing, hydrophilic, hydrophobic, and bulky groups.⁹ We have found that compounds **3–9** are more potent than compound **1** and CDDO in the RAW cell assay (Table 1). Particularly, compounds **3–5** and **7** with acetylene groups are the most potent. Their potency is higher than that of **1** and CDDO, and equal to that of CDDO-Im and CDDO-CN, which have the highest potency amongst semi-synthetic triterpenoids previously evaluated in the same assay.³ In addition, compound **4** is orally and highly active for induction of the anti-inflammatory and cytoprotective enzyme, heme oxygenase-1 (HO-1)¹⁰ in the liver and stomach (Figures 1 and 2). The in vivo potency is also much higher than that of **1** and CDDO.

In this communication, we report our initial results on the synthesis and biological potency of some new C-8a functionalized tricyclic compounds.

Chemistry

Compounds **3** and **4** in racemic form were synthesized by the sequence shown in Scheme 1. Compound **10** was obtained in quantitative yield by ketalization of **2c** with ethylene glycol (EG) in the presence of pyridinium *p*-toluenesulfonate (PPTS) in benzene, followed by Swern oxidation.¹¹ Compound **11** was prepared in 80% yield as a mixture of *E/Z* chlorovinyl isomers (*E:Z* = 4:1) by a Wittig reaction of **10** with (chloromethyl)triphenylphosphonium chloride.¹² Dehydrochlorination of **11** with MeLi, followed by trapping the acetylide with chlorotrimethylsilane (TMSCl) gave **12** in 93% yield.¹³ Deketalization of **12** under acidic conditions, followed by a chromium-mediated allylic oxidation,¹⁴ afforded **13** in 63% yield. Compound **3** was synthesized in 61% yield by double cyanation of **13** with LDA and *p*-toluenesulfonyl cyanide (*p*-TsCN),¹⁵ followed by DDQ oxidation (29% overall yield from **2c**). The TMS group was removed by tetra-*n*-butylammonium fluoride (TBAF)¹⁶ to give **4** in 71% yield (20% overall yield from **2c**).

Optically active (–)- and (+)-**3** and **4** were synthesized by the sequence shown in Scheme 2. The resolution of **2c** in racemic form was carried out in the manner described by Grieco.¹⁷ Treatment of **2c** with (–)-(*R,R*)-2,3-butanediol afforded the pair of diastereomers **14** and **15**. Separation of the two diastereomers was achieved by iterative flash column chromatography to give **14** (including 8% of **15**) and **15** (including 10% of **14**) in 29% and 26% yield, respectively. The diastereomeric purity was determined by ¹H NMR (300 MHz, CDCl₃) using the integration values of the methyl signals (γ 0.92 and 0.88 of **14**, and 0.96 and 0.86 of **15**) for the two diastereomers. Diastereomer **14** was then treated with acidic methanol to give (+)-**2c** in 91% yield. The deketalization conditions required much longer time (45 min) than deketalization of **12** (5 min) due to the bulky ketal of **14**. Similarly, **15** afforded (–)-**2c** in 88% yield under the same conditions. Based on the diastereomeric purity, we concluded that (+)-**2c** includes 8% of (–)-**2c** [enantiomeric excess (ee), 84%] and (–)-**2c** includes 10% of (+)-**2c** (ee, 80%). The circular dichroism (CD) values for the two enantiomers (+)-**2c** and

(-)-**2c** are $\Delta\epsilon_{288} = +0.22$ (c 0.0025, EtOH), and $\Delta\epsilon_{288} = -0.22$ (c 0.0025, EtOH), respectively. Based on these CD values and application of the octant rule,¹⁸ we have determined that (+)-**2c** has the same configuration as that of CDDO and (-)-**2c** has the opposite configuration.

We did not synthesize optically active **3** and **4** directly from **14** (or **15**) because the TMS group might be cleaved under the conditions for removal of the bulky ketal. Thus, we synthesized optically active **3** and **4** from (+)-**2c** (or (-)-**2c**) by the same sequence as for **3** and **4** in racemic form. Optically active (-)-**3** was obtained in 12% yield from (+)-**2c**. Removal of the TMS group was achieved by treatment of (-)-**3** with TBAF to give (-)-**4** in 48% yield. Similarly, (+)-**3** was synthesized in 15% yield from (-)-**2c**, and then removal of the TMS group of (+)-**3** afforded (+)-**4** in 37% yield.

Compounds **5–7** in racemic form were synthesized by the sequence shown in Scheme 3. Dehydrochlorination of **11** with MeLi, followed by quenching the acetylide with aqueous NH_4Cl solution afforded **16** in 95% yield.¹² Insertion of the methyl group into the acetylene moiety was achieved by treating **16** with MeLi and trapping the resulting anion with methyl iodide, to give **17** in 91% yield. The ketal **17** was subjected to acidic conditions to give **20** in 86% yield. Allylic oxidation of **20** afforded **23** in 59% yield. Double cyanation of **23**, followed by DDQ oxidation, gave **5** in 48% yield (21% overall yield from **11**). Ethylacetylene **18** was obtained in 69% yield by quenching lithium acetylide of **16** with iodoethane. Compound **6** was synthesized in 7% yield from **18** by the same methods as for **5**. Treatment of the acetylide of **16** with phenyl cyanate (PhOCN) afforded **19** in 64% yield.¹⁹ Compound **7** was obtained in 7% yield from **19** by the same procedures as for **5**.

Compound **8** in racemic form was synthesized by the sequence shown in Scheme 4. The aldehyde **10** was treated with MeMgBr in THF to give a diastereomeric mixture (2:1) of alcohol **26** in 87% yield. Ratcliffe oxidation²⁰ of **26** gave **27** in 73% yield. A forced Wolff-Kishner reduction of **27** with anhydrous hydrazine and KOH in diethylene glycol (at 165 °C for 2 h and, then at 217 °C for 24 h),²¹ followed by removal of the ketal, produced **28** in 51% yield. Compound **8** was synthesized in 48% yield from **28** by the same sequence as for **5** (16% overall yield from **10**).

Compound **9** in racemic form was synthesized by the sequence shown in Scheme 5. A Wittig reaction on **10** with methyltriphenylphosphonium iodide in the presence of potassium *t*-butoxide in THF gave **29** in 74% yield. Removal of the ketal of **29** afforded **30** in quantitative yield. Compound **9** was obtained in 48% yield via **31** from **30** by the same methods as for **5** (36% overall yield from **10**).

Biological Results and Discussion

We have evaluated the inhibitory activity of new C-8a functionalized tricyclic compounds **3–9** including optically active analogues in the RAW cell assay. The IC_{50} values are shown in Table 1. All new compounds **3–9** are more potent than the lead compound **1**, CDDO, and DXM. Noteworthy is that compounds **3–5** and **7** having acetylene groups are significantly more potent than **1** and CDDO. The potency is equal to CDDO-CN and CDDO-Im, which have the highest potency amongst semi-synthetic triterpenoids previously evaluated in the same assay.³

It is often the case that one enantiomer of a drug has greater potency and/or less toxicity than its antipode. Indeed, we have found that (-)- and (+)-**1** have different biological properties.⁷ In this assay, we have not found a significant difference between (-)- and (+)-enantiomers of **3** and **4**.

Presently, we are engaged in synthesizing various new tricyclic compounds having acetylene groups in order to develop clear structure–activity relationships for these compounds.

Subsequently to the in vitro assay, we have evaluated the potency of compound **4**, which is one of the most potent compounds in the RAW cell assay, for induction of the anti-inflammatory and cytoprotective enzyme, heme oxygenase-1 (HO-1) in the liver and stomach (in vivo, oral administration). There is major interest in stimulating HO-1 as a protective enzyme in many chronic disease conditions in which inflammation and oxidative stress play an important role.¹⁰ As shown in Figures 1 and 2, oral dosing of 1 μ mol of compound **4** causes significant induction of HO-1 in the liver and stomach, while compound **1** and CDDO are both markedly less potent at this low dose. Compound **4** is as potent as CDDO-Im in the liver but is clearly more potent than CDDO-Im in the stomach, which is the most potent semi-synthetic triterpenoid we have developed for induction of HO-1.²²

In conclusion, although a series of tricyclic compounds was designed based on CDDO, the structure of **4** is clearly different from that of CDDO, because **4** has an acetylene group at C-8a. Tricyclic compounds, diterpenoids, and triterpenoids with an acetylene group at C-8a (C-8 in terpenoid nomenclature) have not been reported prior to our synthesis of **4**. Currently, **4** is one of the most bioactive compounds in in vitro and in vivo bioassays in our pool of drug candidates including semi-synthetic triterpenoids and tricyclic compounds. Therefore, **4** may represent a new class of potential drug candidates having an entirely new structure for inflammatory diseases and cancer chemoprevention. Further syntheses and biological evaluation of new tricyclic compounds having acetylene groups at C-8a, including water-soluble analogues, are in progress.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We thank Charlotte Williams, Renee Risingsong, and Darlene Royce (Dartmouth Medical School) for expert technical assistance. This investigation was supported by funds from NIH grants R03-CA105294 and R01-CA78814, from the National Foundation for Cancer Research, and from Reata Pharmaceuticals. M. B. S. is Oscar M. Cohn Professor.

Abbreviations

CDDO	2-cyano-3,12-dioxooleana-1,9(11)-dien-28-oic acid
CDDO-Me	Methyl 2-cyano-3,12-dioxooleana-1,9(11)-dien-28-oate
CDDO-Im	1-[2-cyano-3,12-dioxooleana-1,9(11)-dien-28-oyl]imidazole
CDDO-CN	2-cyano-3,12-dioxooleana-1,9(11)-dien-28-onitrile
DXM	dexamethasone

References

1. Honda T, Gribble GW, Suh N, Finlay HJ, Rounds BV, Bore L, Favaloro FG Jr, Wang Y, Sporn MB. Novel synthetic oleanane and ursane triterpenoids with various enone functionalities in ring A as inhibitors of nitric oxide production in mouse macrophages. *J Med Chem* 2000;43:1866–1877. [PubMed: 10794703]

2. Honda T, Rounds BV, Bore L, Finlay HJ, Favaloro FG Jr, Suh N, Wang Y, Sporn MB, Gribble GW. Synthetic oleanane and ursane triterpenoids with modified rings A and C: A series of highly active inhibitors of nitric oxide production in mouse macrophages. *J Med Chem* 2000;43:4233–4246. [PubMed: 11063620]
3. Honda T, Honda Y, Favaloro FG Jr, Gribble GW, Suh N, Place AE, Rendi MH, Sporn MB. A novel dicyanotriterpenoid, 2-cyano-3,12-dioxooloana-1,9(11)-dien-28-onitrile, active at picomolar concentrations for inhibition of nitric oxide production. *Bioorg Med Chem Lett* 2002;12:1027–1030. [PubMed: 11909709]
4. Suh N, Wang Y, Honda T, Gribble GW, Dmitrovsky E, Hickey WF, Maue RA, Place AE, Porter DM, Spinella MJ, Williams CR, Wu G, Dannenberg AJ, Flanders KC, Letterio JJ, Mangelsdorf DJ, Nathan CF, Nguyen L, Porter WW, Ren RF, Roberts AB, Roche NS, Subbaramaiah K, Sporn MB. A novel synthetic oleanane triterpenoid, 2-cyano-3,12-dioxooleana-1,9-dien-28-oic acid, with potent differentiating, anti-proliferative, and anti-inflammatory activity. *Cancer Res* 1999;59:336–341. [PubMed: 9927043]
5. Place EA, Suh N, Williams CR, Risingsong R, Honda T, Honda Y, Gribble GW, Leesnitzer LM, Stimmel JB, Willson TM, Rosen E, Sporn MB. The novel synthetic triterpenoid, CDDO-imidazolidine, inhibits inflammatory response and tumor growth in vivo. *Clin Cancer Res* 2003;9:2798–2806. [PubMed: 12855660]
6. Favaloro FG Jr, Honda T, Honda Y, Gribble GW, Suh N, Risingsong R, Sporn MB. Design and synthesis of tricyclic compounds with enone functionalities in rings A and C: A novel class of highly active inhibitors of nitric oxide production in mouse macrophages. *J Med Chem* 2002;45:4801–4805. [PubMed: 12383005]
7. Honda T, Favaloro FG Jr, Janosik T, Honda Y, Suh N, Sporn MB, Gribble GW. Efficient synthesis of (–)- and (+)-tricyclic compounds with enone functionalities in rings A and C. A novel class of orally active anti-inflammatory and cancer chemopreventive agents. *Org Biomol Chem* 2003;1:4384–4391. [PubMed: 14685310]
8. Honda T, Honda Y, Yoshizawa H, Gribble GW. An efficient synthesis of tricyclic compounds, (±)-(4aβ,8aβ,10aα)-1,2,3,4,4a,6,7,8,8a,9,10,10a-dodecahydro-1,1,4a-trimethyl-2-oxophenanthrene-8a-carboxylic acid, its methyl ester, and (±)-(4aβ,8aβ,10aα)-3,4,4a,6,7,8,8a,9,10,10a-decahydro-8a-hydroxymethyl-1,1,4a-trimethylphenanthren-2(1*H*)-one. *Org Prep Proced Int* 2005;37:546–550.
9. These analogues will be published in a full account.
10. Ryter SW, Alam J, Choi AM. Heme oxygenase-1/carbon monoxide: from basic science to therapeutic applications. *Physiol Rev* 2006;86:583–650. [PubMed: 16601269]
11. Omura K, Swern D. Oxidation of alcohols by “activated” dimethylsulfoxide. A preparative, steric and mechanistic study. *Tetrahedron* 1978;34:1651–1660.
12. Mella M, Panza L, Ronchetti F, Toma L. 1,2-Dideoxy-3,4:5,7-bis-*O*-(1-methylethylidene)-D-*gluco*- and D-*galacto*-hept-1-ynitols: synthesis and conformational studies. *Tetrahedron* 1988;44:1673–1678.
13. Corey EJ, Ruden RA. Stereoselective methods for the synthesis of terminal *cis* and *trans* enyne units. *Tetrahedron Lett* 1973:1495–1499.
14. Muzart J. Synthesis of unsaturated carbonyl compounds via a chromium-mediated allylic oxidation by 70% tert. butylhydroperoxide. *Tetrahedron Lett* 1987;28:4665–4668.
15. Kahne D, Collum DB. Kinetic cyanations of ketone enolates. *Tetrahedron Lett* 1981;22:5011–5014.
16. Cai, Chengzhi; Vasella, Andrea. Oligosaccharide analog of polysaccharides. Part 3. A new protecting group for alkynes: orthogonally protected dialkynes. *Helv Chim Acta* 1995;78:732–757.
17. Grieco PA, Speake JD. Synthetic studies on quassinoids: Total synthesis and biological evaluation of (+)-des-D-chaparrinone. *J Org Chem* 1998;63:5929–5936. [PubMed: 11672197]
18. Charney, E. *The Molecular Basis of Optical Activity*. Wiley Interscience; New York: 1979. p. 167-191.
19. Murray RE, Zweifel G. Preparation of phenyl cyanate and its utilization for the synthesis of α,β-unsaturated nitriles. *Synthesis* 1980:150–151.
20. Ratcliffe R, Rodehorst R. Improved procedure for oxidations with the chromium trioxide-pyridine complex. *J Org Chem* 1970;35:4000–4002.

21. Barton DHR, Ives DAJ, Thomas BR. A Wolff-Kishner reduction procedure for sterically hindered carbonyl groups. *J Chem Soc* 1955:2056.
22. Liby K, Hock T, Yore MM, Suh N, Place AE, Risingsong R, Williams CR, Royce DB, Honda T, Honda Y, Gribble GW, Hill-Kapturczak N, Agarwal A, Sporn MB. The synthetic triterpenoids, CDDO and CDDO-Imidazolidine, are potent inducers of heme oxygenase-1 and Nrf2/ARE signaling. *Cancer Res* 2005;65:4789–4798. [PubMed: 15930299]

Figure 1. Tricyclic compound 4 is a potent inducer of HO-1 in liver when given by gavage
Male CD-1 mice (three mice per group) were separately gavaged with 1 μ mol of the following: tricyclic compounds **1** and **4**, CDDO, and CDDO-Im. After 6 h, livers were collected and analyzed by Western blot for HO-1.

Figure 2. Tricyclic compound 4 is a potent inducer of HO-1 in stomach when given by gavage
Male CD-1 mice (three mice per group) were separately gavaged with 1 μ mol of the following: tricyclic compounds **1** and **4**, CDDO, and CDDO-Im. After 6 h, stomachs were collected and analyzed by Western blot for HO-1.

Scheme 1. Synthesis of tricyclic compounds 3 and 4 in racemic form

Reagents and yields: (a) EG, PPTS, PhH, 100%; (b) DMSO, (COCl)₂, Et₃N, 100%; (c) Ph₃PCH₂Cl₂, *n*-BuLi, THF, HMPA, 80%; (d) MeLi, THF; TMSCl, 93%; (e) aqueous HCl, MeOH, 97%; (f) CrO₃, *t*-BuOOH, CH₂Cl₂, 65%; (g) *p*-TsCN, LDA, THF; (h) DDQ, PhH; 61% (i) TBAF, THF, 71%.

Scheme 2.
 Synthesis of optically active (-)- and (+)-**3** and **4**

Scheme 3. Synthesis of tricyclic compounds 5–7 in racemic form

Reagents: (a) MeLi, THF; aq. NH_4Cl ; (b) MeLi, THF; MeI; (c) MeLi, THF; EtI; (d) *n*-BuLi, THF, PhOCN; (e) aqueous HCl, MeOH; (f) CrO_3 , *t*-BuOOH, CH_2Cl_2 ; (g) *p*-TsCN, LDA, THF; (h) DDQ, PhH.

Scheme 4. Synthesis of tricyclic compound 8 in racemic form

Reagents and yields: (a) MeMgBr, THF, 87%; (b) CrO₃, pyr., CH₂Cl₂, 73%; (c) NH₂NH₂, KOH, diethylene glycol; (d) aq. HCl, MeOH; 51% from **27**; (e) CrO₃, *t*-BuOOH, CH₂Cl₂; (f) LDA, *p*-TsCN, THF; (g) DDQ, PhH; 48% from **28**.

Scheme 5. Synthesis of tricyclic compound 9 in acemic form

Reagents and yields: (a) Ph_3PMeI , $t\text{-BuOK}$, THF, 74%; (b) aqueous HCl, MeOH, 100%; (c) CrO_3 , $t\text{-BuOOH}$, CH_2Cl_2 , 83%; (d) LDA, $p\text{-TsCN}$, THF; (e) DDQ, PhH, 58%.

Table 1
Inhibition of NO production in RAW 264.7 cells stimulated with interferon- γ by new tricyclic compounds **3–9**^a

Compound			R	IC ₅₀ (nM)	IC ₅₀ (nM)
	(-)-enantiomer	(+)-enantiomer	Compound		
3^b	-C ≡ C-TMS		7^b	-C ≡ C-CN	4
(+)- 3	-C ≡ C-TMS		8^b	-Et	10
(-)- 3	-C ≡ C-TMS		9^b	-CH ≡ CH ₂	10
4^b	-C ≡ CH		1^b		30
(+)- 4	-C ≡ CH		CDDO		20
(-)- 4	-C ≡ CH		CDDO-Im		1
5^b	-C ≡ C-Me		CDDO-CN		1
6^b	-C ≡ C-Et		DXN		20

^a RAW 264.7 cells were treated with various concentrations of compounds and interferon- γ (10 ng/mL) for 24 h. Supernatants were analyzed for NO by the Griess reaction. IC₅₀ values are an average of two separate experiments.

^b A (1:1) mixture of (-)- and (+)-enantiomers (racemic form).