We consider it difficult to apportion the exact amount of credit due to this treatment and to the lead injections, though each may well have assisted the effects caused by the other.

Below this runs a graph of his weight, conforming to the scale on the right-hand side. Taking 12 st. as his normal weight when well, it will be seen that he had lost over 3 st. before treatment commenced; that he rose in weight steadily until he had regained his normal weight, and that he has remained at this for the last nine months.

The vertical lines rising from the base of the chart denote the weekly doses in numbers of cubic centimetres of the lead colloid in use at that time. Weekly intervals on these lines are dated by the months and years at the top of the chart. The numbers underneath the chart are used for reference to this article.

The injections given between (1) and (2) were of lead N. From (3) onwards he was having weekly doses of 10 c.cm. of lead M.A. pretty regularly, and he was putting on weight at a great pace. Part of this increase was due to the very considerable oedema of the feet and legs, as already mentioned, and at (5) he appeared to be suffering from early symptoms of lead poisoning. The injections were therefore discontinued for a short time. At (6) the lead M.A. was recommenced. From there till (7) doses were given intramuscularly; from (7) onwards the intravenous route was used. Owing to the fact that he appeared to be doing quite well as he was, doses were given less frequently from (8) to (9). He was having doses of α rays regularly at this time, but the growth showed signs of increasing in size rather than diminishing, and he was not feeling quite so well. At (9), therefore, full treatment was commenced again, and he at once became better in every respect.

On November 14th, 1924, he gave a *plus* two Wassermann reaction. On November 7th, 1924, Dr. Fletcher examined a portion of the growth, and reported as follows:

"This is undoubted squamous-celled carcinoma. The cancer cells are arranged in solid masses, freely infiltrating the sub-epithelial tissue. There are no cell nests."

This man has had up to the present 368 c.cm. of colloidal lead containing 0.36 gram, or $5\frac{1}{2}$ grains, of lead.

The condition of his blood has been extremely interesting. At the commencement of treatment the blood counts, taken several times, showed an average red count of 2,500,000 and a white count of 12,000. The blood on withdrawal looked like thin washy damson juice of a peculiar purplish colour. No normal red cells were to be seen, extreme poikilocytosis being present. His red cell count has increased to 3,600,000 and the white count is 10,000 odd. The differential white count is normal. The freshly shed blood is bright red, and the red cells are remarkably even and regular, though somewhat oval. It is noteworthy that, in spite of the large amount of lead which he has received, his red cells have increased in number and become normal in shape. No stippled cells have been found in his blood.

Full blood counts must be made in all cases, and stippled cells must be frequently sought for. They are quite common, together with polychromasia, in the blood of many cases, often after quite small doses of lead. We are accustomed to record them as the number of stippled cells counted over the number of fields examined, using a 1/12 oil-immersion and a No. 3 eyepiece. Urinalysis is essential when giving full doses.

Occasionally some protein shock has occurred a few hours after an injection—slight shivering, sweating, some rise of temperature, and more rarely some vomiting.

During the cold weather the preparation of lead M.A. is liable to form a gel, but in any case it should always be thoroughly warmed before being injected intravenously.

The injections have all been made with Record syringes of convenient capacity and No. 17 needles. The syringes can be boiled and used immediately while still hot.

No special technique is necessary, but in comparison with similar injections in the case of asthmatics and other patients the giving of an intravenous injection to a cancer patient is often a matter of extreme difficulty. The veins become much shrunken as the patient wastes, and there is a shortage of blood to distend them; they sometimes become hardened pipes of greater substance than an ordinary artery. The lack of supporting fat allows the vein to roll away from the needle as it attempts to pierce the vessel through the leathery skin.

In the series of cases here dealt with the dosage has on the whole been small, and we consider that there is a large margin of safety within which injections of lead can be increased in size and frequency before toxic symptoms from the lead need be expected. But the dosing of any particular case will depend upon a great many different

factors. Patients of large weight with no cachexia will call for large doses. Males will take larger doses than females. Emaciated patients and those with growths which have ulcerated to the surface will require small doses, at any rate to commence with. As any growth may abort, special care must be taken to make such abortion as aseptic as possible. In this connexion the blood supply to the part is probably an important factor. In growths of the gut abortion must be particularly guarded against.

Summary.

We are using a very stable colloidal suspension of lead and mercury (lead M.A.). This can be injected intravenously with the minimum danger of immediate shock or toxicity from the lead. It has undoubtedly had a beneficial effect upon the patient whose case is reported in full in this paper.

We now have a series of cases under treatment with much heavier doses, and the results obtained will be published at a later date.

Arrangements are now being made whereby a supply of lead M.A. will shortly be available for general use.

THE SURGICAL TREATMENT OF TIC-DOULOUREUX.*

BY

G. PERCIVAL MILLS, F.R.C.S.,

SENIOR ASSISTANT SURGEON, GENERAL HOSPITAL, BIRMINGHAM.

Of the various procedures which have been devised for the cure of this most distressing malady two only have stood the test of experience. They are: (1) Some form of alcohol injection. (2) Operation on the Gasserian ganglion by the Hartley-Krause method of approach.

Injection treatment has obvious advantages: it is a minor operation which involves no long convalescence, there is no risk to life, and serious complications are very rare. On the other hand, it is always more or less uncertain in its effects, and the relief from pain is often only temporary.

Open operation on the ganglion has one advantage only, but an important one—with the rarest exceptions it cures the patient, and cures him permanently. My own practice has been to try injection first, and, if this proves unsatisfactory, to operate. During the last five years I have performed fifty-two injections and thirteen operations. In most of the operation cases I had tried injection previously, but in a few cases the patients had been injected elsewhere and refused to have any more. Consequently it would appear that, out of every four cases, three have been reasonably comfortable with injection and one has needed open operation.

TECHNIQUE OF INJECTION.

There are two distinct routes of approach in attempting to inject the Gasserian ganglion—from the front and from the side. From the front it is rather difficult to find the foramen ovale, but, when found, the needle enters easily. It tends, however, at this angle to enter the subarachnoid space rather than the ganglion, and at least one serious disaster has followed injection by this route. The alcohol was evidently injected into this space, and paralysis of a number of important cranial nerves followed.

From the side it is easier to find the foramen ovale, but, on account of the obliquity of approach, the needle tends to stick in the orifice. In some skulls, even with the soft parts removed, it is impossible to pass a needle through the foramen. I believe, however, that alcohol injected into the mouth of the foramen infiltrates through into the ganglion, for after such injections anaesthesia has occurred in the first and second divisions as well as in the third.

I have confined myself to lateral injections, largely on account of the serious disaster already referred to as following an anterior injection. A local anaesthetic is employed so that the sensations of the patient may be

* Abbreviated from a paper read before the Midland Medical Society on November 25th, 1925.

used as a guide. This merely anaesthetizes the skin at the site of puncture, and the patient feels no pain till the point of the needle strikes the third division as it emerges from the foramen ovale. If at the same moment the needle is felt to enter a narrow cavity it is probably in correct position. The syringe is then attached and a few drops of alcohol injected to test the degree of resistance. If the piston slides down without effort the needle has passed to one or other side of the nerve, and the alcohol is not entering the ganglion. The needle is withdrawn a few millimetres and reinserted until a position is found in which there is a definite resistance felt on pushing down the piston. After a few drops have been injected sensation is tested by pricking the skin of the face with a fine needle. The development of numbness confirms the position, and the injection is then completed. I am satisfied if I get anaesthesia of the second and third divisions with numbness of the first, and many cases have been relieved of their pain in which I obtained numbness of the second and third divisions only. Of fifty-two injections I failed in eleven to get satisfactory lasting anaesthesia. This is roughly one case in five. In some of these I reinjected successfully and in the others I operated. I was guided in my choice by the difficulty experienced in injecting, and to some extent by the wishes of the patient.

Results.

Taking cases which were injected at least a year ago, twenty-five have been traced and eleven of these are still free from recurrence. The periods since the injection of these eleven are 12 months, 15 months, 17 months (2), 18 months, 2 years, 2 years and 3 months, 3 years (2), 3 years and 8 months, and 5 years respectively. Cases in which recurrence is known to have taken place are fourteen in number, and the average period of relief was one year and nine months. This hardly gives a fair estimate of the benefit conferred on the patient, for it has frequently happened that the recurrence of pain has been slight and temporary, so that further treatment has not been necessary. For example, a patient reported that she had been quite free from pain except for an attack a year ago, which lasted only a fortnight; this patient regards herself as cured, and is very grateful, but she appears in my statistics as having only one year's relief. It may fairly be said, therefore, of injections that four out of five are successful in giving relief for an average period of twenty-one months, and that there is quite a reasonable chance of the relief being permanent. How great this chance is can hardly be estimated from so small a number of cases, but a rough idea may be obtained by considering the figures quoted above. Of twenty-five cases traced, eleven have remained free, the average time since injection being twenty-eight months. As the cases which recur do so on an average in twenty-one months, it is not unreasonable to hope that most of these will be permanently cured. The only complications experienced in these cases treated by injection were that two patients complained of partial deafness, one in the same and one in the opposite ear. There were no cases of conjunctivitis.

OPERATION.

Operation was reserved for patients whom I failed to relieve by injection, or who had had repeated injections elsewhere and desired more radical treatment. The Hartley-Krause method of approach was used, and in the first few cases I contented myself with dividing the second and third divisions of the nerve inside the skull and injecting alcohol under direct vision into the ganglion. The recurrence of pain in a case treated on these lines by another surgeon led me to doubt the reliability of this procedure, and in subsequent cases two-thirds of the ganglion was removed. The method of dealing with the middle meningeal artery advocated by Jordan Lloyd was adopted, the artery being plugged in the foramen spinosum with a suitable peg (Lloyd used a match), and consequently only needing to be tied in one place instead of two. This makes its division much easier, so long as the peg does not come out.

Results.

I have performed this operation on thirteen occasions, and though some of the patients were over 70 years of age there has been no mortality. Not only was there no mortality, but in only one case was there any cause for anxiety. This was a case in which the peg came out of the foramen spinosum. I had found Jordan Lloyd's matches too soft and had tried an oak peg. This must have been too hard to fit snugly, for violent haemorrhage occurred on the following day, and in my absence from Birmingham my colleague Mr. Rose very kindly tied the external carotid, with an entirely satisfactory result. With this exception, all the cases had a good convalescence. There was a remarkable freedom from ophthalmic complications, and only one case of conjunctivitis has to be recorded. This, however, was severe, and, owing to the patient's refusal of treatment, resulted eventually in loss of vision in that eye. Other complications experienced were a seventh nerve paralysis, due, I think, to forcible retraction of the flap, and a temporary third nerve paralysis which I cannot explain.

All these cases were relieved of their pain with a single exception, but, as my first operation was only done two years ago, I can produce no evidence of permanent cure from my own experience. It is, however, well recognized that in all but the rarest cases removal of the ganglion cures the condition permanently. In fact, I have felt justified in telling my patients that, though the operation is severe, the cure is certain. This confidence was rudely shaken a few months ago by the last case I operated upon. The operation proved exceptionally easy, and I had such a perfect exposure of the ganglion and its roots that I was led to try the operation of division of the sensory root instead of resection of the ganglion. The patient had absolutely no relief whatever. As this was my first failure I naturally attributed it to the different operation I had performed, and said hard words to myself for trying to paint the lily. I explained the position, and suggested that I should make sure of relief by removing the ganglion. I then removed the lower two-thirds of the ganglion with a part of the mandibular and maxillary nerves, and had the removed tissue microscopically demonstrated the ganglion cells. There was not the slightest relief from pain! This was either a case in which the lesion was on the central side of the ganglion or one of purely subjective pain. The patient showed obvious signs of mental derangement, but these had naturally been attributed to the severity of her suffering.

Of the remaining eleven cases the pain has recurred in one. This was one of the earliest operations in which the ganglion was injected and not removed, and occurred in the patient who developed conjunctivitis. It has finally destroyed any hope I had that injection of alcohol under direct vision might prove a satisfactory alternative to excision.

It has been suggested that though the operation relieves the pain it turns the patient into a chronic invalid. Of two patients who recently wrote to me, one is working as a collier in a mine, and the other is the successful head mistress of a large school for girls.

ACTINOMYCOSIS SUCCESSFULLY TREATED BY IODINE IN MILK.

BY

HUBERT CHITTY, M.S.LOND., F.R.C.S.ENG.,
SURGEON, BRISTOL ROYAL INFIRMARY.

THE rapid and uninterrupted recovery of four consecutive cases afflicted by this distressing and usually fatal malady seems worthy of record.

The treatment adopted was evolved as the result of an observation made during the course of the following fatal case.

Late Treatment: Death.

A. S., a farm labourer, was admitted to the Bristol Royal Infirmary under my care in February, 1923, suffering from what appeared to be acute appendicitis. The appendix was removed. It was situated behind the caecum, and was bound down by dense adhesions. After an uneventful recovery the patient was