glands large enough to be easily palpable. We have not once seen a purulent discharge from the nostrils, but two cases, which both first received the serum on the third day of illness, did develop otorrhoea of mild type. There has not been a case of late nephritis.

I wish again to emphasize that this report deals only with cases which have come under treatment early in the disease.

As the result of the experience I have had with this treatment, I am very hopeful that we have now a remedy which gives promise of great usefulness, and if the estimate we have formed is found to be the general experience we may look for a reduced case mortality, a great reduction in the very severe complications and after-effects of scarlatina, and also, with the cutting down of the necessary quarantine time, we may hope for much relief to the strain which scarlet fever so often puts on the accommodation of isolation hospitals, and for a material reduction in the cost of such institutions. As a prophylactic this new method should prove of great service.

In the preparation of this serum there are at present great difficulties. The fact that the ordinary laboratory animals cannot be made use of to measure the antitoxic value of the serum is a great handicap. The need of the moment is a reliable and simple method of estimating the potency of any particular batch of serum. At present this can only be done in human beings, either by the effect on the intradermal reactions with toxin in healthy susceptible individuals or by the Schultz-Charlton reaction in early cases with scarlatina rash. Even by these methods it must be acknowledged that the estimate which can be formed is very crude. No doubt time will solve the difficulty, but until it is solved there is, I fear, danger that serum of low potency may find its way on to the market. We know that this has occurred with other antitoxic serums.

It would appear that the two essentials in securing maximum results with this method are—serum of high standard, and very early cases.

REFERENCE.
¹ *Lancet*, December 23rd, 1923.

SALIVARY CALCULUS.

BY

G. H. EDINGTON, D.Sc., C.M., F.R.F.P.S.G.,
 SURGEON, WESTERN INFIRMARY, GLASGOW.

THE occurrence of calculi in the salivary apparatus is well known, although infrequent as compared with the incidence of stone in the biliary and urinary systems. This comparative infrequency, hardly to be termed "rarity," is such that the presence of salivary calculi may be overlooked and the condition escape recognition when the patient is first seen.

In almost all textbooks of surgery mention, more or less detailed, is made of salivary calculi, and it is now possible to formulate a fairly typical syndrome. It happens, however, that patients sometimes do not spontaneously furnish a typical history, and the latter may or may not be elicited by interrogation. In such cases as the last the discovery of calculi is dependent on the surgeon being alive to the possibility of their presence, and keeping in his mind the salient clinical features of the condition.

Clinical Course.

The patient usually complains of pain in the floor of the mouth, associated with a swelling there which appears or increases during mealtimes, at which times the pain also is aggravated. Such a history is almost more than suggestive of obstruction in Wharton's duct, the common situation of a salivary calculus. Should, however, the calculus be in Steno's duct, pain will be experienced in the cheek and swelling may be observed there.

When the stone has existed for some time, and has reached a size sufficient to produce obstruction of the duct, enlargement of the salivary gland concerned may be expected. In the case of the submaxillary there may be a visible swelling below the jaw. Swelling of the gland indicates more or less

inflammation of that organ, and on palpation it is found to be of firm consistence. As with the duct, pain may be experienced in the swollen gland during mealtimes, and helps to differentiate the firm swelling from that of malignant disease of upper cervical lymphatic glands. Further help in the differentiation is obtained by bimanual palpation, which shows the external to be continuous with an intrabuccal swelling in the posterior part of the floor of the mouth.

The patient sometimes complains also of a disagreeable taste in the mouth, in which case a streak of pus will usually be seen exuding from the duct orifice, or further back in the floor of the mouth. The occurrence may or may not be associated with an acute inflammatory exacerbation. If marked inflammation exists the presence of calculus may be very successfully masked.

Inspection of the floor of the mouth in a suspected case may show swelling and redness of the orifice of Wharton's duct, and if a probe introduced into the duct strikes the stone the diagnosis is made certain. Sometimes one end of a lenticular-shaped calculus may protrude from the orifice, the inflamed or ulcerated margin of which grasps the stone. In the absence of such findings, palpation of the floor of the mouth may demonstrate a hard body in the course of the duct. In some cases mobility of the concretion within the dilated duct is a characteristic feature. In another type of case the calculus may be palpated far back and apparently lodged in the substance of the gland.

Spontaneous extrusion of the calculus may occur; more often removal has to be undertaken. This is accomplished by slitting the orifice of the duct, or incising directly over the concretion. In the latter case the duct is steadied by fixation forceps applied immediately posterior to the concretion. When the incision is made pus not infrequently wells out, carrying the stone with it. The calculus may not immediately escape through the incision, in which case I have found Lister's ear-hook of assistance. If the calculus be mobile it sometimes retreats into a loculus, may for long defy the surgeon's efforts, and be dislodged only with great difficulty. In cases where the above measures fail, and in cases where the stone is situated in the gland or in the commencement of the duct and cannot be extracted in the floor of the mouth, extirpation of the submaxillary gland from without is called for. While a comparatively easy operation as a rule, extirpation is sometimes rendered very difficult by inflammatory matting of the parts, and not only may the buccal cavity be opened into, but the lingual nerve may not escape injury.

Physical and Chemical Characters of Calculi.

Of yellowish-white colour, and sometimes of faintly greenish hue, salivary calculi may be spherical, torpedo-shaped, or cylindrical. Their surface may be smooth, but is often finely tuberculated. While usually the calculus is solitary, examples of multiple concretions are occasionally met with. Their composition is made up by mineral salts and animal matter. The salts concerned are variously given by different authors. According to Erichsen,¹ Waterhouse,² Ivor Back,³ and H. Bailey,⁴ they are principally calcium phosphate and carbonate; Butlin and Spencer⁵ say chiefly calcium phosphate with some carbonate; Syme⁶ mentions calcium phosphate alone, and Pearce Gould⁷ calcium phosphate and some ammonio-magnesium phosphate. Analysis was made in four of my cases (vide infra), and showed either phosphates alone or along with mere traces of carbonate.

Such salts as are found in the calculi occur also in the saliva, the composition of which is as follows: calcium phosphate and carbonate, sodium carbonate, potassium sulphate, and chlorides of sodium and potassium; carbonates are particularly abundant; mucin is abundant in submaxillary and sublingual, and absent in parotid, saliva (Howell,⁸ with whom Flack and Hill⁹ in the main agree).

Etiology.

The calculi are the result of the precipitation of the salts of the saliva. The exciting cause of the precipitation is not definitely known. Butlin and Spencer (loc. cit.) think that calculus formation is apparently connected with the presence of a plug of inspissated mucus. The same view is elaborated

by Back (loc. cit.), who considers that the genesis of the calculi is intimately connected with inflammation of the duct of the gland concerned, a plug of mucus containing bacteria being formed, and salts deposited thereon; such inflammation depends on septic conditions of the mouth. Be that as it may, the surgeon meets with many septic mouths without calculus formation; and, on the other hand, it is often found in cases of calculus that the mouth does not show any special foulness. It seems to me that Back's view may be taken as a working hypothesis, while the cause of the precipitation is not definitely known.

Situation.

It is not known whether precipitation occurs primarily in the duct or in the gland substance. Clinically, however, the concretions are usually found in the duct of the gland concerned, although in the case of the submaxillary gland it is not rare to meet with them apparently in the parenchyma. Their relative frequency in Wharton's duct may depend on the accessibility of that structure to palpation.

All writers are agreed that calculus is much more frequently found in the submaxillary duct than in those of the other salivary glands. Andrews¹⁰ gives the frequency as about five times greater than in any other gland. Bailey gives the ratio as a little more than 50 to 1; in my cases it is 8 to 1. This probably depends on the relative abundance of mucin in submaxillary saliva (Howell, supra). Bailey (loc. cit.) quotes Christopherson's observation that in the Sudan parotid are much commoner than submaxillary calculi.

CASES.

The following nine cases, seen during the past twenty-five years, illustrate and are grouped according to the signs and symptoms furnished by individual patients. The submaxillary was implicated eight times, the parotid once.

A. SUBMAXILLARY.

(1) "Mealtime" Syndrome.

CASE I.¹¹—Female, aged 30, seen 1906. Submaxillary swelling, right, for six weeks; three weeks later dental extraction without improvement; swelling painful at mealtimes; disagreeable taste in mouth; gland stony hard, and pressure on it caused pus and mucus to exude from duct orifice. Attempted probing failed; no calculus felt on palpating duct; incision in floor of mouth failed to identify duct. Gland excised through external incision, and found to contain calculus, size of small pea, in dilated commencement of duct.

CASE II.—Male, aged 24, seen September, 1922. Discomfort in right submaxillary region, for nine years, especially during meals; twelve months ago had abscess below tongue. Prominent mucosa at orifice of Wharton's duct, and in line of duct. Palpation revealed hard body in duct; mucosa fixed by forceps, and incised, evacuating stone measuring 7 by 10 mm., with smooth hard coat; distal end bluntly tapered, proximal end thicker and bare; subsequent probing negative.

CASE III.—Male, aged 29, seen May, 1925. Pain and swelling in left side floor of mouth, and in neck, worse at mealtimes; swallowing hindered. Stony hard swelling in left submaxillary gland; sublingual thickening, juicy, in floor of mouth; no calculus felt; no glossitis. Calculus passed spontaneously, and discomfort passed away. Calculus, spherical, measured 7 by 5 by 4 mm. Analysis (Dr. Cappell) showed composition to be almost entirely of calcium phosphate, with traces of carbonates and organic matter; no sulphate present, and only a trace of magnesium and iron.

(2) Swelling in Floor of Mouth.

CASE IV.¹²—Male, aged 16, seen November, 1900. Swelling "under tongue" for ten days. Redness and swelling of orifice of right Wharton's duct; in centre of reddened area, small yellowish-white point, calcareous when struck by probe; felt very hard between finger and thumb. Mucosa proximal to swelling fixed by forceps, and incision freed stone, which was then squeezed out by finger and thumb, and followed by stringy mucus; no distension of duct, or swelling of gland. Concretion flattened ovoid, 6 by 4 by 3 mm., nucleus visible through cracks in rind. Analysis (Dr. W. B. Brodie) showed composition to be entirely calcium phosphate, with some albuminoid matter (mucus); carbonates, chlorides, and sulphates absent.

(3) Swelling in Floor of Mouth, and Submaxillary Swelling.

CASE V.—Male, aged 38, seen June, 1925. Swelling under tongue and in right submaxillary region, for one week. Swelling and signs of inflammation in line of right Wharton's duct; probe struck calculus. Incision evacuated pus in which escaped a small cylindrical calculus measuring 9 by 4 mm. Analysis (Dr. Cappell) gave precisely similar results as in Case iii. An element of confusion in

this case when first seen by his medical attendant was the presence of an enlarged gland of two years' duration at angle of left mandible. This gland was clearly tuberculous. There was a history of enlarged glands in childhood, and scarring was present in the left anterior triangle, and over the sterno-mastoid.

(4) Submaxillary Swelling, Acutely Inflamed.

CASE VI.—Male, aged 55, seen May, 1924. Submaxillary swelling, left side, of some years' standing; rapid enlargement of one week's duration, with pain on swallowing and on protruding tongue, and carache on same side. Firmness of swelling suggested malignancy; but no lesion could be seen or felt in tongue or fauces. No evidence of calculus. Increase of swelling and onset of oedema in floor of mouth posteriorly led to external incision, evacuating flakes of pus. During convalescence the mouth became very foul. Three weeks after incision pus observed far back in floor of mouth. Later, swelling in floor of mouth subsided with discharge of pus, and reaccumulated. Pus came from opening in inflamed area far back in floor, without induration of cancer. Later (February, 1925) palpation detected stony hardness anteriorly to fauces; bimanual palpation showed continuity with enlarged submaxillary gland externally, the whole mass moving freely on mandible. Extirpation of submaxillary gland was carried out with difficulty, owing to matting, and the floor of the mouth was opened and the lingual nerve damaged during removal. Healing took place after slow convalescence, and with slight "anchoring" of tongue to floor of mouth and anaesthesia. Examination of gland after removal showed embedded in intrabuccal part a large ovoid calculus, 14 by 20 mm., with tuberculated surface. Similarly tuberculated calcareous material was present on wall of containing cavity. Histological examination of gland (made by Dr. Cappell) showed involution and atrophy of secreting tissue, chronic inflammatory changes, and much fibrous tissue.

(5) Concretion Presenting at Orifice of Duct.

CASE VII.—Male, aged 31, seen September, 1922, with tip of salivary calculus projecting from orifice of left Wharton's duct. Inflammation of fold in floor of mouth, and enlargement of submaxillary gland. Portion of concretion, measuring 12 by 7 mm., extracted; remainder felt by probe in duct, and extracted the following day by Lister's ear-hook. It measured 10 by 7 mm., was cylindrical in form, and with cup-like extremity towards duct orifice. Probed duct subsequently to 1½ in., with negative result.

(6) History of Concretions being passed from Duct.

CASE VIII.—Male, aged 22, seen December, 1916. Had history of "enlarged gland at left angle of mandible, opening into mouth by a sinus, sometimes emitting a concretion." There had been long-continued discharge from left Wharton's duct, and periodic swelling of submaxillary gland; no calculus palpable. Submaxillary and portions of sublingual gland excised; firmly adherent to tissues of neck.

B. PAROTID.

CASE IX.—Male, aged 20, seen November, 1916. For three years had had swelling of left cheek, varying in size. On palpation, calculus felt at commencement of Steno's duct (posteriorly). Removed through horizontal incision in skin of cheek over stone; not ascertained if in duct or gland tissue. No ulceration or inflammation at orifice of duct. No sutures; wound healed on ninth day. Swelling of gland persisted, especially in anterior part below level of duct. Duct probed with negative result. Analysis (Captain W. Campbell, R.A.M.C.) showed inner core of cretaceous matter, outer coating of inspissated mucus. Ovoid shape, measured 7.5 by 6.5 mm. Composition: calcium, magnesium, and phosphorus pentoxide.

REMARKS.

The above series is far too small for statistical purposes. It illustrates, however, the comparatively high incidence of calculus in the submaxillary apparatus; the high proportion of males; the absence of constant clinical picture; the obscuration of diagnosis by inflammation (Case vi); and in the four specimens analysed the part taken by phosphates.

It is interesting to note, also, that Cases ii and vii were brothers, and that the solitary parotid case (ix) occurred in a soldier in Egypt, although the trouble had probably started before he went there.

REFERENCES.

- ¹ Erichsen: *Science and Art of Surgery*, tenth edition, London, 1895, vol. ii, p. 702.
- ² Treves's *System*, London, 1896, vol. ii, p. 491.
- ³ Choyce and Beattie's *System*, London, 1912, vol. ii, p. 243.
- ⁴ Bailey: *Brit. Journ. of Surg.*, 1924, vol. xii, p. 321.
- ⁵ Butlin and Spencer: *Diseases of the Tongue*, London, 1900, p. 236.
- ⁶ Syme: *Principles of Surgery*, second edition, Edinburgh, 1837, p. 427.
- ⁷ Pearce Gould: *Internat. Textbook of Surgery*, London, 1900, vol. ii, p. 165.
- ⁸ Howell: *Textbook of Physiology*, third edition, London, 1909, p. 727.
- ⁹ Flack and Hill: *Textbook of Physiology*, London, 1919, pp. 371-2.
- ¹⁰ Keen's *Surgery*, London, 1908, vol. iii, p. 323.
- ¹¹ *British Dental Journal*, London, December, 1906.
- ¹² *Glasgow Medical Journal*, vol. iv, 1901, p. 118.