Purvis Oration

ON

THE SURGERY OF MALIGNANT DISEASE
OF THE PHARYNX.DELIVERED BEFORE THE WEST KENT MEDICO-CHIRURGICAL
SOCIETY ON DECEMBER 11TH, 1925,

BY

WILFRED TROTTER, M.S.LOND., F.R.C.S.ENG.,
SURGEON, UNIVERSITY COLLEGE HOSPITAL.

(With Coloured Plate.)

WHATEVER may be our hopes about the discovery in the treatment of malignant disease of methods more specific and less harsh, surgical operation must be admitted to be the most useful and trustworthy means we now possess. The extension of the field to which it can be applied and the increase in the refinement with which it can be used are therefore among the most important duties of the surgeon. The development of the operative treatment of malignant disease in whatever department is not, however, a matter only concerned with surgical technique, but is equally dependent on pathological and clinical principles, and therefore has a general as well as a more special interest. It is for this reason that I have ventured to choose for my subject a branch of surgery which may at first sight seem to promise little but narrow technicalities, for I hope to show that it has some relations that concern us all in whatever branch of our art we may practise.

At this point I may perhaps be allowed to say a few words about my personal connexion with the subject, and in justification of the hope that my experience in it may have included some aspects that are not of common knowledge. In some Hunterian Lectures given in 1913 I laid down certain principles which had already proved useful in the treatment of malignant growths of the pharynx and tried to define the chief clinical generalizations that can be made about the disease in question. In the ensuing twelve years one's experience naturally has been largely added to. The steady increase of experience, while it has not perhaps made any changes in practice that amount to alterations of principle, has in this considerable lapse of years allowed of such an accumulation of changes in the technical detail of treatment that the lectures of 1913 no longer represent what is, in my judgement, the best we can do for the unfortunate sufferers from carcinoma of the pharynx. As my treatment of the subject on this second occasion of dealing with it is to be in the nature of a revision rather than a primary exposition, I shall hope to be able to keep in sight the aspects of more general interest that it possesses and to pass over its minor technicalities.

There are two further limitations I shall venture to set upon my handling of the material I have to put before you. First, I shall deal only with malignant disease of that part of the pharynx which is called by anatomists the laryngeal part. This includes the lower half of the length of the tube, and excludes matters like growths of the tonsil, which are more or less familiar to all, and growths of the naso-pharynx, which constitute a group having very different pathological relations. Limiting ourselves to the laryngeal pharynx we shall be concerned with two groups of cases—the growths about the upper opening of the larynx (epilaryngeal group), and the growths of the post-cricoid region (hypopharyngeal group). In both these groups we are concerned only with the squamous epithelioma. The second limitation I shall venture to set will be in the avoidance of statistical statements. Where experience is necessarily small—and the total number of the cases in question that I have ever seen, much less operated on and followed up, must probably be limited to a few hundreds—statements in figures are apt to possess more the air of authority than the substance of it, and to put more responsibility on the person who observes and classifies the facts than he may be competent or willing to

assume. These considerations are even stronger when the subject under inquiry is a changing and growing one, and as such I have always regarded the surgery of the pharynx. I shall therefore avoid numerical evidence in supposed confirmation of the conclusions I come to, and shall allow these to depend on a judgement that in the circumstances cannot perhaps be regarded as erring characteristically in the direction of haste.

THE GENERAL CHARACTERS OF MALIGNANT DISEASE
OF THE PHARYNX.

It has commonly happened in the evolution of medicine that when a certain part of the body has come within surgical access, this mere fact has quickly reacted on the pathology and symptomatology of the part in an unexpectedly fertilizing way. This sequence, which we are all familiar with on the great scale in the cases of the abdomen, the chest, and the brain, has shown itself with precision in the minor example of the pharynx. As soon as it became evident that in the operation of lateral (trans-thyroid) pharyngotomy we have a means that gives easy and adequate access to the laryngo-pharynx and allows satisfactory treatment of a large number of the growths found there, it began also to be evident that our clinical and pathological knowledge of the early phases of malignant disease in this situation could and must be largely extended and clarified. We found that there are characteristic starting-points for the disease, that there are variations in type and progress, and that all these facts have a direct and practical bearing on operability and on post-operative prognosis. It is to the conclusions derived from such observations that I am particularly anxious to call your attention.

(a) *Frequency*.—Malignant disease of the laryngo-pharynx is not at all rare. In my opinion the disease is not very much less common here than it is in the mouth or oesophagus, so that our subject must take its place among the great primary groups of malignant disease the incidence of which is favoured by some local condition of normal life. Any attempt to make an exact statement of its frequency is quite useless, because even yet relatively few cases of growths in the cervical alimentary tract are submitted to exact diagnosis, and because the clinical record of cases is confused by the description of an indefinite number as "extrinsic carcinoma of the larynx." Whatever the exact figure may be, there can be no doubt at all that the disease is common enough frequently to escape diagnosis for many months. Its position in relation to diagnosis is like that of rectal carcinoma; it is common, its symptoms may well be those of much commoner trivial diseases, and to exclude it involves an examination that if rarely difficult is often a little troublesome. Some such comfortable diagnosis as that of "relaxed throat" is therefore apt to tempt our less conscientious moments.

(b) *Rate of Progress*.—Epithelioma in this region does not usually progress with great rapidity, and there is generally after the first appearance of definitely suspicious symptoms a period of several months during which the case is still operable. There is one very important factor which must always be allowed for in determining the apparent extent and progress of these growths, and that is the influence of sepsis. Sepsis in the pharynx and oesophagus is determined by the teeth, and, it must be added, not so much by their condition as by their presence. It is by the comparison of the appearances of epitheliomata occurring in those with teeth with the appearances of such growths occurring in the edentulous that one is able to estimate the influence of sepsis in exaggerating the apparent extent and gravity of the lesion. The two most important facts that experience has established in this regard are, first, that a growth that seems hopeless may become obviously operable within a few weeks of the clearance of the mouth, and secondly, that in patients already edentulous epitheliomata are apt to run an insidious and relatively symptomless course for several months.

(c) *Incidence according to Sex and Age*.—Epithelioma of the laryngo-pharynx shows perhaps the most remarkable example known in the pathology of malignant disease of the primary seat of the tumour being determined by the

sex of the patient. It is a familiar fact that the post-ericoid or hypopharyngeal carcinoma is practically limited to women, while growths around the upper laryngeal opening (epilaryngeal carcinoma) on the one hand, and of the cervical oesophagus on the other, although not wholly limited to men, show a strong preference for that sex. In addition to being almost exclusively a disease of women, the post-ericoid carcinoma is peculiar in appearing at a relatively early age. The age at which it appears may be estimated as at least ten years younger than that at which the epilaryngeal carcinomata of men begin to be common. My experience suggests that carcinoma beginning definitely on the posterior pharyngeal wall belongs to the post-ericoid group as regards sex and age incidence. This is a comparatively rare type; it arises on the posterior wall well above the cricoid region, and often is directly visible through the mouth. It is interesting from its incomprehensible association in sex incidence with the post-ericoid growth, and it has a certain technical interest in demanding for its treatment an operation of special design.

Malignant disease of the laryngo-pharynx, then, although including an indefinite number of minor variations, falls naturally into two broadly defined main groups that we name for convenience epilaryngeal and hypopharyngeal. The differences as to sex incidence between these two go with clinical and therapeutic peculiarities, so that the natural classification is also one of practical value (Figs. 1 and 2).

GROWTHS OF THE EPI-LARYNGEAL TYPE.

These growths occur on and immediately around the upper opening of the larynx. The great majority of them occur in the male sex, and of cases of malignant disease of the laryngo-pharynx in that sex the great majority are of this type. They occur in men who have reached or passed middle age. Four common starting-points can be made out by the study of early cases; these are: (1) the epiglottis; (2) the aryepiglottic fold, especially toward its arytenoid extremity; (3) the lateral pharyngeal wall outside and away from the arytenoid; and (4) the pyriform sinus. When these starting-points are known a little experience will enable one from the laryngoscopic examination to estimate with some precision with which type one has to deal in a given case. The decision is important and worth an effort to attain, as it has a direct bearing on treatment and prognosis.

It may be stated at once that growths starting at any of the three points first mentioned (the epiglottis, the fold, the lateral wall) lend themselves to satisfactory operative treatment. At early and even moderately advanced stages a cure or freedom from the disease for many years can be looked for with reasonable confidence, without great operative dangers, and with no permanent mutilation or disability. The case is different with growths that start in the pyriform sinus. In this case the situation of the growth makes it symptomless in the early stages, and also leads to early involvement of the lateral wall of the larynx on the one hand, and of the thyroid cartilage on the other. It often happens, therefore, that by the time diagnosis is possible even an extensive and mutilating operation is of little use (Fig. 3).

Diagnosis.

I do not propose to discuss the finer shades of diagnosis among these types. The uses of practical work will be met if we know, first, when to suspect malignant disease of

the laryngo-pharynx; secondly, how to make a *reasonably probable diagnosis* of its presence; and thirdly, how to estimate in a general way the *gravity of the case* and the prospects of treatment.

In middle-aged and elderly men any kind of abnormal sensation persistently felt in the same part of the throat should be regarded seriously. The sensation may be a tickling; it may be that of a crumb or larger body of food lodged in the throat; it may be a discomfort in swallowing saliva alone while there is no difficulty at meals. There is usually no pain in these early stages, no trouble in swallowing food, no alteration of the voice, except perhaps to a very finely discriminating ear, and no affection of the general health. Occasionally enlarged glands—not necessarily carcinomatous—appear very early. For this reason palpation of the cervical glands is an art that should be cultivated very carefully, and one should not be content to regard the glands as normal in a given case unless the patient has been examined lying down and with all the muscles of the neck relaxed.

Laryngoscopic examination is, however, of course the essential method. It must be admitted that malignant disease of the pharynx is overlooked rather often in its early stages. I am convinced that this is due to no difficulty in the diagnostic problem itself, but rather to the modesty of medical men in tending to regard the laryngoscope as an instrument always and only for the expert. It is perfectly true that to get a good view of the interior of the larynx, and especially of the anterior part of the vocal cords, is sometimes very difficult. There is, however, no need for this in the cases we are discussing. All that is necessary is to get an adequate view of the upper opening of the larynx and to estimate the mobility of the arytenoids. This is easy, and in the majority of cases demands no more than that the mirror should be used with a good light, some gentleness and patience, and a general idea of what one may expect to see.

There are three appearances to be on the look-out for when a growth of the epilaryngeal region is suspected: (a) an ulcer with a raised edge, which may, of course, be seen only in part, but of which the merest glimpse definitely caught is diagnostic; (b) a collection of muco-pus which the patient cannot get rid of by swallowing; (c) a fixed arytenoid. A fixed arytenoid, if much ulcerating growth can be seen, generally indicates a late stage of disease; if little ulceration can be seen it points to the dreaded pyriform sinus type. A freely movable arytenoid, even when the growth seems of formidable size, is always a favourable sign. (Plate, Fig. 1.) The amount of fungation of a growth is not necessarily a measure of its gravity; sometimes the pharynx seems filled by an ugly sloughy mass which at operation proves to be semi-pedunculated and arising from a relatively localized epithelioma.

GROWTHS OF THE HYPOPHARYNGEAL TYPE.

These growths occur in the narrow tubular part of the pharynx behind the cricoid cartilage. They are practically limited to women, and include the great majority of epitheliomata of the pharynx in that sex. They appear in the adult and middle-aged, and are not uncommon in the fourth decade. A remarkable feature of my experience of these cases is that the history given by the patient often shows that there has been some difficulty in swallowing for many years, and that on this slight disability the serious dysphagia has gradually supervened. Difficulty and awkwardness in swallowing are common complaints in

FIG. 1.—Distribution and sex incidence of epithelioma of laryngo-pharynx. The epilaryngeal group, the ordinary type occurring in men.

FIG. 2.—Distribution and sex incidence of epithelioma of laryngo-pharynx. The hypopharyngeal group, practically limited to women.

FIG. 3.—Diagram of section through pyriform sinus of pharynx to show its relation externally to the thyroid cartilage and internally to the lateral wall of the larynx.

women, and are doubtless often an expression of "nervousness." It is perhaps for this reason that in a good many cases of post-cricoid carcinoma the early stages of definite dysphagia are not always taken very seriously. This is especially to be regretted, as the condition is a very formidable one, so that satisfactory treatment is more than usually dependent on early diagnosis. Symptoms that must always be regarded with very grave suspicion are *definite obstructive dysphagia even if regurgitation is only occasional and slight, frequent choking fits during meals, and chronic laryngitis with huskiness* from leakage of food into the larynx. Signs to be made out by examining the neck are all late and of little value; they are forward displacement of the larynx, palpability of the primary growth, and enlarged glands.

Laryngoscopic examination, if we know what to look for, is of great value in spite of the fact that by it a definite diagnosis cannot always be made. Three stages in the laryngoscopic appearances can usually be made out; examination, of course, is directed to the retroarytenoid region, and is quite easy. (a) In the earliest stages there is no growth visible and no swelling, but there is more or less chronic laryngitis, and in the pharynx behind the arytenoids is visible a pool of frothy mucus or muco-pus which the patient cannot clear away by swallowing. (Plate, Fig. 2.) (b) Later there is added to this picture a purplish, glistening swelling of oedema over the arytenoids and in the inter-arytenoid region. (Plate, Fig. 3.) (c) Finally, on the anterior wall of the pharynx below the top of the arytenoids there appears the raised edge of an epitheliomatous ulcer. *Only quite a small segment of the ulcer may be visible, and it may emerge from the pool of mucus only momentarily and at the height of a strong effort of phonation.* (Plate, Fig. 4.) Once such an appearance has been made out clearly the diagnosis is settled. The fixation of an arytenoid is common, but is usually a late phenomenon.

In the presence of strong reason to suspect post-cricoid carcinoma a characteristic laryngoscopic picture should not be waited for, but a direct examination should be made with the bronchoscope under an anaesthetic. On no account, however, should the surgeon be tempted to decide the vertical extent of the growth by forcing the tube past it. In my experience the operability of a given case of post-cricoid carcinoma can be decided definitely only by an actual exploration, though a negative decision can often be made by clinical examination in advanced cases. When an operation is begun in doubt as to whether it can be carried through, the operability of the case is to be judged without the pharynx being opened. In these circumstances retreat is possible without any serious damage having been inflicted.

TREATMENT.

There is one method of general utility in the operative treatment of carcinoma of the laryngo-pharynx. This is the operation of lateral (trans-thyroid) pharyngotomy, which is the foundation on which all the different procedures, from the simplest to the most elaborate, must be built up. It is primarily a method of access, and depends on the anatomical fact that the laryngo-pharynx lies within the laryngeal skeleton. If the great cornu of the hyoid bone and the ala of the thyroid cartilage are removed, the wall of this part of the pharynx is disclosed unopened, and through it the situation and extent of a growth can be determined by palpation with some precision. It can now be decided if a radical procedure is possible, and what it should be (Fig. 4). These are the essential facts, and I do not propose to enter here into technical details. We must consider, however, the further general lines on which we may be called upon to proceed.

The cases I have had to deal with in an experience of nearly twenty years' work in this field have fallen into three broadly distinguishable classes.

A. Inoperable Cases.—This class, always and still deplorably large, comprises probably a good half of all the cases I have had to do with. Most are recognizable for what they are at a glance; some have been, and a few still are, after every effort of foresight, submitted to fruitless exploratory operations. Here we have this satisfaction that the preliminaries that are necessary in such formidable cases are the best palliative measures that can be devised, and in themselves give great relief and some prolongation of life. These preliminaries are tracheotomy (in such circumstances under local anaesthesia) and a complete clearance of the mouth; they are measures always worth considering, even when there can be no thought of radical operation. There can be no doubt that a large proportion of the inoperable cases I have seen were at some part of their course favourable in type and within the reach of cure.

B. Operable Cases of the More Serious Types.—These are cases in which a more or less formidable operation offers a reasonable prospect of freedom from the disease for a period that may extend to several or many years. In order,

however, that the life thus saved or prolonged shall be tolerable, it is necessary to deal with the mutilation the extensive operation has entailed by more or less complicated plastic procedures to restore the functions of respiration, speech, and deglutition. (Plate, Fig. 5.) It is impossible to describe here in detail any of the plastic measures that may be necessary, but we may indicate that some such procedure must invariably be foreseen in operations for post-cricoid growths, for growths on the posterior pharyngeal wall, and for growths of the epilaryngeal group when any large resection of the party wall between pharynx and larynx has been made. These cases, therefore, demand from the surgeon a certain amount of special experience, a readiness on his part to evolve the design and the carrying out of the operation together, and to improvise necessary modifications during its actual course. Of the cases that come to professedly radical operation probably about half belong to this class. They involve difficult and dangerous procedures, and have an apparently endless capacity for disappointing at any moment the most plausible hopes. Nevertheless, and perhaps especially for this reason, the small proportion of really complete success they provide is quite enough to encourage the effort to increase it. A special difficulty of this

FIG. 4.—To illustrate the principle of the operation of lateral trans-thyroid pharyngotomy. The outline of the laryngeal skeleton is shown. The finely dotted lines indicate the relative positions of the aryepiglottic fold, the arytenoid cartilage, and the vocal cord. The thickly dotted lines show where the great cornu and the thyroid ala are divided in the operation. When these parts have been removed the laryngo-pharynx is accessible.

group is that, as many of the cases are advanced, extensive gland dissections often must be done to obtain access to the primary growth. Such procedures are, in my experience, best combined in one operation, and it is often possible, by making use of the sterno-mastoid muscle as a partition in the wound, to preserve the raw area left by the gland dissection from infection from the pharynx and to secure primary union in it.

C. Operable Cases of the More Favourable Types.—This class includes as its most important members those growths of the epilaryngeal group that I have already mentioned as the most favourable—namely, those of the epiglottis, aryepiglottic fold, and lateral wall. These may amount to about half of all the operable cases. The epithelioma that occurs in the three sites mentioned is not of a very malignant form; even in the fairly advanced stages in which it is still so often first seen by the surgeon it is frequently not beyond cure; the situation is anatomically such that large excisions of the pharyngeal wall are possible without the need for plastic measures; the tendency to gland infection is not very strong, and is inclined to be limited to one side of the neck. For these reasons cases of this class can be approached by the surgeon with every reasonable confidence that decidedly satisfactory results will be obtained. There is probably none of the major types of malignant disease in general in which surgical treatment—with some allowance for the special local difficulties—is on the whole more satisfactory than it is in this far from uncommon group of epilaryngeal growths. This favourable estimate implies, of course, not

merely that there is a reasonable expectation of cure or prolonged freedom from the disease in all fairly early cases, but also that such results are to be obtained without the infliction of any serious mutilation or even disability. The tumour is to be dealt with by the operation of lateral pharyngotomy in its simplest form, and as in these circumstances it is reduced to a straightforward and almost formal procedure I shall venture to mention the chief points in the routine to the use of which, after innumerable variations, I have more or less definitively settled.

Lateral Pharyngotomy in the Early Epilaryngeal Case.

Preliminary Preparation.—As one confidently expects to be able to close the pharynx so that leakage into the neck cannot occur—if at all—for several days, I do not now insist on a preliminary clearance of the mouth in these cases. Thus the patient is saved a good deal of extra distress and about three weeks' delay.

Anaesthetic.—Chloroform first through the mouth, later through the tracheotomy tube, is always used. To keep down the total amount used—a most important point—cocaine solution (10 per cent.) is injected into the trachea before this is opened and applied to the interior of the pharynx in plugs as soon as the lumen is exposed. Small hypodermic injections of morphine (1/6 to 1/8 gr.), given during the operation, are also a help in reducing the dose of chloroform.

Tracheotomy.—This is always necessary. The immemorial procedure is modified in three important ways. The thyroid isthmus is never drawn aside, but if met with is freely incised or excised; the trachea is not merely incised but a disc one-third of an inch in diameter is taken out of its anterior wall. Thus the patient should be able to breathe freely through the tracheotomy wound before the tube is inserted. This is the only certain way of preventing dangerous emergencies or even fatalities when the tube is accidentally dislodged or deliberately changed. Finally, StClair Thomson's form of Durham's tube in its largest size is always used and is indispensable. The tube will be necessary for about a week; it is removed as soon as the patient can breathe easily through the larynx while the tube is stopped. It is a small and obvious point that the tube, when it is to be discarded, should be taken out in the morning. Healing of a tracheotomy wound is usually quicker when a disc has been removed than when the trachea has been incised, for there is much less likely to be necrosis of the rings.

Gland Operation.—Only unilateral gland infection has to be provided against. When glands are palpably involved the usual complete unilateral dissection is done with removal of the jugular vein and ligature and division of the external carotid close to the bifurcation, but with preservation of the sterno-mastoid. The anterior edge of this muscle is then stitched to the prevertebral fascia behind the pharynx and the gland wound behind it drained by a separate puncture. If the glands are not obviously involved a rather limited gland dissection is done in the anterior triangle and the sterno-mastoid stitched over the vessels as mentioned above.

The Pharyngotomy.—A vertical incision is made over the middle of the lateral aspect of the larynx down to the cartilages and the infrahyoid muscles and constrictors are

turned forwards and backwards so that the great cornu of the hyoid and the thyroid ala are completely exposed. The structures on the deep surfaces of these are separated from them with a raspatory, and then the great cornu is removed to its joint, as well as the posterior two-thirds or more of the ala. The lax pharyngeal wall is now exposed, and the situation and extent of the growth are easily made out through it. The wound is carefully protected from the contact of pharyngeal mucus, and the pharynx is incised in such a way that this incision will serve in part for the isolation of the tumour. The removal of the latter with an adequate margin is then completed. The gap left is carefully closed by stitches, and in the same procedure the incision in the pharyngeal wall is stitched up. The constrictors and infrahyoid muscles are brought together closely over the line of suture in the pharynx. The neck wound is left widely open. It is filled with sterile boric powder and plugged over an apron of green protective. It should be the object and expectation of the surgeon to avoid altogether infection of the neck wound from the pharynx, both during the operation and after. A small rubber catheter is left in the pharynx and brought out through the mouth for feeding. It is left in for about five days and then passed for individual feeds.

In about a month healing should be complete, swallowing normal, and speech rapidly recovering its usual strength. This estimate applies only to favourable cases; there are, of course, many variations in the length of convalescence, and the patient may need skilled nursing and surgical care for some weeks more.

SOME FINAL CONSIDERATIONS.

When we turn to consider in general the results of the surgery of malignant disease of the laryngo-pharynx, it is but fair to remind ourselves in the first place of what the results of the disease are without sur-

gery. There is no more dreadful form of malignant disease, and we are to remember that the favourable easy types left untreated are spared nothing in the end. Again, we may well remind ourselves that the treatment we can use is no surgery of mutilation, but looks to leave the patient capable of leading a normal unmarked life. Such considerations may allow us to regard with a certain indulgence the large gaps between what we should like to do and what we can do in this department.

At the present time two conclusions seem to me to be fairly definite. In the first place we have technical methods which seem soundly established and capable of progressive refinement and extension. In the second place we can say that, in cases carefully but not very rigorously selected as favourable, we can look for satisfactory results as to the disease, secured by drastic but not very dangerous operations which leave no serious disability.

One of the accompaniments of progress in the surgery of any kind of malignant disease seems always to have been the disclosure of the extent to which late recurrences are possible. In my experience the surgery of the pharynx has followed this rule. I have, for example, comparatively recently seen two cases in which after seven years of complete freedom there has been a return or fresh outbreak of the disease. To my mind such events are not wholly discouraging, for they show how our efforts are helped by the natural resistive powers of the body, and even to put the case on the humblest level, health for seven years is no inconsiderable gift.

DESCRIPTION OF COLOURED PLATE.

FIG. 1.—Epithelioma of lateral wall; redrawn from a sketch of the laryngoscopic appearances. At first sight the growth seemed to have started in the pyriform sinus and the case appeared to be unfavourable. The arytenoid cartilage was, however, freely movable, and this suggested that the growth might in fact have begun on the lateral wall. At operation this was found to be the case, and a satisfactory local excision was possible without damage to the arytenoid. The case illustrates the deceptive appearances given to growths of the lateral wall by their mobility.

FIGS. 2, 3, AND 4.—Laryngoscopic appearances of post-cricoid carcinoma. These drawings are not intended to reproduce individual cases but to give a generalized idea of the three stages referred to in the text. In Fig. 2 the only abnormality is the pool of frothy mucus behind the arytenoids that indicates obstruction of the hypopharynx. In Fig. 3 oedema of the arytenoid region has appeared. In Fig. 4 the upper edge of an epitheliomatous ulcer is visible. In strong phonation this would show more extensively.

FIG. 5.—To illustrate the type of plastic procedure necessary in connexion with the more extensive operations. Drawing made some weeks after primary operation for a large growth. The excision involved removal of a large part of the pharynx and larynx, and, of course, of the party wall between them. Repair of the defect has been prepared for by the implantation of a large folded flap of skin presenting epithelial surfaces to the pharynx and larynx. At the second operation this flap will be used to re-form the upper laryngeal opening and the wall between larynx and pharynx, and the wound will be completely closed.

FIG. 1.

FIG. 2.

FIG. 3.

FIG. 4.

FIG. 5.